

Le Jardin de Mère Nature dans une Petite Planète

(Mother Nature's Garden on a Small Planet)

a SPECIES DISTRIBUTION LISTING for

TOWNSHIP 14 SOUTH, RANGE 09 EAST PIMA COUNTY, ARIZONA Gila and Salt River Baseline and Meridian

August 31, 2010 Update

“An increasing need for careful husbandry of the earth's natural resources has renewed interest in the classification and mapping of ecosystems. The inventory of our remaining biotic entities is particularly urgent because the increased aspirations of a constantly growing world population are placing ever greater stress on these generous, but finite, living resources.”

United States Department of Agriculture, Forest Service, General Technical Report RM-73



This photograph was taken looking east northeast toward the Roskruge Mountains.
William T. Kendall, March 9, 2007

“To know the desert involves an acquaintance with all its aspects, and all its physical features, as well as all of the animals and plants that have learned how to find in it a congenial place to live. The most significant lesson that the desert dweller can learn from a familiarity with its plant and animal life is to regard himself not as an exile from some better place, but as a man at home in an environment to which his life can be adjusted without physical or intellectual loss.”

Forest Shreve, *The Cactus and Its Home*, found in *Discovering the Desert*, by William G. McGinnies

MAJOR CONTRIBUTORS AND SOURCES OF INFORMATION

William T. Kendall *WTK (date of observation)*

Southwest Environmental Information Network (SEINet) *85 (a date of a search for information on the species)*

E. Lendell Cockrum. 1960. *The Recent Mammals of Arizona: Their Taxonomy and Distribution*, The University of Arizona Press, Tucson, Arizona. This inclusion is based on the general distribution maps and statements. *118 (distribution note, map - Figure Number and Page Number)*

Charles H. Lowe. 1964. *The Vertebrates of Arizona with Major Section on Arizona Habitats*, The University of Arizona Press, Tucson, Arizona. *55*

SPECIES DISTRIBUTION LISTINGS

Species Distribution Listings are being developed to encourage and promote the conservation of local native animals and plants. The listings are developed for legally defined geographic areas, and larger bodies of water. The listings include species reported as having been observed in or reported from the described area. Exotic and non-local landscaped plants are not included in the listings unless they have become naturalized into the surrounding native environment. Neither “Man” nor our domesticated animals, except for feral animals, have been included in the listings of species; however, they have had an impact on all natural areas, the future degree of this impact must be managed in order to restore and provide for the continuation of the natural interrelationships between all species.

Due to the continuing addition of species, the listings should be considered works in progress. In the listings, and most often in the listing of animals, species have been included based on general distribution mapping and/or statements and not on an observation made in a specific location. It is recommended that we consider a species as being “confirmed” to a township or general listing area only after we have at least three recorded sightings, cited in the footnotes, with no more than one of those records being based on general distribution mapping for the species. Note that the Southwest Environmental Information Network (SEINet) *85* may have several collections recorded for a species within any given township or listing area, and that the date shown in parentheses is a date of the search of their records and not a date of

recorded sighting. Note also that many of the individual species collection records found in SEINet include additional associated species. For assistance with the identification of a plant, contact the University of Arizona Herbarium (520-621-7243; FAX: 520-621-7186; P.O. Box 210036 Herring Hall, 1130 East South Campus Drive, Tucson, Arizona 85721).

Individual species records are presented alphabetically by division, class, family and genus within their kingdoms. Following the scientific name is the authority, common synonym(s), common name(s), a general description of the species, a general description of the habitat, the biotic communities in which it has reportedly been observed and footnotes. An attempt is being made to identify the range in mature (flowering/fruiting) heights reported for the plants. Wherever possible the flowering period is given as it has been reported and is inclusive to early month (1st-10th), mid-month (11th-20th) and late month (21st-end). The habitat description is provided in order to help you visualize the types of natural habitats a species is found in. Descriptions have been developed from and are based on herbarium records and general descriptions of habitat. The habitat description provided should not be considered as limiting as to the type of habitat that a plant might occupy. The terms “streambed”, “creekbed”, “riverbed” or “lakebed” refer to their dry aspects. Plants reported as occurring in recently burned areas were observed in the area within one year following a fire. The range in elevation has been rounded off to the nearest 100 feet up for the higher elevation, or down for the lower elevation. Species reported from within 0 to 100 feet as their lower elevation limit have been recorded as occurring “from sea level”. The reporting of the ecological formations follows the mapping presented in the “Biotic Communities of the Southwest” by David E. Brown and Charles H. Lowe, August 1980, with the exception of the “wetlands” which are being reported as an ecological formation in the listings. Species not considered to be native to Arizona are shown as being **EXOTIC**, printed in red. Exotic plants are not recommended for use in landscaping or restoration projects. Plants that may be an attractive component of a restored native habitat are so noted. Plants reported as having been used by native peoples of North America and which might be investigated to determine their value as a home garden or commercial food, fodder, beverage, spice, fiber, and/or dye crop may be so noted; much of this information is based on the records of the Native American Ethnobotany website [University of Michigan - Dearborn], footnote *127*. Species once reported as having occurred within the described area, but that no longer occurs there, may be shown as **EXTIRPATED**. Disjunct species, outliers and plants on the edge of the main population, as observed by the surveyor, may be noted as being **PERIPHERAL**. When describing the “native range” of plants in North America northwestern refers to Alaska, northern refers to northern Canada (the Yukon Territory, Northwest Territories and Nunavut), northeastern refers to Greenland, central refers to southern Canada (north-central: British Columbia, Alberta, Saskatchewan, Manitoba, Ontario, Quebec, Newfoundland and Labrador, New Brunswick, Nova Scotia and Prince Edward Island) and the United States (south-central), and southern refers to Mexico, below which is Central America and South America. In the footnotes, the source(s) used for the inclusion of the species in a listing is printed in either green *00* (indicating that the entry is based on an actual sighting) or blue *00* (indicating that the entry is based on a general distribution description and/or mapping). Plants listed in the book “Livestock-Poisoning Plants of Arizona” by Ervin M. Schmutz, Barry M. Freeman and Raymond E. Reed and published in 1968 (80) as being either “**Major Poisonous Range Plants**” or “**Secondary Poisonous Range Plants**” are further identified by their listing heading being printed in red in the footnotes; plants considered to be “Rarely Poisonous and Suspected Poisonous Range Plants” and “Poisonous Cropland and Garden Plants” have also been noted. In order to facilitate referencing to T.H. Kearney and R. H. Pebbles’ “Arizona Flora” the corresponding page number(s) for the species has been provided in the footnote (*46 (Page #)*).

Local native plants are recommended for use in landscape and restoration projects. Once established many native species require little, if any, irrigation. The inclusion of a plant in the township listing does not necessarily mean that the plant is suitable for the site in which you want to plant it. Ideally restoration should include those species of plants that were native to the property. The source material, of plants and seed, used in the project should be as local as possible. In order to determine what plants were native try

to locate photographs of the area prior to clearing or look for natural areas and remnant populations and plants adjacent to where the restoration is to take place. Plants should be planted in their approximate original habitat and density and taking into consideration the original local native site and elevation of occurrence.

The use of local native plants in landscape and restoration projects encourages native animals to remain in the area and helps us to retain the area's natural beauty and unique identity and heritage.

The Species Distribution Listings have been created and maintained by William T. Kendall. Any questions, concerns, corrections and/or comments, including the reporting of unrecorded species and information relating to historical distributions, may be sent to the following address: William T. Kendall, P.O. Box 86091, Tucson, Arizona 85754-6091

DISCLAIMER: The information presented under "Township Notes" has been obtained from large scale mapping and should be used only as a general guide. The listings are not meant to take the place of on-site surveys for species. Information used in the listings is accepted from biologists and individuals interested in helping to promote the conservation of our natural resources. Mistakes are made in the identification of species, the interpretation of data and in the recording of information, and changes in nomenclature occur. For these reasons I can not and do not warrant the accuracy of these listings. Attempts are made to keep the information contained in the Species Distribution Listings as accurate as possible; however, I disclaim any implied warranty or representation about its accuracy, completeness, or appropriateness for any particular purposes. Users of the information found in the listings assume full responsibility for their use of the information and understand that I not responsible or liable for any claim, loss, or damage resulting from its use.

CAUTION: Many native desert plants have sharp thorns and spines. Care should be given when handling these plants and consideration should be given to public safety at sites where they are to be planted. Range plants having a known toxic or poisonous property may be so noted. Footnotes for plants whose sources may have cautionary statements, comments and information on rarely poisonous or suspected poisonous properties may be shown in red *00*. Many poisonous plants are similar in appearance to edible ones. No field collected plant should be eaten unless you know for a fact that it is safe for you to do so.

CONTENTS

Township Notes

Conservation Related Organizations and Nurseries

Listing of Plants

Kingdom Plantae: The Plant Kingdom

Subkingdom Tracheobionta: The Vascular Plants

Superdivision Spermatophyta: The Seed Plants

Division Magnoliophyta: The Flowering Plants

Class Liliopsida: The Monocots

Class Magnoliopsida: The Dicots

Listing of Animals

Kingdom Animalia: The Animal Kingdom
Subkingdom Metazoa: The Multicellular Animals
Section Deuterostomia: The Deuterostomes
Phylum Chordata: The Chordates
Subphylum Vertebrata: The Vertebrates
Class Mammalia: The Mammals

Acknowledgements

Footnotes and References for the Species Distribution Listings

TOWNSHIP NOTES

LOCATION: This township is located in east-central Pima County in south-central Arizona. A portion of the Tohono O’Odham Nation is located within this township.

Historic Ranching Activities: General ranching activities included the placement of corrals, windmills and stock tanks. Named stock tanks: the Indio Tank, the West Tank.

Historic Mining Activities: General mining activities included prospecting.

LANDMARKS: A portion of the Roskrige Mountains is located in this township. Named peaks in the township include Pescadero Mountain (3,717 feet) and the north half of La Tortuga Butte (3,050 feet). This township is located in Aguirre Valley. Named washes include the Pescadero Wash and the Aguirre Wash.


ELEVATION: Elevations range from approximately 2,265 feet in a wash located on the north township line just west of the northeast corner to approximately 3,739 feet on an unnamed (Roskrige) peak in the Roskrige Mountains.

PHYSIOGRAPHIC PROVINCE: This township is located within the Sonoran Desert Section of the Basin and Range Physiographic Province (2).

SOILS: Soils have been described as being Hyperthermic (very hot) Arid Soils (soils with mean annual soil temperatures of more than 72 degrees Fahrenheit (22 degrees Centigrade) and less than 10 inches (25 cm) mean annual precipitation) of the Mohall-Laveen-Coolidge Association (deep soils on the valley plains and old terraces) and Thermic (hot) Arid Soils (soils with mean annual temperatures of 59 degrees to 72 degrees Fahrenheit (15 degrees to 22 degrees Centigrade) and 5 to 10 inches (13 to 25 cm) mean annual precipitation) and/or Thermic (hot) Semiarid Soils (soils with mean annual temperatures of 59 degrees to 72 degrees Fahrenheit (15 degrees to 22 degrees Centigrade) and 10 to 16 inches (25 to 41 cm) mean annual precipitation) of the Mohave-Tres Hermanos-Anway Association (deep, arid soils on the valley plains); Pinaleno-Nickel-Palos Verdes Association (deep, arid, gravelly soils on deeply dissected uplands), and the Rock Outcrop-Lampshire-Cellar Association (rock outcrop and very shallow and

shallow semiarid soils of the mountains and foothills) with isolated areas of Rock Outcrop and shallow soils (3).

BIOTIC COMMUNITY: Portions of this township are located within the Lower Colorado River and Arizona Upland Subdivisions of the Sonoran Desertscrub Regional Formation of the Desertscrub Formation with associated Wetlands (4).



Map of Township and Adjacent Sections

LISTED BELOW ARE A FEW OF THE NATIVE PLANTS REPORTED AS OCCURRING IN THIS TOWNSHIP THAT MIGHT BE CONSIDERED

FOR USE IN LANDSCAPE AND RESTORATION PROJECTS

Trees and Large Shrubs (over 7 feet maximum height)

Netleaf Hackberry (*Celtis laevigata* var. *reticulata* - 6' to 60')

Saguaro (*Carnegiea gigantea* - 5' to 60')

Velvet Mesquite (*Prosopis velutina* - 2' to 56')

Organ Pipe Cactus (*Stenocereus thurberi* - 5' to 40')

Blue Paloverde (*Parkinsonia florida* - 40" to 40')

Ocotillo (*Fouquieria splendens* - 5' to 33')

Foothill Paloverde (*Parkinsonia microphylla* - 40" to 26')

Catclaw Acacia (*Acacia greggii* var. *greggii* - 40" to 25')

Desert Hackberry (*Celtis ehrenbergiana* - 3' to 20')

Whitethorn Acacia (*Acacia constricta* var. *constricta* - 1' to 20')

Whitethorn Acacia (*Acacia constricta* var. *paucispina* - 1' to 20')

Chain-fruit Cholla (*Cylindropuntia fulgida* var. *fulgida* - 3' to 15')

Staghorn Cholla (*Cylindropuntia versicolor* - 3' to 15')

Greythorn (*Ziziphus obtusifolia* var. *canescens* - 3' to 13')

Kearney Snakewood (*Condalia warnockii* var. *kearneyana* - 20" to 13')

Creosote Bush (*Larrea tridentata* var. *tridentata* - 20" to 13')

Fishhook Barrel Cactus (*Ferocactus wislizeni* - 1' to 11')

Desert Broom (*Baccharis sarothroides* - 3' to 10')

Teddybear Cholla (*Cylindropuntia bigelovii* - 20" to 10')

Cane Cholla (*Cylindropuntia spinosior* - 16" to 10')

Santa Catalina Prairie Clover (*Dalea pulchra* - 20" to 8')

Desert Pricklypear Cactus (*Opuntia engelmannii* var. *engelmannii* - 20" to 8')

Vines and Climbers

Slender Janusia (*Janusia gracilis* - 16" to 10')

Shrubs (2 to 7 feet maximum height)

Limberbush (*Jatropha cardiophylla* - 1' to 7')

Tulip Pricklypear Cactus (*Opuntia phaeacantha* - 10" to 7')

Wright Beebrush (*Aloysia wrightii* - 20" to 78")

Desert Christmas Cactus (*Cylindropuntia leptocaulis* - 1' to 6')

Black-spined Pricklypear Cactus (*Opuntia macrocentra* var. *macrocentra* - 2' to 5')

Desert Mistletoe (*Phoradendron californicum* - 8" to 5', see note)

Fairyduster (*Calliandra eriophylla* - 4" to 5')

Triangleleaf Bursage (*Ambrosia deltoidea* - 1' to 4')

Grasses

False Rhodes Grass (*Trichloris crinita* - 24" to 60")

Tanglehead (*Heteropogon contortus* - 8" to 60")

Whiplash Pappusgrass (*Pappophorum vaginatum* - 16" to 52")

Plains Bristlegrass (*Setaria vulpiseta* - 12" to 48")
Sand Dropseed (*Sporobolus cryptandrus* - 12" to 48")
Bush Muhly (*Muhlenbergia porteri* - 10" to 44")
Tobasa (*Pleuraphis mutica* - 12" to 36")
Black Grama (*Bouteloua eriopoda* - 8" to 30")
Desert Fluffgrass (*Dasyochloa pulchella* - ½" to 6")

Shrubs (under 2 feet maximum height), Subshrubs, Herbs and Small Succulents

Brownfoot (*Acourtia wrightii* - 1' to 5')
Desert Tobacco (*Nicotiana obtusifolia* var. *obtusifolia* - 12" to 42", see note)
Bundle Hedgehog Cactus (*Echinocereus fasciculatus* - 2" to 18")
Graham Pincushion Cactus (*Mammillaria grahamii* - 1" to 12")
Desert Holly (*Acourtia nana* - 2" to 10")

CONSERVATION RELATED ORGANIZATIONS AND NURSERIES

Arizona Department of Agriculture

<http://www.azda.gov/>

Native Plant Crimes HOTLINE: 602-364-0907

The mission statement of the Arizona Department of Agriculture is to regulate and support Arizona agriculture in a manner that encourages farming, ranching, and agribusiness while protecting consumers and natural resources.

NOTICE OF INTENT TO CLEAR LAND

The Arizona Department of Agriculture enforces the sections of the Arizona Revised Statutes commonly referred to as the "Arizona Native Plant Law". The statutes require, in part, that anyone who is clearing land notify the State of Arizona in advance of the clearing. Some land owners involved in the clearing of land allow for nurseries and people who are interested in salvaging plants to do so prior to the clearing. The Arizona Department of Agriculture posts these notifications in their county offices. You may also contact the Arizona Department of Agriculture and, for a fee, be put on a mailing list of people receiving copies of the Notices of Intent to Clear Land.

Contact Information: Arizona Department of Agriculture, 1688 West Adams Street, Phoenix, Arizona 85007. Telephone number: 602-542-4373.

Arizona Game and Fish Department

<http://www.gf.state.az.us/>

Operation GAME THIEF: 602-942-3000

~~The mission statement of the Arizona Game and Fish Department is to conserve, enhance, and restore Arizona's diverse wildlife resources and habitats through aggressive protection and management programs, and to provide wildlife resources and safe watercraft and off-highway vehicle recreation for the enjoyment, appreciation, and use by present and future generations.~~

As part of their conservation program the Arizona Game and Fish Department provides ideas on how to learn to live with, and landscape for, wildlife:

LIVING WITH WILDLIFE

http://www.azgfd.gov/w_c/urban_wildlife.shtml

Contact Information: Arizona Game and Fish Department, 5000 West Carefree Highway, Phoenix, Arizona 85086-5000. Telephone number: 602-942-3000

Arizona Native Plant Society

<http://aznps.org/>

The Arizona Native Plant Society is a statewide nonprofit organization devoted to Arizona's native plants. Its mission is to promote knowledge, appreciation, conservation, and restoration of Arizona native plants and their habitats. They work with the Southwest Rare Plant Task Force to develop strategies for protecting rare species and their habitats; they keep abreast of conservation issues concerning native plants species and responds to those through their Conservation Committee; they promote the use of native species in residential and commercial landscapes; they publish the Plant Press, support the publication of scholarly works and maintains a website with information and links about native plant, and they host a series of statewide events that provide forums to learn from professionals. Member activities and benefits include chapter and statewide gatherings; field trips and educational presentations; conservation through education, outreach and restoration; habitat restoration projects; informative website, newsletters and journals, and interactions with plant experts and enthusiasts.

LISTING OF SOURCES FOR NATIVE PLANTS AND SEEDS

The Arizona Native Plant Society maintains a listing of Native Plant and Seed Sources at:

<http://www.aznps.org/sources.html>

Contact Information: Arizona Native Plant Society, PO Box 41206, Tucson, Arizona 85717.

Tucson Cactus and Succulent Society

<http://www.tucsoncactus.org/>

The Tucson Cactus and Succulent Society is a non-profit organization dedicated to educating, teaching and learning about cacti and succulent plants. Their monthly programs feature knowledgeable individuals who can educate you and help you understand more about these fascinating plants. They conduct and sponsor native cactus and succulent rescue operations, plant sales, field trips, nursery and garden visits, conventions and conferences as well as other activities throughout the year.

NATIVE PLANT RESCUE NOTICE

The Tucson Cactus and Succulent Society expends a tremendous amount of time and effort in the organizing and overseeing of their native plant rescue events. The native plant rescues carried out by the dedicated members of the Society provide an immeasurable service to our community.

Members of the Tucson Cactus and Succulent Society organize native plant rescues in areas being cleared for development. If interested in rescuing plants and/or obtaining local native plants for your landscaping or restoration project join the Society and become a rescue crew member.

Contact Information: Tucson Cactus and Succulent Society, PO Box 64759, Tucson, Arizona 85728-4759. Telephone number: 520-885-6367.

Desert Survivors Native Plant Nursery

<http://www.desertsurvivors.org/nursery.asp>

The Desert Survivors Native Plant Nursery sells many local native plants and is willing to consider growing any native plant for which there is a buyer.

Contact Information: Desert Survivors Native Plant Nursery, 1020 West Starr Pass Boulevard, Tucson, Arizona 85713. Telephone number: 520-791-9309.

Native Seeds/SEARCH

<http://www.nativeseeds.org>

The Native Seeds/SEARCH is a nonprofit conservation organization that seeks to preserve the crop seeds that connect the Native American cultures to their lands. The mission of the Native Seeds/SEARCH is to conserve, distribute and document the adapted and diverse varieties of agricultural seeds, their wild relatives and the role these seeds play in the cultures of the American Southwest and Northwest Mexico.

Contact Information: Native Seeds/SEARCH, 526 North Fourth Avenue, Tucson, Arizona 85705. Telephone number: 520-622-5561 or toll free at 866-622-5561; FAX 520-622-5561; e-mail: info@nativeseeds.org

LISTING OF PLANTS

**STRICTLY ENFORCED LAWS PROTECT MANY OF ARIZONA'S NATIVE
PLANTS FROM COLLECTION, MUTILATION AND DESTRUCTION**

Native Plant Crimes HOTLINE: 602-364-0907

Kingdom Plantae: The Plant Kingdom
Subkingdom Tracheobionta: The Vascular Plants
Division Magnoliophyta: The Flowering Plants

CLASS LILIOPSIDA: The MONOCOTS

Poaceae (Gramineae): The Grass Family

Andropogon contortus (see *Heteropogon contortus*)

***Bouteloua eriopoda* (J. Torrey) J. Torrey: Black Grama**

COMMON NAMES: Black Grama, Black Grass, Hairyfoot Grama, Navajita Negra (Spanish), Woolly Foot Grama, Woollyfoot Grama. DESCRIPTION: Terrestrial perennial graminoid (a bunchgrass with decumbent culms 8 to 30 inches in height, plants were described that were 6 to 8 inches in height and 2 to 4 inches in width, plants 8 inches in height and 2 to 4 inches in width were observed and recoded); the foliage is grayish-green curing to gray; the flowers (spikelets) may be maroon, purplish, reddish-brown or red-maroon; the anthers are orange to yellow; flowering generally takes place between late June and early November (high amounts of spring rainfall may induce an earlier flowering; additional records: four for late May). HABITAT: Within the range of this species it has been reported from mountains; gravelly mountaintops; mountainsides; bouldery, rocky, gravelly, gravelly-sandy, sandy and sandy-silty mesas; sandy plateaus; rims of gorges; rocky canyons; rocky canyon sides; bedrock, bouldery and sandy canyon bottoms; gorges; rocky talus slopes; crevices in rocks; pockets of sandy soil; rocky knolls; ledges; gravelly ridges; gravelly-loamy ridgetops; sandy meadows; cinder cones; crater walls; rocky foothills; bouldery, rocky, rocky-gravelly, rocky-sandy and sandy hills; gravelly-clayey hilltops; bouldery and rocky hillsides; sandy bases of escarpments; bouldery, bouldery-gravelly, bouldery-sandy, rocky, rocky-gravelly, rocky-loamy, rocky-clayey, shaley, stony, cobbly, cindery, gravelly, gravelly-sandy, gravelly-sandy-clayey, gravelly-loamy, gravelly-clayey, sandy, sandy-loamy and clayey-loamy slopes; gravelly-sandy-loamy alluvial fans; rocky outcrops; amongst boulders, rocks and stones; rims of craters; lava flows; sandy sandhills; sandy dunes; sandy hummocks; sandy prairies; sandy, sandy-clayey and clayey-loamy plains; stony-sandy, cindery and sandy-loamy flats; esplanades; basins; valley floors; sandy valley bottoms; along rocky, rocky-gravelly, rocky-sandy, gravelly, gravelly-sandy, gravelly-loamy, sandy and loamy roadsides; sandy arroyos; bottoms of arroyos; draws; rills; along creeks; along and in bouldery, rocky, gravelly, sandy and clayey washes; within bouldery-gravelly, rocky and gravelly drainages; bedrock depressions; swales; sandy beaches; gravelly-sandy benches; gravelly shelves; gravelly terraces; floodplains, and waste places growing in dry, well drained desert pavement; bouldery, bouldery-gravelly, bouldery-sandy, rocky, rocky-gravelly, rocky-gravelly-sandy, rocky-sandy, shaley, stony, stony-sandy, cobbly, cindery, gravelly, gravelly-sandy and sandy ground; rocky loam, cobbly-sandy loam, gravelly loam, gravelly-sandy loam, sandy loam, sandy-clayey loam, sandy-silty loam, clayey loam, silty loam and loam ground; rocky-clayey, gravelly clay, gravelly-sandy clay, sandy clay and clay ground, and sandy silty ground, occurring from 1,800 to over 8,200 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, established plants are tolerant of short droughts and the plants may form "fairy rings". The plants may form large patches through the spreading and rooting of stoloniferous runners, it also regenerates through tillering and layering. Even though it may be difficult to establish, consideration should be given to including this plant in re-seeding mixtures, seedling establishment is reduced by drought. Black Grama is considered excellent forage for wildlife, and Cactus Wrens (*Campylorhynchus brunneicapillus*) will use this grass in the construction of their nests. *Bouteloua eriopoda* is native to

southwest-central and southern North America. *5, 6, 15, 30, 33 (Pages 149-151), 43 (081910), 46 (Page 128), 48, 58, 63 (081910 - color presentation), 77, **85** (081910 - color presentation), 105*

Chloris crinita (see *Trichloris crinita*)

***Dasyochloa pulchella* (K.S. Kunth) C.L. von Willdenow ex P.A. Rydberg: Low Woollygrass**

SYNONYMY: *Erioneuron pulchellum* (K.S. Kunth) T. Tateoka, *Tridens pulchellus* (K.S. Kunth) A.S. Hitchcock, *Triodia pulchella* K.S. Kunth. COMMON NAMES: Desert Fluffgrass, Fluff Grass, Fluff-grass, Fluffgrass, Low Woollygrass, Oerennuak Grass, Zacate Borreguero. DESCRIPTION: Terrestrial perennial (often appearing to be an annual and has also been described as being a short-lived perennial) tufted graminoid (a bunchgrass (clumpgrass) ½ to 6 inches in height, plants were observed that were 2 to 4 inches in height and 2 to 4 inches in width, plants were observed that were 4 inches in height and 12 inches in width); the foliage is bluish-green curing to a gray-white; the flowers are green, silvery or white; flowering generally takes place between late March and late October (additional record: one for early December). HABITAT: Within the range of this species it has been reported from mountains; rocky-sandy, gravelly, sandy-loamy and clayey mesas; rocky, gravelly and sandy canyons; gravelly-sandy canyon bottoms; rocky talus slopes; sandy soils in crevices in rocks and rock slabs; knolls; rocky and gravelly ridges; clayey ridgetops; ridgelines; meadows; foothills; rocky, gravelly and sandy hills; rocky, rocky-sandy and gravelly hillsides; sandy bases of escarpments; sandy edges of escarpments; bouldery, rocky, rocky-gravelly, stony, cindery-clayey, gravelly, gravelly-sandy, gravelly-sandy-loamy, gravelly-sandy-clayey-loamy and sandy slopes; rocky alluvial fans; rocky-sandy, gravelly and sandy bajadas; rocky outcrops; amongst boulders and rocks; rocky-sandy coves; lava rincons; sand hills; sand dunes; breaks; gravelly steppes; sandy and clayey plains; rocky, cindery, gravelly, gravelly-sandy, sandy, sandy-loamy and clayey flats; valley floors; along railroad right-of-ways; along bouldery-rocky, rocky, gravelly, gravelly-sandy-loamy, gravelly-loamy, sandy and sandy-loamy roadsides; arroyos; sandy bottoms of arroyos; gravelly draws; rocky gullies; streambeds; creekbeds; along and in rocky-sandy, gravelly, gravelly-sandy and sandy washes; along and in sandy drainages; playas; marshes; clayey depressions; along banks of washes; edges of washes; rocky-sandy shores of lakes; benches; gravelly and sandy terraces; rocky-sandy and loamy bottomlands; floodplains; rocky lowlands; sandy riparian areas, and disturbed areas growing in moist or dry desert pavement; bouldery, bouldery-rocky, rocky, rocky-gravelly, rocky-sandy, stony, cobbly, cindery, gravelly, gravelly-sandy and sandy ground; gravelly loam, gravelly-sandy loam, gravelly-sandy-clayey loam, sandy loam, clayey loam and loam ground; rocky clay, cindery clay, gravelly-sandy clay and clay ground, and sandy silty ground, occurring from 100 to 7,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This low, densely tufted perennial may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America; it was noted as having been used as a drug or medication. This plant is browsed by the Desert Bighorn Sheep (*Ovis canadensis mexicana*); however, it has been reported that this plant is generally avoided by grazing animals. *Dasyochloa pulchella* is native to southwest-central and southern North America. *5, 6, 15 (recorded as *Erioneuron pulchellum* (H.B.K.) Tateoka), 16 (recorded as *Erioneuron pulchellum* (H.B.K.) Tateoka), 33 (recorded as *Tridens pulchellus* (H.B.K.) Hitchc., Page 97), 43 (071309), 46 (recorded as *Tridens pulchellus* (H.B.K.) Hitchc., Page 90), 58 (recorded as *Erioneuron pulchellum* (H.B.K.) Tateoka), 63 (100609 - color presentation), 77 (recorded as *Erioneuron pulchellum* (H.B.K.) Tateoka), 85 (100609 - color presentation of dried material), 105 (recorded as *Tridens pulchellus* (H.B.K.) Hitchc.), 127, **WTK** (March 9, 2007)*

***Eragrostis cilianensis* (C. Allioni) F. Vignolo-Lutati ex E.E. Janchen: Stinkgrass**

SYNONYMY: *Eragrostis megastachya* (G.L. Koeler) J.H. Link. COMMON NAMES: Amoresco (Hispanic), Candy Grass, Candy-grass, Candygrass, Éragrostide Fétide, Graminha (Portuguese), Großes Liebesgras (German), Gray Love Grass, Lovegrass, Stink Grass, Stinkgrass, Stinking Lovegrass, Strong-scented Lovegrass, Strongscented Lovegrass, Watergrass, Zacate Apestoso (Hispanic), Zacate Apestoso (Hispanic), Zacate Borreguero (Hispanic), Zacate de Amor Hediondo

(Hispanic), Zacate Estepario (Hispanic). DESCRIPTION: Terrestrial annual tufted graminoid (a bunchgrass with prostrate to erect culms 3 to 36 inches in height); the foliage is gray-green or light green; the spikelets (flowers) are greenish, white or whitish with green veins turning tawny with age, the anthers are yellow; flowering generally takes place between late July and late October (additional records: one for mid-March, one for late March, one for mid-May, one for late May, one for early June, three for early July, one for mid-November, one for late November and one for mid-December). HABITAT: Within the range of this species it has been reported from mountains; sandy mesas; gravelly and sandy canyons; sandy canyon bottoms; sandy bases of cliffs; bluffs; knolls; ledges; sandy meadows; bouldery foothills; rocky hills; rocky and gravelly hillsides; rocky, rocky-gravelly, gravelly, gravelly-sandy-loamy, gravelly-loamy, sandy-loamy, sandy-clayey, loamy, clayey-loamy and clayey slopes; rocky outcrops; amongst rocks; coves; plains; sandy and sandy-silty flats; valley floors; valley bottoms; along gravelly, gravelly-loamy and sandy roadsides; arroyos; bottoms of arroyos; draws; gulches; gullies; gravelly-sandy seeps; springs; along streams; along and in cobbly-sandy streambeds; along creeks; creekbeds; sandy riverbeds; along and in gravelly, sandy and silty-clayey washes; within drainage ways; clayey lakebeds; cienegas; marshes; bedrock depressions; silty swales; along rocky-sandy, sandy and sandy-loamy banks of creeks, rivers, washes and drainages; sandy edges of streams, ponds, lakes and marshes; sand bars; sandy benches; gravelly-loamy terraces; loamy bottomlands; sandy and clayey floodplains; sandy mesquite bosques; around and in stock tanks (charcos, repressos); sandy ditches; sandy riparian areas; waste places, and disturbed areas growing in wet, moist, damp and dry bouldery, rocky, rocky-gravelly, stony, cobbly-sandy, gravelly, gravelly-sandy and sandy ground; rocky-clayey loam, gravelly loam, gravelly-sandy loam, sandy loam, clayey loam, clayey-humusy loam and loam ground; sandy clay, silty clay and clay ground, and sandy silty and silty ground, occurring from sea level to 8,000 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: **EXOTIC** Invasive Plant. *Eragrostis cilianensis* is native to middle, eastern and southern Europe; western, eastern and southern Asia, and Africa. *5, 6, 15, 16, 30, 33 (recorded as *Eragrostis megastachya* (Koel.) Link, Pages 82-83), 43 (101009), 46 (Page 86), 58, 63 (101009 - color presentation of seed), **68**, **77**, **80** (This plant is listed as a Rarely Poisonous and Suspected Poisonous Range Plant. "This annual grass was reported to cause poisoning in horses when fed in large quantities over a long period of time."), **85** (101009 - color presentation of sried material), 101 (color photograph), 105 (recorded as *Eragrostis megastachya* (Koel.) Link)*

Eragrostis megastachya (see *Eragrostis cilianensis*)

Erioneuron pulchellum (see *Dasyochloa pulchella*)

***Heteropogon contortus* (C. Linnaeus) A.M. Palisot de Beauvois ex J.J. Roemer & J.A. Schultes:**
Tanglehead

SYNONYMY: *Andropogon contortus* C. Linnaeus. COMMON NAMES: Barba Negra, Black Spear Grass, Bunch Spear Grass, Common Tangleweed, Hierba Torcida (Spanish), Piligrass (Hawaii), Tangle Grass, Tangle-head, Tanglehead, Tanglehead Grass, Retorcido Moreno, Spear Grass, Speergras (German), Zacate Colorado. DESCRIPTION: Terrestrial perennial tufted graminoid (a bunchgrass (clumpgrass) 8 to 60 inches in height); the foliage is bright green curing to orange-brown; the spikelets (flowers) may be brown or purple; flowering generally takes place between early January and late May and again between early August and early December (flowering records: one for early January, one for late January, one for late February, one for mid-March, one for early May, one for late May, three for early August, five for late August, three for early September, four for mid-September, three for late September, one for mid-October, four for early November, one for mid-November and one for early December). HABITAT: Within the range of this species it has been reported from mountains; rocky mountainsides; mesas; cliffs; cliff faces; along rocky canyons; along canyon walls; along bouldery and gravelly canyon bottoms; bases of cliffs; crevices in rocks; ledges; rocky ridges; bouldery ridgetops; foothills; rocky hills; rocky and gravelly-clayey hillsides; rocky, gravelly, gravelly-sandy and sandy

slopes; rocky outcrops; amongst boulders and rocks; lava flows; rocky and sandy plains; gravelly flats; valley floors; along sandy roadsides; along and in rocky arroyos; rocky-sandy bottoms of arroyos; along draws; within gullies; ravines; around seeping streams; streambeds; creekbeds; along and in rocky, rocky-sandy, cobbly, gravelly-sandy and sandy washes; within gravelly-sandy-loamy drainages; within rocky and sandy drainage ways; bedrock tinajas; around pools; (silty) banks of streams and rainwater basins; edges of washes; sandy beaches; terraces; floodplains; riparian areas, and disturbed areas growing in dry bouldery, bouldery-sandy, rocky, rocky-sandy, cobbly, gravelly, gravelly-sandy and sandy ground; gravelly-sandy loam and loam ground; gravelly clay ground, and silty ground, occurring from sea level to 7,500 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. *Heteropogon contortus* is native to south-central and southern North America and possibly to other tropic sub-tropic and warm-temperate regions of the world. *5, 6, 15, 16, 33 (Page 302), 43 (101209), 46 (Page 144), 48, 58, 63 (101209 - color presentation), 77, 85 (101209 - color presentation), 105 (Reports that Tanglehead Grass “is one of the easiest grasses to establish under conditions of low rainfall.”), **WTK** (March 9, 2007)*

Hilaria mutica (see *Pleuraphis mutica*)

***Muhlenbergia porteri* F.L. Scribner ex W.J. Beal: Bush Muhly**

COMMON NAMES: Bakú (Tarahumara), Bush-grass, Bush Muhly, Hoe Grass, Hoegrass, Liendrilla Amacollada (Hispanic), Mesquite Grass, Mesquitegrass, Porter’s Muhlenbergia, Telaraña (Hispanic), Zacate Aparejo (Hispanic). DESCRIPTION: Terrestrial perennial tufted graminoid (a bunchgrass with geniculate culms 10 to 44 inches in height/length and 18 inches to 10 feet in width, several plants were described as being 3 feet in height and 10 feet in width); the stems are dull green; the leaves are green, purplish-green or yellow-green curing to buff; the panicles (compound inflorescences) are usually purple; the spikelets (flowers) are green becoming purple when mature; anthers are purple to yellow; flowering generally takes place between late February and late October (additional records: one for late November and one for early December); the caryopsis (fruit) is yellowish-brown the aggregate of which covers the plants in a misty shroud. HABITAT: Within the range of this species it has been reported from mountains; rocky and stony-sandy mountainsides; mesas; rocky cliffs; bouldery and rocky canyons; rocky canyonsides; rocky-sandy and gravelly canyon bottoms; gorges; talus slopes; crevices in rocks; buttes; along sandy-silty and silty ledges; rocky ridge tops; foothills; rocky and sandy hills; bouldery-sandy and rocky hillsides; rocky escarpments; along bouldery, bouldery-rocky, rocky, rocky-loamy, gravelly, gravelly-loamy, sandy and sandy-loamy slopes; bajadas; rocky outcrops; amongst boulders and rocks; alcoves; sandy lava flows; lava fields; sand dunes; dune-like areas of fine blow-sand deposits; gravelly plains; rocky, gravelly-sandy, sandy and sandy loamy flats; open sandy ground amongst Ephedra and Larrea; basins; sandy valley floors; valley bottoms; along rocky, rocky-gravelly, gravelly, gravelly-loamy and sandy roadsides; rocky arroyos; clefts in rocky hillsides; within draws; gulches; ravines; springs; bouldery streambeds; along rivers; along and in rocky, rocky-gravelly, rocky-sandy, gravelly, gravelly-sandy and sandy washes; along drainages; bouldery-cobbly and rocky drainage ways; around ponds; gravelly-sandy banks; margins of washes; sandy-silty and silty benches; gravelly terraces; sandy floodplains; sandy mesquite bosques; riparian areas, and disturbed areas often growing in the protection of shrubs and trees in damp and dry rocky desert pavement; bouldery, bouldery-rocky, bouldery-cobbly, bouldery-sandy, rocky, rocky-gravelly, rocky-sandy, stony-sandy, cindery, gravelly, gravelly-sandy and sandy ground; rocky loam, rocky-clayey loam, cobbly-sandy loam, gravelly loam, sandy loam, clayey loam and loam ground; gravelly clay, sandy clay and clay ground, and cobbly-sandy silty, sandy silty and silty ground, occurring from 700 to 7,500 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. In areas where it occurs naturally, consider including Bush Muhly seed in reseeding mixtures. According to the USDA Forest Service Fire Effects Information System, Bush Muhly germinates best when temperatures are at 86 degrees Fahrenheit (30 degrees Centigrade). When re-vegetating desert washes consider planting Bush Muhly along with Whitethorn Acacia (*Acacia*

constricta), Catclaw Acacia (*Acacia greggii* var. *greggii*), Limberbush (*Jatropha cardiophylla*), Triangleleaf Bursage (*Ambrosia deltoidea*) and White Bursage (*Ambrosia dumosa*). Bush Muhly is browsed by the Desert Bighorn Sheep (*Ovis canadensis* subsp. *mexicana*). *Muhlenbergia porteri* is native to southwest-central and southern North America. *5, 6, 15, 16, 30, 33 (“Bush Muhly originally existed in extensive stands on the open range lands of southern Arizona but now occurs for the most part in the protection of shrubs and subshrubs and is seldom locally abundant. It is highly palatable and well liked by livestock despite the wiry culms.”, Pages 201-202), 43 (101709), 46 (Page 111), 48, 58, 63 (101709 - color presentation), 77, 85 (101709 - color presentation), 105 (“This was formerly one of the most abundant and important grasses of southern Arizona, but is found now largely as individual plants under the protection of shrubs. ... Where possible this grass should be allowed to set a full crop of seed during the summer growing season at least every second or third year. Deferment of grazing during July and August every year is recommended on run-down ranges.”), **WTK** (March 9, 2007)*

Panicum capillare var. *hirticaule* (see *Panicum hirticaule* var. *hirticaule*)

Panicum capillare var. *pampinosum* (see *Panicum hirticaule* var. *hirticaule*)

***Panicum hirticaule* C.B. Presl var. *hirticaule*: Mexican Panicgrass**

SYNONYMY: *Panicum capillare* C. Linnaeus var. *hirticaule* (J.S. Presl) F.W. Gould, *Panicum capillare* C. Linnaeus var. *pampinosum* (A.S. Hitchcock & M.A. Chase) F.W. Gould, *Panicum pampinosum* A.S. Hitchcock & M.A. Chase, *Panicum sonorum* W.J. Beal. COMMON NAMES: Chiri Chiri (Spanish), Mexican Panicgrass, Roughstalk Witchgrass, Sauhui (Spanish), Sonora Panic, Sowi Millet, Triguillo (Spanish), Witchgrass, Zacate de Año (Spanish), Zacate Peludo Perdís (Spanish). DESCRIPTION: Terrestrial annual graminoid (erect-spreading culms 2 to 40 inches in height); flowering generally takes place between mid-August and mid-October (flowering beginning as early as July has been reported). HABITAT: Within the range of this species it has been reported from mountains; sandy mesas; canyons; gravelly, gravelly-sandy and sandy canyon bottoms; rocky ledges; openings in woodlands; meadows; rocky hills; rocky, rocky-clayey and gravelly hillsides; rocky and gravelly slopes; bajadas; amongst boulders; dunes; sandy plains; clayey flats; basins; valley floors; along rocky-loamy roadsides; sandy draws; along seeps; along streams; streambeds; along and in oases; gravelly, gravelly-sandy, sandy and silty washes; within sandy drainage ways; sandy-silty depressions; clayey-loamy and silty swales; along margins of washes; along gravelly-sandy floodplains; mesquite bosques; ditches; sandy riparian areas, and disturbed areas growing in wet, moist and dry bouldery, rocky, gravelly, gravelly-sandy and sandy ground; rocky loam, gravelly loam and clay loam ground; rocky clay and gravelly clay and clay ground, and sandy silty and silty ground, occurring from sea level to 6,800 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: The species, *Panicum hirticaule*, was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. *Panicum hirticaule* var. *hirticaule* is native to southwest-central and southern North America; Central America, and South America. *5, 6, 30 (species), 33 (recorded as *Panicum capillare* L. var. *hirticaule* (Presl) Gould, Page 283; *Panicum capillare* L. var. *pampinosum* (Hitchc. & Chase) Gould, Page 284, and *Panicum sonorum* Beal, Page 282), 43 (101809), 46 (recorded as *Panicum pampinosum* (A.S. Hitchcock & M.A. Chase, Page 136), 63 (101809 - *Panicum hirticaule* J. Presl var. *hirticaule*), 77, **80** (Species of the genus *Panicum* are listed as Rarely Poisonous and Suspected Poisonous Range Plants. Species of this genus have been reported to cause loss in livestock due to photosensitization and nitrate poisoning.), **85** (101809 - *Panicum hirticaule* var. *hirticaule* J. Presl), 127*

Panicum pampinosum (see *Panicum hirticaule* var. *hirticaule*)

Panicum sonorum (see *Panicum hirticaule* var. *hirticaule*)

Pappophorum apertum (see *Pappophorum vaginatum*)

Pappophorum mucronulatum (see *Pappophorum vaginatum*)

***Pappophorum vaginatum* S.B. Buckley: Whiplash Pappusgrass**

SYNONYMY: *Pappophorum apertum* W. Munro ex F. Lamson-Scribner, *Pappophorum mucronulatum* auct. non C.G. Nees von Esenbeck. COMMON NAMES: Mucronulate Pappusgrass, Pappusgrass, Pima Pappusgrass, Whiplash Pappusgrass. DESCRIPTION: Terrestrial perennial graminoid (a bunchgrass (clumpgrass) 16 to 52 inches in height); the foliage is gray-green or light green; the inflorescences may be tinged with purple; based on few flowering records available, flowering generally takes place between late March and late October (flowering records: two for late March, one for late April, one for early July, one for late August, three for early September, one for mid-September and one for late October). HABITAT: Within the range of this species it has been reported from mountains; mountaintops; mesas; rocky canyons; bases of cliffs; ridgetops; foothills; rocky hillsides; rocky, sandy and clayey slopes; bajadas; rocky plains; gravelly and sandy-silty flats; basins; valley floors; valley bottoms; coastal dunes; sandy coastal flats; along railroad right-of-ways; along stony and sandy roadsides, along sandy gullies; along creeks; along and in gravelly washes; along drainage ways; depressions; banks of washes; along edges of washes; margins of washes; floodplains; dams; in sandy ditches, and disturbed areas growing in moist and dry rocky, stony, gravelly and sandy ground; clayey loam ground, clay ground, and sandy silty ground, occurring from sea level to 4,800 feet in elevation in the grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. *Pappophorum vaginatum* is native to southwest-central and southern North America and southern South America. *5, 6, 15, 16, 33 (recorded as *Pappophorum mucronulatum* Nees., Page 103), 43 (101809), 46 (recorded as *Pappophorum mucronulatum* Nees., Page 91), 48, 63 (101809 - color presentation), 77, **85** (101809 - color presentation of dried material), 105 (recorded as *Pappophorum mucronulatum* Nees.)*

***Pleuraphis mutica* S.B. Buckley: Tobosagrass**

SYNONYMY: *Hilaria mutica* (S.B. Buckley) G. Benth. COMMON NAMES: Tobosa, Tobosa Grass, Tobosagrass. DESCRIPTION: Terrestrial perennial graminoid (a bunchgrass 1 to 3 feet in height); the foliage is dull bluish-green or gray-green curing to gray; the inflorescence is purplish, straw or white; the spikelets (flowers) are greenish-tan or tinged with pink; flowering generally takes place between May and October; however, under favorable conditions, flowering may take place throughout the year (flowering records: one for early April, one for mid-April, one for early July, three for mid-August, one for late August, three for mid-September, one for mid-October, one for late October and two for early November). HABITAT: Within the range of this species it has been reported from mountains; mountaintops; bouldery and gravelly-sandy mesas; canyons; buttes; gravelly-sandy-clayey ridges; ridgetops; foothills; rocky and sandy-loamy hills; rocky hilltops; rocky and sandy hillsides; across bouldery, bouldery-rocky-clayey, rocky, rocky-clayey, cobbly-clayey, gravelly-sandy-clayey, sandy, sandy-loamy and clayey slopes; alluvial fans; bajadas; bouldery and rocky outcrops; amongst boulders and rocky-gravels; lava hills; lava fields; prairies; gravelly plains; sandy, sandy-clayey, sandy-silty, loamy and clayey flats; basins; rocky and sandy valley floors; valley bottoms; along gravelly-sandy roadsides; arroyos; gullies; along creeks; along and in bedrock, rocky and sandy washes; drainages; along drainage ways; clayey depressions; swales; along margins of washes; benches; terraces; floodplains; lowlands; mesquite bosques, and ditches growing in dry rocky desert pavement; bouldery, rocky, rocky-gravelly, rocky-sandy, gravelly, gravelly-sandy and sandy ground, rocky-clayey loam, gravelly-clayey loam, sandy loam, sandy-clayey loam, clayey loam and loam ground; bouldery-rocky clay, rocky clay, cobbly clay, gravelly-sandy clay, sandy clay, silty clay and clay ground, and rocky-gravelly silty and silty ground, occurring from 1,100 to 6,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. *Pleuraphis mutica* is native to southwest-central and southern North America. *5, 6, 16 (recorded as

Hilaria mutica (Buckl.) Benth.), 33 (recorded as *Hilaria mutica* (Buckl.) Benth., Page 161), 43 (102009, *Hilaria mutica* Benth.), 46 (recorded as *Hilaria mutica* (Buckl.) Benth., Page 122), 48, 63 (102009 - color presentation), 77 (recorded as *Hilaria mutica* (Buckl.) Benth.), 80 (The Ergot Fungus (*Claviceps* sp.) is listed as a Secondary Poisonous Range Plant. Tobosa (*Hilaria mutica*) can be a host of the Ergot Fungus. “Ergot contains poisonous alkaloids and other compounds that may cause chronic poisoning (gangrenous ergotism) in the extremities when consumed in small amounts, or convulsive poisoning when large amounts are eaten. Animals may be poisoned by feeding on mature, infected grain or hay. Livestock, especially cattle, and humans are susceptible. ... Pastures causing ergot poisoning should be mowed or the animals removed. Mildly poisoned animals will usually recover if removed from the infested pastures, kept quiet, and supplied with good feed and water. In Arizona, some losses may be expected on rangelands during wet years, but most losses have occurred from grazing pastures of Dallas Grass (*Paspalum dilatatum*.” See text for additional information. Tobosa is listed as a Rarely Poisonous and Suspected Poisonous Range Plant. This perennial grass “may become infected with Ergot (*Claviceps*) and cause Ergot poisoning of livestock.”), 85 (102009 - color presentation of dried material), 105 (recorded as *Hilaria mutica* (Buckl.) Benth.)*

Setaria macrostachya (see NOTES and related footnotes 33, 46, 85 and 105 under *Setaria vulpiseta*)

***Setaria vulpiseta* (J.B. de Lamarck) J.J. Roemer & J.A. Schultes: Plains Bristlegrass**

COMMON NAMES: Assaak, Plains Bristlegrass, Xikkaa Kiix, Zacate Tempranero, Zacate Temprano. DESCRIPTION: Terrestrial perennial tufted graminoid (a bunchgrass with somewhat geniculate culms 1 to 4 feet in height, one plant was described as being 2 inches in width at the base, several plants were described as being 8 to 16 inches in width at the base); the stems and leaves are pale to bright green sometimes with a bluish tinge curing to an orange-brown; the flowers may be orange and purple; flowering generally takes place between late April and mid-October (additional records: one for early March and one for mid-November). HABITAT: Within the range of this species it has been reported from mountains; cliffs; rocky canyons; rocky canyon sides; canyon bottoms; canyonettes; rocky talus; bases of cliffs; crevices in rocks; amongst rocky buttes; crests of buttes; rocky ledges; ridges; openings in woodlands; foothills; hills; hilltops; hillsides; rocky and gravelly slopes; bajadas; rocky outcrops; amongst boulders and rocks; sandy dunes; sandy mesquite hummocks; plains; gravelly flats; valley floors; along gravelly roadsides; arroyos; gravelly-sandy-loamy draws; streambeds; sandy creeks; along and in gravelly washes; within drainages; drainage ways; depressions; gravelly-sandy banks of streambeds, sandy riverbeds; creeks and washes; (rocky) edges of streambeds and washes; benches; sandy-loamy bottomlands; sandy floodplains; mesquite bosques; stock tanks; riparian areas, and disturbed areas growing in muddy and moist and dry bouldery, rocky, gravelly, gravelly-sandy and sandy ground; gravelly loam, gravelly-sandy loam and sandy loam ground; sandy clay and clay ground, and cobbly-sandy silty ground sometimes in the partial shade of shrubs and trees, occurring from sea level to 6,000 feet in elevation in the woodland, scrub, grassland, desert scrub and wetland ecological formations. NOTES: *Setaria vulpiseta*, the Plains Bristlegrass has been recorded in many texts as *Setaria macrostachya*; however, it has been reported that *Setaria macrostachya*, with the common name Large-spike Bristlegrass is an EXOTIC species that may also be found in Arizona. There appears to be some confusion as to what's what with this species with regard to its taxonomy. The native Plains Bristlegrass may be an attractive component of a restored native habitat, and the plant is reportedly a good soil binder. Plains Bristlegrass is an important forage grass with a high palatability; however, it is often selectively grazed over other range grasses and does not stand up well to heavy grazing. *Setaria vulpiseta* is native to south-central (again, some say that it is native and some say that it isn't) and southern North America; Central America, and South America. *5, 6, 15 (recorded as *Setaria macrostachya* H.B.K.), 16 (recorded as *Setaria macrostachya* H.B.K.), 33 (recorded as *Setaria macrostachya* H.B.K., Plains Bristlegrass., Page 270), 43 (102409), 46 (recorded as *Setaria macrostachya* H.B.K., Plains Bristlegrass, Page 139 and reidentified as in the Supplement, Page 1041), 48 (recorded as *Setaria macrostachya*), 58 (recorded as *Setaria macrostachya* H.B.K.), 63 (102409 - color presentation of seed), 77 (recorded as *Setaria*

macrostachya H.B.K.), 85 (102409 - *Setaria macrostachya* Kunth and *Setaria vulpiseta* (Lam.) Roemer & J.A. Schultes), 105 (recorded as *Setaria macrostachya* H.B.K.)*

***Sporobolus cryptandrus* (J. Torrey) A. Gray: Sand Dropseed**

COMMON NAMES: Covered-spike Dropseed, Dropseed, Drop Seed Grass; Drop-seed Grass, Large-panicle Vilfa, Larfe-panicled Vilfa, Lesser Dropseed, Prairie Grass, Prairie-grass, Sand Drop-seed, Sand Dropseed, Sand Rush Grass, Sand Rush-grass, Sand Rushgrass, Sporobole à Fleurs Cachées (French, alternate spelling Sporobole à Fleures Cachées also observed), Vai Tava'i (Yaqui, also called this grass Vaso which is the Yaqui generic name for grass), Zacate de Arena. DESCRIPTION: Terrestrial perennial graminoid (a bunchgrass (clumpgrass) with erect, rarely decumbent, culms 1 to 4 feet (one record of 6½ feet) in height and 1 to 8 inches in width at the base, plants 40 inches in height and 4 to 6 inches in width were reported); the foliage may be bluish-green, light green, dark green or purple curing to light straw-yellow; the spikelets (flowers) may be brownish, purplish, bright red-maroon or yellow; flowering generally takes place between late April and early June and between late July and late October (additional records: one for late January, one for early April and one for late May). HABITAT: Within the range of this species it has been reported from mountains; gravelly mountaintops; bouldery, rocky, gravelly-sandy and sandy mesas; sandy plateaus; rocky and sandy rims of canyons; cliffs; rocky and gravelly-loamy canyons; along bouldery-cobbly-sandy and sandy canyon bottoms; gorges; bouldery talus; sandy crevices in boulders and rock walls; bluffs; along tops of bluffs; buttes; rocky ledges; along rocky, gravelly-loamy and sandy ridges; glades; sandy and clayey meadows; tops of cinder cones; sandy foothills; gravelly and gravelly-sandy hills; sandy hillsides; escarpments; along bedrock, bouldery, rocky, rocky-gravelly, rocky-sandy-loamy, shaley, cobbly, gravelly, sandy, sandy-loamy and silty-clayey slopes; rocky outcrops; amongst boulders and rocks; sheltered nooks of rim rock; sandy lava flows; sand hills; sand dunes; sand hummocks; sandy and sandy-loamy prairies; pebbly, gravelly-sandy, sandy and sandy-clayey plains; bouldery, rocky, rocky-sandy, gravelly, sandy, clayey, clayey-loamy and silty-clayey flats; basin floors; gravelly-sandy valley floors; valley bottoms; coastal dunes; sandy coastal plains; sandy coastal flats; gravelly railroad right-of-ways; sandy roadways; along rocky-sandy, cindery, gravelly, gravelly-loamy, sandy, sandy-loamy, sandy-clayey, sandy-silty and clayey roadsides; sandy and clayey arroyos; bottoms of arroyos; draws; bottoms of draws; springs; gravelly-loamy soils along streams; along streambeds; rocky creekbeds; along rivers; along and in sandy riverbeds; along and in rocky, rocky-sandy, gravelly, sandy, sandy-loamy, clayey and silty-clayey washes; within drainages; drainage ways; clayey playas; gravelly-sandy depressions; clayey swales; (sandy and sandy-loamy) banks of arroyos, rivers, washes and lakes; (rocky, gravelly and sandy) edges of draws, gullies; streams, drainage ways, pools and depressions; (sandy) shores of lakes; gravelly and sandy beaches; sandy benches; stony-loamy, sandy and sandy-loamy terraces; loamy bottomlands; sandy floodplains; lowlands; mesquite bosques; around stock tanks (charcos); along and in loamy ditches; rocky and gravelly-sandy riparian areas; loamy waste places, and disturbed areas growing in dry bouldery, bouldery-cobbly-sandy, rocky, rocky-gravelly, rocky-sandy, shaley, cobbly, cindery, gravelly, gravelly-sandy, pebbly and sandy ground; stony loam, gravelly loam, sandy loam, sandy-clayey loam, sandy-silty loam, silty-clayey loam and loam ground; gravelly clay, gravelly-sandy clay, sandy-clay, silty clay and clay ground, and gravelly silty, sandy silty and silty ground, occurring from sea level to 10,300 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, established plants are drought resistant. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food and/or fiber crop. When using this plant in landscaping and re-vegetation projects use plants and/or seed collected from as local a population as possible. Rocky Mountain Bighorn Sheep (*Ovis canadensis*) browse this plant, Scaled Quail (*Callipepla squamata*), Black-tailed Jackrabbits (*Lepus californicus*), Black-tailed Prairie Dogs (*Cynomys ludovicianus*) feed on this plant, small mammals and birds also utilize this plant. *Sporobolus cryptandrus* is native to central and southern North America and southern South America (report for Argentina found in the Germplasm Resources Information Network). *5, 6, 15, 16, 33 (very similar to *Sporobolus flexuosus* and difficult to distinguish

without having mature panicles, Pages 226-227), 43 (102409 - *Sporobolus cryptandrus* A. Gray), 46 (Page 114), 48, 58, 63 (102409 - color presentation), 77, **85** (102409 - color presentation of dried material), 105, 127*

***Trichloris crinita* (M. Lagasca y Segura) L.R. Parodi: False Rhodes Grass**

SYNONYMY: *Chloris crinita* M. Lagasca y Segura, *Trichloris mendocina* (R.A. Philippi) F. Kurtz. COMMON NAMES: False Rhodes Grass, False Rhodesgrass, Feather Fingergrass, Multiflowered Chloris, Rhodes Grass, Twoflower Chloris, Twoflower Trichloris. DESCRIPTION: Terrestrial perennial graminoid (2 to 5 feet in height); the foliage is green or reddish; based on few flowering records observed, flowering generally takes place between late April and mid-October (flowering records: one for late April, one for mid-July and one for mid-October, flowering has been generally described as taking place from late spring to fall). HABITAT: Within the range of this species it has been reported from mountains; mesas, canyons, crevices in rocks; pockets of soil; rocky hills, slopes; bajadas; plains; gravelly-sandy flats; coastal plains; along railroad right-of-ways; along sandy-loamy roadsides; along bottoms of arroyos; along seeps; along sandy washes; drainages; depressions; loamy benches; terraces; floodplains; sandy lowlands, and disturbed areas growing in dry rocky, gravelly-sandy and sandy ground and loam ground, occurring from sea level to 4,000 feet in elevation in the scrub, grassland, and desertscrub ecological formations. NOTES: This large, showy grass may be an attractive component of a restored native habitat. *Trichloris crinita* is native to southwest-central and southern North America and western and southern South America. *5, 6, 15 (recorded as *Chloris crinita* (Lag.) Parodi), 33 (recorded as *Trichloris mendocina* (Phil.) Kurtz, Page 134), 43 (102509), 46 (Page 126), 63 (102509 - color presentation), 58 (recorded as *Chloris crinita* Lag.), **85** (102509 - color presentation)*

Trichloris mendocina (see *Trichloris crinita*)

Tridens pulchellus (see *Dasyochloa pulchella*)

Triodia pulchella (see *Dasyochloa pulchella*)

CLASS MAGNOLIOPSIDA: The DICOTS

Asclepiadaceae: The Milkweed Family

***Matelea parvifolia* (J. Torrey) R.E. Woodson: Spearleaf**

SYNONYMY: *Gonolobus parvifolius* J. Torrey. COMMON NAMES: Angle-pod, Anglepod, Littleleaf Matelea, Little Leaf Milk Vine, Milkweed Vine, Small-leaf Anglepod, Small-leaved Milkvine, Spearleaf. DESCRIPTION: Terrestrial perennial shrub or vine (a clambering, climbing and twining vine 16 inches to 5 feet in length); the stems are gray-green or green; the leaves are green; the flowers may be black, dark brownish-purple, green, greenish-purple, dark purple or purple-brown; based on few flowering records examined, flowering generally takes place between early March and mid-May and between mid-October and early December (flowering records: three for late January, three for early March, three for mid-March, four for late March, three for early April, two for mid-April, one for early May, one for mid-May, one for mid-October, one for late October, five for early November, one for mid-November, one for late November and one for early December); the fruits are long, warty, green seed pods. HABITAT: Within the range of this species it has been reported from rocky mountains; rocky mountainsides; mesas; rocky canyons; canyon bottoms; rocky ridge tops; ridgelines; rocky and stony-gravelly hills; rocky and rocky-gravelly hillsides; bouldery and rocky slopes; bajadas; amongst boulders and rocks; bouldery, cobbly, gravelly and gravelly-sandy flats; along roadsides; along arroyos; springs;

rivers; along and in rocky washes; along drainages; edges of washes; floodplains, and rocky riparian areas growing in dry bouldery, rocky, rocky-gravelly, stony-gravelly, cobbly and gravelly soils, occurring from 1,200 to 5,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: *Matelea parvifolia* is native to southwest-central and southern North America. *5, 6, 15, 16, 43 (110809), 46 (recorded as *Gonolobus parvifolius* Torr., Page 665), 63 (110809), 77, **85** (110809 - color presentation)*

Gonolobus parvifolius (see *Matelea parvifolia*)

Asteraceae (Compositae): The Aster Family

***Acourtia nana* (A. Gray) J.L. Reveal & R.M. King: Dwarf Desertpeony**

SYNONYMY: *Perezia nana* A. Gray. COMMON NAMES: Ban Auppa-ga (Gila River Pima), Desert Holly, Desert-holly, Dwarf Desertpeony. DESCRIPTION: Terrestrial perennial forb/herb (2 to 10 inches in height, plants 4 to 5 inches in height and 3 to 6 inches in width were reported); the holly-like leaves are pale grayish-green or olive-green; the flowers may be cream, pale lavender-pink, lavender, lavender-pink, maroon and white, pale pink-lavender, pink, pink-purple, purple, white or white-pink; flowering generally takes place between late March and early June (additional records: one for late January, one for late February, one for early July, one for late July, one for early August, one for mid-August, one for early September, two for late September, one for mid-October, one for mid-November and two for mid-December). HABITAT: Within the range of this species it has been reported from rocky mountains; sandy mesas; gravelly-loamy canyons; talus slopes; rocky foothills; rocky and gravelly hills; rocky and gravelly hillsides; bouldery, rocky, stony, gravelly, gravelly-sandy and sandy slopes; bajadas; amongst boulders and rocks; gravelly breaks; gravelly plains; rocky, gravelly, gravelly-sandy, sandy, sandy-loamy and clayey flats; basins; basin bottoms; rocky valley floors; valley bottoms; gravelly-loamy roadsides; arroyos; bottoms of arroyos; rocky gullies; gravelly-loamy creekbeds; riverbeds; within gravelly, gravelly-sandy and sandy-clayey washes; drainage ways; playas; sandy-loamy, sandy-clayey-loamy and clayey-loamy swales; (clayey-loamy) banks of washes; benches; gravelly and sandy terraces; floodplains; mesquite mosques; riparian areas, and disturbed areas growing in dry desert pavement; bouldery, rocky, rocky-sandy, stony, gravelly, gravelly-sandy and sandy ground; gravelly loam, sandy loam, sandy-clayey loam, clayey loam, silty loam and loam ground, and sandy clay and clay ground often in the shade of trees and shrubs, occurring from 1,200 to 7,100 feet (one record for 8,500 feet) in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. Consider using Desert Holly as a ground cover under larger shrubs and trees. The flowers give off a fragrance similar to that of violets or lilacs. *Acourtia nana* is native to southwest-central and southern North America. *5, 6, 15, 16, 28 (color photograph), 43 (110809), 46 (recorded as *Perezia nana* Gray, Page 957), 58, 63 (110809 - color presentation), 77, **85** (110909 - color presentation), 115 (color presentation)*

***Acourtia wrightii* (A. Gray) J.L. Reveal & R.M. King: Brownfoot**

SYNONYMY: *Perezia wrightii* A. Gray. COMMON NAMES: Brownfoot, Desert Holly, Perezia, Pink Perezia, Pink Perezia, Wright's Desertpeony. DESCRIPTION: Terrestrial perennial forb/herb (1 to 5 feet in height, one plant was recorded as being 1 foot in height with a crown 1 foot in width); the holly-like leaves are dark green; the flowers may be lavender, pink, pink-brown, pink-lavender, pink-purple, pale purple, purple, white, white & pink, whitish-maroon or white & purple; flowering generally takes place between early February and early July and sometimes in autumn between early September and early November (additional records: one for mid-August, one for late November and one for early December). HABITAT: Within the range of this species it has been reported from mountains; plateaus; rock cliffs; crater walls; rocky canyons; rocky canyon bottoms; talus slopes; bases of cliffs; along crevices in boulders; buttes; along ledges; ridges; ridgetops; foothills; rocky, stony-gravelly and sandy hills; rocky

and rocky-gravelly-loamy hillsides; bouldery-rocky, rocky, rocky-gravelly, shaley, gravelly and sandy slopes; sandy alluvial fans; gravelly and sandy bajadas; along bedrock and rocky outcrops; amongst boulders and rocks; bases of boulders; in shaded alcoves; rocky plains; rocky and silty flats; railroad right-of-ways; rocky and gravelly-sandy-clayey-loamy roadsides; along rocky arroyos; draws; gullies; ravines; seeps; rocky springs; along creeks; along rocky, gravelly and sandy washes; along drainage ways; rocky banks of streams and washes; edges of washes; mudflats; beaches; floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry desert pavement; bouldery, bouldery-rocky, rocky, rocky-gravelly, rocky-sandy, stony-gravelly, gravelly and sandy ground; rocky-gravelly loam, rocky silty loam, gravelly-sandy-clayey loam, sandy loam, silty-clayey loam and silty loam ground, and silty ground, occurring from 700 to 7,300 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, the flowers are reported to be fragrant. This plant was reported to have been utilized by native peoples of North America; it was noted as having been used as a drug or medication. *Acourtia wrightii* is native to southwest-central and southern North America. *5, 6, 15, 16, 28 (color photograph), 43 (110809), 46 (recorded as *Perezia wrightii* Gray, Page 957), 58, 63 (110909 - color presentation), 77, 85 (110909 - color presentation), 115 (color presentation), 127, **WTK** (March 9, 2007)*

***Ambrosia deltoidea* (J. Torrey) W.W. Payne: Triangle Bur Ragweed**

SYNONYMY: *Franseria deltoidea* J. Torrey. COMMON NAMES: Burrobush, Bur-sage, Bursage, Chamizo Forrajero, Chicurilla, Rabbit Bush, Kokomak Segoi (Pima), Shegoi (Pima), Todshag (Papago), Triangle Bur Ragweed, Triangle Burr Ragweed, Triangle Bursage, Triangle-leaf Bursage, Triangle-leaved Bursage, Triangle-leaf Burr Ragweed. DESCRIPTION: Terrestrial perennial evergreen (or drought-deciduous) subshrub or shrub (1 to 4 feet in height, one plant was described as being 2 feet in height and width); the leaves are gray, gray-green or green; the flowers are greenish, greenish-yellow, purple, white or yellow; flowering generally takes place between early January and early May (additional records: three for late May; flowering as late as July has been reported). HABITAT: Within the range of this species it has been reported from mountains; rocky mesas; rocky canyons; canyon bottoms; bases of cliffs; buttes; ridges; rocky foothills; rocky hills; rocky hillsides; rocky, gravelly and gravelly-clayey slopes; bajadas; lava flows; dunes; sandy plains; rocky, stony-chalky, gravelly and sandy flats; basins; rocky valley floors; along rocky-sandy roadsides; shallow arroyos; runnels; riverbeds; along and in stony-gravelly, gravelly and sandy washes; within drainages; (rocky and sandy) banks of creeks and washes; edges of dry lakes (playas); margins of washes; gravelly terraces; bottomlands; floodplains; riparian areas, and disturbed areas growing in moist and dry desert pavement; rocky, rocky-gravelly, rocky-sandy, stony-gravelly, gravelly, gravelly-sandy and sandy ground; gravelly-sandy loam and loam ground; rocky clay, gravelly clay and sandy clay ground, and stony chalky ground, occurring from 100 to 4,000 feet in elevation in the grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat and may be useful in the restoration of disturbed habitat. It may live to be about 50 years of age. The Triangleleaf Bursage serves as a nurse plant for Saguaro (*Carnegiea gigantea*), Ocotillo (*Fouquieria splendens*), Foothill Paloverde (*Parkinsonia microphylla*) and other woody plants. The Triangleleaf Bursage is one of the first plants to colonize in open spaces. *Ambrosia deltoidea* is native to southwest-central and southern North America. *5, 6, 13, 15, 16, 28 (color photograph), 43 (070910), 46 (recorded as *Franseria deltoidea* Torr., Page 896), 63 (111009 - color presentation), 77 (color photograph #68), 85 (111009 - color presentation), 91, 115 (color presentation), **WTK** (March 9, 2007)*

***Baccharis sarothroides* A. Gray: Desertbroom**

COMMON NAMES: Amargo, Broom Baccharis, Caasot Caocl (Seri), Desert Broom, Desert-broom, Desertbroom, Escoba, Greasewood, Groundsel, Hierba del Pasma, Mexican Broom, Romerillo, Rosin Brush, Rosin-brush, Rosin Bush, Shooshk Vakch ("Wet Shoes" - Pima), Soosk Vaks ("Wet Shoes" - questionably Maricopa), Wet Shoes. DESCRIPTION: Terrestrial perennial deciduous shrub (3 to 10 feet in height, one plant was described as being 40 inches in height and 40 inches in width, one plant was

described as being 7 feet in height and 8 feet in width); the foliage is bright green or yellow-green; the disc (no ray flowers) flowers (dioecious) are cream, golden, rust, white or yellow; flowering generally takes place between mid-September and late February (additional records: one for mid-March, two for late March, two for mid-April, one for late April, one for mid-July, one for early August and one for late August). HABITAT: Within the range of this species it has been reported from mountains; mesas; canyons; canyon bottoms; chasms; ridges; foothills; silty-clayey hills; rocky hillsides; rocky slopes; bajadas; debris fans; sandy plains; rocky flats; valley floors; coastal plains; along rocky, rocky-gravelly-sandy-clayey-loamy, rocky-gravelly-sandy-silty-clayey-loamy and sandy roadsides; along arroyos; along sandy and clayey bottoms of arroyos; draws; gulches; gullies; springs; along streams; streambeds; along gravelly-sandy creeks; gravelly and sandy creekbeds; along rivers; along bouldery-cobbly-sandy, rocky-sandy, gravelly and sandy riverbeds; along and in cobbly, gravelly and sandy washes; along drainages; along drainage ways; waterholes; playas; oases; along (gravelly and sandy) banks of arroyos, rivers and washes; edges of washes; beaches; sandy benches; terraces; bottomlands; floodplains; lowlands; mesquite bosques; along canals; along ditches; muddy and sandy riparian areas, and disturbed areas growing in muddy and damp and dry bouldery-cobbly-sandy, rocky, rocky-sandy, cobbly, gravelly, gravelly-sandy and sandy ground; rocky-gravelly-sandy-clayey loam and rocky-gravelly-sandy-silty-clayey loam ground; silty clay and clay ground, and sandy silty ground, occurring from sea level to 6,200 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, consider planting only male plants to eliminate seed production. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial beverage and/or fiber crop; it was also noted as having been used for making tools and as a drug or medication. *Baccharis sarothroides* is native to Southwestern and southern North America. *5, 6, 13, 15, 16, 18, 26 (color photograph), 28 (color photograph), 43 (111209), 46 (Page 883), 48, 58, 63 (111209), 77, 85 (111309 - color presentation), 115 (color presentation), 127, **WTK** (March 9, 2007)*

***Gutierrezia* M. Lagasca y Segura: Snakeweed**

COMMON NAMES: Snake-weed, Snakeweed *43 (062310), 46 (Pages 852-853), 63 (022207), **WTK** (March 9, 2007)*

Franseria deltoidea (see *Ambrosia deltoidea*)

Perezia nana (see *Acourtia nana*)

Perezia wrightii (see *Acourtia wrightii*)

Brassicaceae (Cruciferae): The Mustard Family

***Arabis perennans* S. Watson: Perennial Rockcress**

SYNONYMY: *Arabis eremophila* E.L. Greene, *Boechera perennans* (S. Watson) W.A. Weber. COMMON NAMES: Perennial Rockcress, Rock Cress, Stiff-arm Rock Cress, Stiffarm Rock Cress. DESCRIPTION: Terrestrial perennial forb/herb or shrub (8 to 40 inches in height, plants 8 to 12 inches in height and 6 to 17 inches in width were reported); the leaves are gray-green; the flowers may be pale blue-lavender, bluish-purple, cream, lavender, pink, pink-lavender, pinkish-purple, dull mauve, pale purple, purple, purple-magenta, purplish-pink, purplish-rose, reddish-violet, rose-magenta, violet-lavender, white & lavender or white-purple; flowering generally takes place between early February and early July (additional records: one for early January, one for mid-January, one for early August, one for early October and one for early December). HABITAT: Within the range of this species it has been reported from mountains; along mountaintops; rocky mountainsides; sandy mesas; sandy plateaus; rocky cliffs; rock faces; bouldery, rocky and rocky-sandy canyons; rocky and shaley-sandy canyon walls;

bedrock, rocky, gravelly-sandy and sandy canyon bottoms; talus slopes; along sandy bases of cliffs; crevices in rocks; bluffs; rocky knobs; summits of laccoliths; rocky ledges; sandy ridges; ridgetops; rocky openings in woodlands; meadows; rocky-gravelly foothills; rocky hills; bouldery and rocky hillsides; sandy bases of escarpments; bedrock, bouldery, bouldery-gravelly, bouldery-sandy, rocky, rocky-shaley, rocky-gravelly, rocky-sandy, rocky-loamy, cobbly-sandy, cobbly-loamy, cindery, gravelly, gravelly-silty, sandy, sandy-loamy and loamy slopes; bajadas; rocky outcrops; amongst boulders and rocks; bases of boulders; lava flows; rocky mounds; flats; basins; along sandy valley floors; along roadbeds; along gravelly roadsides; rocky walls of arroyos; along draws; gulches; bouldery-sandy and rocky ravines; springs; along streams; gravelly streambeds; along creeks; along rivers; along and in rocky, rocky-gravelly, gravelly and sandy washes; within drainages; bouldery-cobbly drainage ways; marshes; (rocky) banks of gullies, streams and washes; (rocky-loamy and gravelly) edges of arroyos, streams and washes; gravelly terraces; rocky and gravelly-sandy riparian areas, and disturbed areas growing in moist and dry bouldery, bouldery-cobbly, bouldery-gravelly, bouldery-sandy, rocky, rocky-shaley, rocky-gravelly, rocky-sandy, shaley, shaley-sandy, cobbly-sandy, cindery, gravelly, gravelly-sandy and sandy ground; rocky loam, rocky-sandy loam, cobbly loam, gravelly-clayey loam, sandy loam, clayey loam and loam ground; sandy clay and clay ground, and gravelly silty and silty ground, occurring from 600 to 9,500 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant was reported to have been utilized by native peoples of North America; it was noted as having been used as a drug or medication. *Arabis perennans* is native to southwest-central and southern North America. *5, 6, 15, 16, 28 (color photograph), 43 (010310), 46 (Page 353), 58, 63 (010310 - color presentation), 77, **85** (010410 - color presentation), 115 (color presentation), 127*

Arabis eremophila (see *Arabis perennans*)

Boechera perennans (see *Arabis perennans*)

Cactaceae: The Cactus Family

***Carnegiea gigantea* (G. Engelmann) N.L. Britton & J.N. Rose: Saguaro**

SYNONYMY: *Cereus giganteus* G. Engelmann. COMMON NAMES: Giant Cactus, Giant Cereus, Ha:saan (Tohono O'odham), Ha Shun (Pima), Mashad (Tohono O'odham), Pitahaya (Spanish Conquistadors), Sage-of-the-desert, Saguaro (Spanish), Sahuaro. DESCRIPTION: Terrestrial perennial stem-succulent tree (erect stems 5 to 60 feet in height and 6 to 30 inches in diameter); the plants are green; the spines are yellow or reddish-brown aging to gray or gray-black; the flowers (2 to 3 inches in diameter) are a waxy creamy-white opening at about 8 p.m. and closing at about 5 p.m. the next day with around four blooms opening per day over a 30 day period; flowering generally takes place between late April and mid-June (additional records: one for late March, one for early July, one for mid-July, two for early September and one for early October), the ripe fruits (2¼ to 3 inches in length and 1 to 1½ inches in diameter) split into 2 to 6 segments that curl back to reveal the red inner lining of the rinds which are sometimes mistakenly thought to be red flowers. HABITAT: Within the range of this species it has been reported from mountains; mesas; canyons; canyon walls; buttes; ridges; ridgelines; rocky foothills; rocky and gravelly hills; rocky hilltops; rocky hillsides; rocky, gravelly, gravelly-loamy and sandy-clayey-loamy slopes; rocky and gravelly bajadas; rocky outcrops; amongst boulders and rocks; stabilized sandy and sandy-powdery dunes; plains; gravelly and sandy flats; valley floors; along arroyos; along and in riverbeds; within sandy washes; drainages; floodplains, and mesquite bosques growing in dry desert pavement; bouldery, rocky, gravelly, sandy and sandy-powdery ground, and gravelly loam and sandy-clayey loam ground, occurring from sea level to 5,100 feet in elevation in the scrub, grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, fodder (seeds),

beverage and/or fiber crop; it was also noted as having been used as tools, ceremonial items and musical instruments, and as an indicator of the changing of the seasons (with the Saguaro harvest marking the beginning of a new year). Saguaro are very slow to establish, a 5 year old plant may be no more than ¼ to ½ inch in height. The growth rate of Saguaro is extremely variable. William G. McGinnies in his book “Discovering the Desert” reports that a plant 36 inches in height may be from 20 to 50 years of age, he also presents a table of typical growth rates reporting the following: 4 inches - 8.0 years, 8 inches - 12.5 years, 16 inches - 19.1 years, 32 inches - 27.3 years, 3.3 feet - 30.3 years, 6.6 feet - 40.5 years, 10 feet - 47.5 years, 13 feet - 54 years, 16 feet - 60.0 years, 18 feet - 74.0 years, 20 feet - 83.0 years, 25 feet - 107.0 years, 30 feet - 131.0 years, and 35 feet - 157.0 years. The growth rate of propagated and cultivated saguaros is much faster. One of the largest known saguaros, located in Saguaro National Monument, was reported to be 52 feet in height, had 52 arms, weighed an estimated 10 tons and was thought to be 235 years of age. Cristate forms have been reported. The Broad-billed Hummingbird (*Cynanthus latirostris*), Broad-tailed Hummingbird (*Selasphorus platycercus*), Costa’s Hummingbird (*Calypte costae*), Curved-billed Thrasher (*Toxostoma curvirostre*), Lesser Long-nosed Bat (*Leptonycteris curasoae* subsp. *yerbabuena*) and Rufous Hummingbird (*Selasphorus rufus*) have been observed visiting the flowers. Coyotes (*Canis latrans*), Desert Mule Deer (*Odocoileus hemionus* subsp. *crooki*), Desert Bighorn Sheep (*Ovis canadensis* subsp. *mexicana*), Javelina (*Peccari tajacu*) and White-winged Doves (*Zenaida asiatica*) as well as other animals and birds feed on the saguaro fruit and seeds. the Gila Woodpecker (*Melanerpes uropygialis*) and Gilded Flicker (*Colaptes chrysoides*) make holes in this plant for their nests which are later utilized by the Ash-throated Flycatcher (*Myiarchus cinerascens*), Cactus Wren (*Campylorhynchus brunneicapillus*), Elf Owl (*Micrathene whitneyi*), House Finch (*Carpodacus mexicanus*), Lucy’s Warbler (*Vermivora luciae*), Purple Martin (*Progne subis*) and Cactus Wren (*Campylorhynchus brunneicapillus*). Red-tailed Hawks (*Buteo jamaicensis*), White-winged Doves (*Zenaida asiatica*) and other birds nest on the arms of the plant. *Carnegiea gigantea* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Cereus giganteus* Engelm., Pages 108-111, color photographs including habitat), 13 (color photographs including habitat with associated species: Plates C.2 and D.3), 15 (color photograph on Page 77 includes habitat and associated species), 16, 18, 26 (color photograph), 27 (recorded as *Cereus giganteus*, Pages 64-65, color photographs: Plates 39, 39A & 39B, Page 102), 28 (recorded as *Cereus giganteus*, color photograph), 38 (color photograph), 43 (011610), 45 (color photograph), 46 (Page 569), 48 (recorded as *Cereus giganteus*), 52 (recorded as *Cereus giganteus*, color photograph), 53 (recorded as *Cereus giganteus* Engelm.), 58 (recorded as *Cereus giganteus* Engelm.), 63 (011610 - color presentation), 77 (color photograph #63), 85 (011610 - color presentation), 86 (recorded as *Cereus gigantea*, color photograph), 91, 107, 115 (color presentation), 119, 127, 134, **WTK** (March 9, 2007)*

Cereus giganteus (see *Carnegiea gigantea*)

Cereus thurberi (see *Stenocereus thurberi*)

***Cylindropuntia bigelovii* (G. Engelmann) F.M. Knuth: Teddybear Cholla**

SYNONYMY: *Opuntia bigelovii* G. Engelmann. COMMON NAMES: Arizona Jumping Cactus, “Ball” Cholla, Cholla Guera, Go'te (Seri), Jumping Cactus, Jumping Cholla, Silver Cholla, Teddybear Cactus, Teddy Bear Cholla, Teddy-bear Cholla, Teddybear Cholla. DESCRIPTION: Terrestrial perennial stem-succulent subshrub or shrub (20 inches to 10 feet in height, one plant was reported to be just over 8 feet in height and 40 inches in width with 2 to 3 main trunks); the central trunk is black or dark brown; older branches are dark-brown; the stems (3 to 10 inches in length and 1¼ to 2½ inches in diameter) are bluish, light green, green or bluish-green; the spines are golden, silvery, tan, pale yellow or yellow aging to dark brown; the glochids are yellow; the flowers (1 to 1½ inches in diameter) may be chartreuse-yellow, cream tinged with rose, green, green-yellow, greenish-yellow, magenta, pink, white-yellow, yellow tinged with red-purple or white tinged with lavender; the anthers are yellow, deep yellow, yellow-orange or deep yellow-orange; the stigma lobes are cream, dark chartreuse-green, green, dark green or

olive green; flowering generally takes place between early March and mid-June (additional records: one for late January, one for early February, one for early September, one for mid-November, two for late November and one for early December); the nearly spineless fruits ($\frac{1}{2}$ to $\frac{3}{4}$ inch in length and $\frac{1}{2}$ to $\frac{3}{4}$ inch in diameter) are greenish-yellow, yellow or yellow-green and fleshy when ripe. HABITAT: Within the range of this species it has been reported from mountains; rocky and sandy mountainsides; canyons; canyon bottoms; cliffs; talus slopes; bluffs; rocky ridges; rocky ridgetops; rocky foothills; rocky and sandy hills; rocky hillsides; rocky, rocky-gravelly, rocky-sandy, gravelly and sandy slopes; rocky alluvial fans; gravelly-loamy bajadas; plains; cobbly-silty, gravelly and silty flats; basins; valley floors; along roadsides; arroyos; along and in rocky, gravelly and sandy washes; sandy drainages; benches; lowlands, and disturbed areas growing in dry desert pavement; rocky, rocky-gravelly, rocky-sandy, gravelly and sandy ground; gravelly loam and silty loam ground; clay ground, and cobbly-silty and silty ground, occurring from sea level to 4,400 feet in elevation in the scrub and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. This is the spiniest of the cholla cacti in Arizona. Thomas Kearney and Robert Peebles in their book *Arizona Flora* had this to say about the Teddybear Cholla: "The combination of barbed spines and densely armed, easily detached joints has earned profound respect for this formidable cholla." Teddy-bear Chollas may live to be 60 or more years of age. The Teddybear Cholla is a preferred nesting site of the Cactus Wren (*Campylorhynchus brunneicapillus*). Pack Rats (*Neotoma* sp.) use the joints of this plant in the construction of their nests. The change in nomenclature in USDA NRCS has not been recognized in BONAP, species remains as *Opuntia bigelovii* (accessed 041806). *Cylindropuntia bigelovii* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Opuntia bigelovii* Engelm., Pages 50, 52 & 54-55), 15 (recorded as *Opuntia bigelovii* Engelm., color photograph on Page 77 includes habitat and associated species), 18, 26 (genus, recorded as *Opuntia*), 27 (Page 5, color photographs including habitat: Plates 5 & 5A, Page 94), 28 (recorded as *Opuntia bigelovii*, color photograph), 43 (052110), 45 (color photograph), 46 (recorded as *Opuntia bigelovii* Engelm., Page 584), 48, 63 (052110 - color presentation including habitat), 77 (recorded as *Opuntia bigelovii* Engelm., color photograph #13), 85 (052110 - color presentation), 86 (color photograph), 91 (recorded as *Opuntia bigelovii* Engelm.), 115 (color presentation), 119 (recorded as *Opuntia bigelovii* Engelm.), 127, **WTK** (March 9, 2007)*

***Cylindropuntia fulgida* (G. Engelmann) F.M. Knuth var. *fulgida*: Jumping Cholla**

SYNONYMY: *Opuntia fulgida* G. Engelmann, *Opuntia fulgida* G. Engelmann var. *fulgida*. COMMON NAMES: Chain Cholla, Chain-fruit Cholla, Cholla, Cholla Brincadora, Choya, Jumping Cahin-fruit Cholla, Jumping Cholla, Sonora Jumping Cholla, Velas de Coyote. DESCRIPTION: Terrestrial perennial stem-succulent shrub or tree (3 to 15 feet in height, one plant was reported as being 4 $\frac{1}{4}$ feet in height and 40 inches in width, one plant was reported as being 4 $\frac{1}{4}$ feet in height and 8 $\frac{1}{4}$ feet in width, one plant was reported as being 6 $\frac{1}{2}$ feet in height and 5 feet in width, one plant was reported as being 10 feet in height and 13 feet in width); the stems are green or purple; the spines are golden-yellow turning brown with age; the flowers ($\frac{3}{4}$ to 1 inch in diameter) are cream-yellow, pink, pink-purple, purple, purple-pink, red-purple, rose-pink or yellow tinged with pink; the anthers are white; flowering generally takes place between mid-April and mid-September (additional record: one for early December); the smooth fleshy fruits ($\frac{3}{4}$ to 2 inches in length and $\frac{3}{4}$ to 1 inch in diameter) are gray-green, green or purple forming clusters or pendant "chains". HABITAT: Within the range of this species it has been reported from mountains; mesas; canyons; ledges; ridges; hills; hillsides; rocky, gravelly-loamy and sandy slopes; gravelly bajadas; plains; rocky-gravelly, gravelly, sandy and sandy-silty flats; along valley floors; along rocky-gravelly and sandy roadsides; along creeks; along and in washes; banks of streams, creeks and washes; edges of washes; terraces, and floodplains growing in dry desert pavement; rocky, rocky-gravelly, gravelly and sandy soils; gravelly loam and silty-clayey loam ground; clay ground, and sandy silty ground, occurring from 800 to 4,100 feet in elevation in the grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant

was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. Each year, following flowering, additional fruits are added to the end of the chains. Chain-fruit Chollas may live to be from 40 to 80 years of age. The Chain-fruit Cholla is a preferred nesting site of the Cactus Wren (*Campylorhynchus brunneicapillus*). The Costa's Hummingbird (*Calypte costae*) has been observed visiting the flowers. Deer and Javelina feed on the fruits. The change in nomenclature in USDA NRCS has not been recognized in BONAP, species remains as *Opuntia fulgida* (accessed 041806). *Cylindropuntia fulgida* var. *fulgida* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Opuntia fulgida* Engelm. var. *fulgida*, Pages 49-52), 15 (recorded as *Opuntia fulgida* Engelm. var. *fulgida*), 16 (recorded as *Opuntia fulgida* Engelm.), 26 (genus, recorded as *Opuntia*), 27 (species, Pages 10-11, color photograph: Plate 10, Page 96), 28 (recorded as *Opuntia fulgida*, color photograph), 43 (011810), 45 (species, color photograph of species), 46 (recorded as *Opuntia fulgida* Engelm., Page 585), 48 (genus, recorded as *Opuntia*), 52 (recorded as *Opuntia fulgida*, color photograph), 53 (recorded as *Opuntia fulgida* Engelm.), 63 (011810 - color presentation), 77 (recorded as *Opuntia fulgida* Engelm. var. *fulgida*), 85 (011810 - color presentation), 91 (recorded as *Opuntia fulgida* Engelm. var. *fulgida*), 115 (color presentation of species), 119 (recorded as *Opuntia fulgida* Engelm.), 127, **WTK** (March 9, 2007)*

***Cylindropuntia leptocaulis* (A.P. de Candolle) F.M. Knuth: Christmas Cactus**

SYNONYMY: *Opuntia leptocaulis* A.P. de Candolle. COMMON NAMES: Agujilla, Alfilerillo (Spanish), Catalineria (Spanish), Christmas Cactus, Christmas Cholla, Darning Needle Cactus, Desert Christmas Cactus, Desert Christmas Cholla, Diamond Cactus, Holycross Cholla, Naf (or Nav?, Gila River Pima), Pencil Cactus, Pencil Cholla, Pencil-joint Cholla, Pipestem Cactus, Rat-tail Cactus, Rattail Cactus, Slender-stem Cactus, Tajasilla, Tasajilla (Hispanic), Tasajillo (Spanish), Tasajo (Spanish), Tesajo (Hispanic), Tesajo Cactus (Christmastree Cacti). DESCRIPTION: Terrestrial perennial stem-succulent shrub (1 to 6 feet in height (sometimes becoming vine-like and growing upwards with support 8 to 15 feet in height), one plant was reported as being 2 feet in height and 2 feet in width, one plant was reported as being 30 inches in height and 5 feet in width, one plant was reported as being 40 inches in height and 5 feet in width, one plant was reported as being 4 feet in height and 8 feet in width, one plant was reported as being 5 feet in height and 8¼ feet in width); the stems are gray-green, green, purplish or yellow-green; the spines gray-brown, purple-brown, red-brown or yellow-brown often being paler toward the tip; the glochids are reddish-brown or yellow; the anthers are yellow; the flowers (3/8 to 3/4 inch in diameter) are bronze, cream, light green-cream, cream-yellow, green, green-yellow, greenish-cream, greenish-yellow, pale yellow, yellow or whitish; flowering generally takes place between late March and late June (additional records: two for mid-July, one for late July, one for early August, one for early October, one for mid-October and one for late October); the spineless (with glochids) fleshy fruits (1/2 to 3/4 inch in length and 1/4 to 7/16 inch in diameter) are coral, orange, orange-red, red, reddish-orange, scarlet, scarlet-red or yellow when mature. HABITAT: Within the range of this species it has been reported from mountains; sandy mountainsides; rocky-sandy and silty mesas; along cliffs; rocky canyons; rocky canyon bottoms; rocky talus slopes; rocky ledges; gravelly ridges; foothills; rocky and rocky-gravelly hills; hilltops; rocky hillsides; rocky, gravelly, gravelly-sandy-loamy, sandy and silty-loamy slopes; clayey-loamy alluvial fans; gravelly, gravelly-silty and sandy bajadas; rocky and gypsum outcrops; amongst cobbles; sand hills; sandy lava flows; lava beds; breaks; sandy and clayey-loamy plains; rocky-sandy, gravelly, gravelly-sandy and sandy flats; basins; valley floors; gravelly and gravelly-sandy roadsides; within gravelly and sandy arroyos; bottoms of arroyos; along ravines; riverbeds; along and in rocky, gravelly and sandy washes; sandy drainages; along (cobbly-sandy) banks of rivers and drainages; edges of arroyos, ravines and washes; rocky and sandy benches; terraces; bottomlands; floodplains; along fencelines; along ditches; riparian areas, and disturbed areas growing in dry desert pavement; rocky, rocky-gravelly, rocky-sandy, cobbly-sandy, gravelly, gravelly-sandy and sandy ground; gravelly-sandy loam, clayey loam, silty loam and loam ground; rocky-sandy clay and loamy clay ground, and gravelly silty and silty ground often found growing within grasses, shrubs or trees, occurring from sea level to 5,900 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations.

NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. The Desert Christmas Cactus is believed to have a life span of about 50 years. A high mortality rate is to be expected with plants coming into contact with fire. Hummingbirds have been observed visiting the flowers; the fruits are eaten by birds and small mammals, and Cochineal Scale (*Dactylopius coccus*) has been observed growing on this plant. The change in nomenclature in USDA NRCS has not been recognized in BONAP, species remains as *Opuntia leptocaulis* (accessed 041806). *Cylindropuntia leptocaulis* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Opuntia leptocaulis* DC., Pages 56-58), 15 (recorded as *Opuntia leptocaulis* DC.), 16 (recorded as *Opuntia leptocaulis* DC.), 18, 26 (genus, recorded as *Opuntia*), 27 (Page 2, color photograph: Plate 2, Page 94), 28 (recorded as *Opuntia leptocaulis*, color photograph), 43 (011910), 45 (color photograph), 46 (recorded as *Opuntia leptocaulis* DC., Page), 48 (genus, recorded as *Opuntia*), 58 (recorded as *Opuntia leptocaulis* DC.), 63 (011910 - color presentation), 77 (recorded as *Opuntia leptocaulis* DC.), **85** (011910 - color presentation), 86 (recorded as *Opuntia leptocaulis*, color photograph), 91 (recorded as *Opuntia leptocaulis* DC.), 115 (color presentation), 119 (recorded as *Opuntia leptocaulis* DC.), 127, **WTK** (March 9, 2007)*

***Cylindropuntia spinosior* (G. Engelmann) F.M. Knuth: Walkingstick Cactus**

SYNONYMY: *Opuntia spinosior* (G. Engelmann) J.W. Toumey. COMMON NAMES: Cane Cholla, Cardenche, Handgrip Cholla, Spiny Cholla, Tasajo, Tourney-cane Cholla (Arizona), Walkingstick Cactus, Walking Stick Cholla. DESCRIPTION: Terrestrial perennial stem-succulent shrub (16 inches to 10 feet in height, one plant was described as being 6½ feet in height and 5 to 6½ feet in width, one plant was described as being 6½ feet in height and 10 feet in width); the stems may be brown-green, grayish-maroon, grayish-purple, green, purple or purplish-green; the spines may be brown, gray, pale pink, pink, purplish-gray, reddish-gray or tan; the glochids may be tan, yellow or yellowish-white aging to gray; the flowers (1¼ to 2 inches in diameter) may be bronze-purple, brown, greenish-yellow, magenta, magenta-red, maroon, orange, pink, dark pink, light purple, purple, purple-pink, red, dark red, red-purple, red & yellow, saffron, salmon-pink, terra-cotta, white or yellow; the anthers are yellow; flowering generally takes place between early April and early August (additional records: three for early January, two for early February and one for late September); the fleshy ripe fruits (1 to 1¼ inches in length and ¾ to 1 inch in diameter) are bright lemon-yellow, red, bright yellow, pale yellow, yellow, yellow-green, yellowish-green or yellow with a reddish cast and remain on the plant for some time. HABITAT: Within the range of this species it has been reported from mountains; mountaintops; mountainsides; mesas; rocky canyons; canyon bottoms; talus, ridgelines; foothills; rocky hills; rocky hillsides; along rocky, rocky-sandy and sandy slopes; bajadas; rock outcrops; amongst rocks; plains; gravelly, gravelly-sandy and silty flats; grassy valley floors; roadsides; arroyos; rocky draws; springs; along creeks; creekbeds; along sandy washes; drainages; along drainage ways; sandy flood channels; terraces; floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry rocky, rocky-sandy, gravelly, gravelly-sandy and sandy ground; silty-clayey loam, silty loam and loam ground, and silty ground, occurring from 900 to 7,200 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. The Cactus Wren (*Campylorhynchus brunneicapillus*) nests in the branches. The change in nomenclature in USDA NRCS has not been recognized in BONAP, species remains as *Opuntia spinosior* (accessed 041806). *Cylindropuntia spinosior* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Opuntia spinosior* (Engelm.) Toumey, Pages 39-43, color photograph), 15 (recorded as *Opuntia spinosior* (Engelm.) Toumey), 16 (recorded as *Opuntia spinosior* (Engelm.) Toumey), 26 (genus, recorded as *Opuntia*), 27 (Page 14, color photograph: Plate 12, Page 96), 28 (color photograph), 43 (063009), 45 (color photographs), 46 (recorded as *Opuntia spinosior* (Engelm. & Bigel.) Toumey, Page 585), 48 (genus, recorded as *Opuntia*), 53, 58 (recorded as *Opuntia spinosior* (Engelm.) Toumey), 63 (011910 - color presentation), 77 (recorded as *Opuntia spinosior*

(Engelm.) Toumey), 85 (012010 - color presentation), 115 (color presentation), 119, 127, **WTK** (March 9, 2007)*

***Cylindropuntia versicolor* (G. Engelmann ex J.M. Coulter) F.M. Knuth: Staghorn Cholla**

SYNONYMY: *Opuntia versicolor* G. Engelmann ex J.M. Coulter. COMMON NAMES: Deer Horn Cactus, Deer Horn Cholla, Deerhorn Cholla, Morada Cholla (Spanish), Staghorn Cholla, Tree Cholla. DESCRIPTION: Terrestrial perennial stem-succulent shrub (3 to 15 feet in height, one plant was reported to be 40 inches in height with a crown 40 inches in width, one plant was reported to be 40 inches in height with a crown 6½ feet in width, one plant was reported to be 50 inches in height with a crown 40 inches in width, one plant was reported to be 51 inches in height with a crown 6½ feet in width, one plant was reported to be 63 inches in height with a crown 87 inches in width, one plant was reported to be 75 inches in height with a crown 87 inches in width); the stems are green, green-purple, greenish-red, maroon, purple, purple-green or dark purple-red; the spines are dark brown, gray, pinkish, purple-brown, dark reddish-brown or whitish; the glochids are reddish-brown, yellow or dark yellow; the flowers (1¼ to 2¼ inches in diameter) are bronze, bronze-red, brown, burnt orange, gold, green, lavender, magenta, orange, orange-brown, orange-red, orange-rust, pink-red, purple, red, rose, rose-purple, yellow, yellow-green or yellow-green-bronze; the anthers are yellow; flowering generally takes place between early April and mid-June (additional records: one for early January, one for early March, one for late August and one for mid-September); the fleshy, spineless or nearly spineless pear-shaped fruits (¾ to 1¾ inches in length and ¾ inch in diameter) are green tinged with lavender, purple, straw-yellow, red, bright yellow or yellowish-green sometimes tinged with purple or red, sometimes forming chains of 2 to 4 fruits. HABITAT: Within the range of this species it has been reported from mountains; rocky mountainsides; mesas; rocky and sandy canyons; rocky canyon bottoms; ridges; foothills; rocky and rocky-gravelly hills; rocky hillsides; rocky slopes; gravelly-sandy alluvial fans; rocky and gravelly-sandy bajadas; sand dunes; plains; gravelly and gravelly-sandy flats; sandy valley floors; along roadsides; along arroyos; ravines; along sandy streambeds; along sandy washes; playas; sandy gravelbars; strands, and riparian areas growing in dry rocky, rocky-gravelly, gravelly, gravelly-sandy and sandy ground, occurring from sea level to 5,000 feet in elevation in the woodland, grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. Varied flower colors between plants, the cascading branches of the larger plants, along with pendulous fruits make this an attractive plant. The change in nomenclature in USDA NRCS has not been recognized in BONAP, species remains as *Opuntia versicolor* (accessed 041806). *Cylindropuntia versicolor* is native to southwest-central and southern North America. *5, 6, 12 (color photograph, recorded as *Opuntia versicolor*, Pages 43 & 45-46), 15 (recorded as *Opuntia versicolor* Engelm.), 16 (recorded as *Opuntia versicolor* Engelm.), 26 (genus, recorded as *Opuntia*), 27 (Pages 6, color photograph: Plates 6, 6A & 6B, Page 95), 28 (color photograph), 43 (012110 - *Cylindropuntia versicolor* (Engelm.) F.M. Knuth), 45 (color photograph), 46 (recorded as *Opuntia versicolor* Engelm., Page 585), 48 (genus, recorded as *Opuntia*), 58 (recorded as *Opuntia versicolor* Engelm.), 63 (012110 - color presentation), 77 (recorded as *Opuntia versicolor* Engelm., color photograph #15), 85 (012110 - color presentation), 115 (color presentation), 119 (recorded as *Opuntia versicolor* Engelm.), 127, **WTK** (March 9, 2007)*

Echinocactus wislizeni (see *Ferocactus wislizeni*)

***Echinocereus fasciculatus* (G. Engelmann ex B.D. Jackson) L.D. Benson: Pinkflower Hedgehog Cactus**

SYNONYMY: *Echinocereus fasciculatus* (G. Engelmann) L.D. Benson var. *fasciculatus*, *Echinocereus fendleri* (G. Engelmann) F. Sencke ex J.N. Haage var. *fasciculatus* (G. Engelmann ex B.D. Jackson) N.P. Taylor, *Echinocereus fendleri* (G. Engelmann) F. Sencke ex J.N. Haage var. *robustus* (R.H. Peebles) L.D. Benson, *Mammillaria fasciculata* G. Engelmann ex B.D. Jackson (possibly incorrectly

applied). COMMON NAMES: Bundle Hedgehog, Bundle Hedgehog Cactus, Bundle-spine Hedgehog, Magenta-flower Hedgehog Cactus, Pinkflower Hedgehog Cactus, Robust Hedgehog, Robust Hedgehog Cactus, Short-spine Strawberry Cactus, Strawberry Cactus. DESCRIPTION: Terrestrial perennial stem-succulent shrub (ascending to erect stems 2 to 18 inches in height and 1½ to 3 inches in width either single or in clusters of up to 30 stems, one plant was reported to have 150 stems); the stems are green or dark green; the spines often with zones of differing colors including black, gray, grayish-black-purplish, reddish-brown, whitish or yellowish turning gray with age; the flowers (2 to 3 inches in diameter) are cerise, lavender-pink, pale magenta, magenta, magenta-maroon, magenta-pink, magenta-purple, magenta-red, pink, pink-purple, purple, reddish-purple, rose-pink or white; the anthers are yellow; the stigma lobes are green, dark green or olive green; flowering generally takes place between late March and late June (additional records: one for early October, one for mid-October, one for late October, two for early November and one for early December); the mature fruits (¾ to 1¼ inches in length and ½ to 1 inch in diameter) are orange-red or bright red. HABITAT: Within the range of this species it has been reported from mountains; mesas; cliffs; canyons; canyon sides; bases of cliffs; buttes; knolls; ledges; ridges; along rocky and stony ridgetops; foothills; rocky, gravelly and sandy hills; rocky hilltops; rocky and sandy hillsides; rocky, stony and gravelly slopes; bajadas; rocky outcrops; amongst rocks and gravels, plains; gravelly flats; valley floors; along cobbly creeks; along and in washes; rocky and sandy banks, and floodplains growing in dry rocky, rocky-gravelly, stony, cobbly, gravelly and sandy ground, occurring from 1,800 to 6,300 feet in elevation in the woodland, scrub, grassland and desert scrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. The species, *Echinocereus fendleri*, was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop. The fruits are eaten by birds and other animals. *Echinocereus fasciculatus* is native to southwest-central and southern North America. *5, 6, 8, 12 (color photograph - recorded as *Echinocereus fasciculatus* (Engelm.) L. Benson var. *fasciculatus*, Pages 132-135), 15 (recorded as *Echinocereus fasciculatus* (Engelm.) L. Benson var. *fasciculatus*), 16 (recorded as *Echinocereus fasciculatus* (Engelm.) L. Benson), 27 (Page 81, color photograph: Plate 50, Page 104), 43 (012110), 45 (color photograph), 46 (recorded as *Echinocereus fendleri* (Engelm.) Rümpler var. *robustus* (Peebles) L. Benson, Page 572 and *Echinocereus fendleri* (Engelm.) Rümpler var. *robustus* (Peebles) L. Benson, Page 572), 48 (genus), 58 (recorded as *Echinocereus fasciculatus* (Engelm.) L. Benson var. *fasciculatus*), 63 (012110), 77 (recorded as *Echinocereus fasciculatus* (Engelm.) L. Benson, color photograph #64), 85 (012110 - color presentation), 115 (color presentation), 119 (species, recorded as *Echinocereus fendleri* (Engelm.) Rümpler), 127, **WTK** (March 9, 2007)*

***Ferocactus wislizeni* (G. Engelmann) N.L. Britton & J.N. Rose: Candy Barrelcactus**

SYNONYMY: *Echinocactus wislizeni* G. Engelmann. COMMON NAMES: Arizona Barrel Cactus, Barrel Cactus, Bisnaga, Biznaga, Biznaga de Agua (Spanish), Biznagre, Candy Barrel, Candy Barrel Cactus, Candy Barrelcactus, Compass Barrel, Compass Plant, Fish-hook Barrel, Fishhook Barrel Cactus, Fishhook Cactus, Southwest Barrel Cactus, Southwestern Barrel Cactus, Visnaga, Viznaga Hembra (Spanish), Wislizenus Barrel, Yellow-spined Barrel Cactus. DESCRIPTION: Terrestrial perennial stem-succulent shrub or tree (erect stem 1 to 11 feet in height and 8 to 40 inches in diameter); the stem is green or blue-gray-green; the central spines and larger radial spines are gray, dull pink, reddish or tan; the smaller radial spines are white; the flowers (1½ to 2½ inches in diameter) are orange, orange-yellow, orange-red, orange-yellow, parchment, pinkish-red, reddish, red-orange, yellow or yellow-orange; flowering generally takes place between mid-July and mid-October (additional records: one for early January, three for early March, five for mid-March, two for late March, one for early April, one for mid-April, one for late April and two for early June); the mature fruits (1¼ to 2 inches in length and 1 to 1½ inches in diameter) are greenish-brown, bright yellow or yellow-green and may remain on the plant until the next flowering period. HABITAT: Within the range of this species it has been reported from mountains; rocky mountainsides; mesas; rocky canyons; canyon walls; sandy canyon bottoms; bluffs; foothills; bouldery, rocky, gravelly and sandy hills; hillsides; rocky, cobbly and clayey-loamy slopes;

rocky, gravelly and sandy alluvial fans; bajadas; rocky outcrops; plains; rocky, gravelly and sandy flats; valley floors; along roadsides; arroyos; sandy bottoms of arroyos; along washes; rocky, gravelly and sandy margins of washes; floodplains, and mesquite bosques growing in dry desert pavement; bouldery, rocky, cobbly, gravelly and sandy ground, and sandy-clayey loam and clayey loam ground, occurring from 500 to 5,600 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food and/or beverage crop; it was also noted as having been used as a tool (the spines were heated and used to make fishing hooks). Fishhook Barrel Cacti are very slow to establish. A 4 year old plant may be no more than 1½ inches in height and 2 inches in width, and an 8 year old plant may be no more than 4¼ inches in height and 4¾ inches in width. The growth rate of propagated and cultivated barrel cacti is much faster. The life-span of Fishhook Barrel Cacti is reported to be from 50 to over 130 years of age. Some plants tend to lean to the south with age. Cristate forms have been reported. The fruits are eaten by Mule Deer (*Odocoileus hemionus*), Javelina (*Peccari tajacu*) and other animals, and the seeds are eaten by birds and rodents. *Ferocactus wislizeni* is native to southwest-central and southern North America. *5, 6, 12 (Pages 166-170, color photograph), 15, 16, 18, 26 (genus, color photograph of genus), 27 (Page 120, color photographs: Plates 60, 60A, 60B & 60C Pages 106), 28, 43 (063009 - *Ferocactus wislizeni* Britton & Rose), 45 (color photograph), 46 (Page 573), 48 (genus), 58, 63 (012210 - color presentation), 77 (color photograph #10), 85 (012210 - color presentation, also recorded as *Ferocactus wislizeni* var. *wislizeni*), 91, 115 (color presentation), 119, 127, **WTK** (March 9, 2007)*

Ferocactus wislizeni var. *wislizeni* (see footnote 85 under *Ferocactus wislizeni*)

Echinocereus fasciculatus var. *fasciculatus* (see *Echinocereus fasciculatus*)

Echinocereus fendleri var. *fasciculatus* (see *Echinocereus fasciculatus*)

Echinocereus fendleri var. *robustus* (see *Echinocereus fasciculatus*)

Lemaireocereus thurberi (see *Stenocereus thurberi*)

Mammillaria fasciculata (see *Echinocereus fasciculatus* and/or *Mammillaria thornberi*)

***Mammillaria grahamii* G. Engelmann: Graham's Nipple Cactus**

SYNONYMY: *Mammillaria grahamii* G. Engelmann var. *grahamii* G. Engelmann, *Mammillaria grahamii* G. Engelmann var. *oliviae* (C.R. Orcutt) L.D. Benson, *Mammillaria microcarpa* G. Engelmann, *Mammillaria oliviae* C.R. Orcutt. *Neomammillaria microcarpa* (G. Engelmann) N.L. Britton & J.N. Rose, *Neomammillaria milleri* N.L. Britton & J.N. Rose, *Neomammillaria oliviae* (C.R. Orcutt) N.L. Britton & J.N. Rose. COMMON NAMES: Arizona Fishhook, Arizona Fishhook Cactus, Biznaguita, Cabeza de Viejo Cekida, Cactus, Corkseed Cactus, Fishhook Cactus, Fishhook Mammillaria, Fishhook Pincushion, Graham Fishhook, Graham Nipple Cactus, Graham's Fishhook Cactus, Graham's Nipple Cactus, Graham Pincushion Cactus, Lizard Catcher, Nipple Cactus, Olive Pincushion, Pin-cushion Cactus. DESCRIPTION: Terrestrial perennial stem-succulent shrub (ascending to erect stems 1 to 12 inches in height and 1 to 3 inches in diameter, one plant was reported to be 1¼ inches in height and 1½ inches in width); the stems are gray-green or green; the central spines are black, golden-brown, purplish-brown or reddish; the radial spines are whitish; the flowers (½ to 1½ inches in diameter) may be lavender, pink, pink with a darker mid-stripe, pink-lavender, rose-pink, rose-purple or white, the anther are yellow; the stigma lobes are green; flowering generally takes place between mid-May and early August and one week after a heavy rains between mid-March and late September; the mature club-shaped fruits (1/2 to 1 1/8 inches in length and 3/16 to 1/2 inch in diameter) are carmine, bright orange, orange-red, bright red,

scarlet or yellow. HABITAT: Within the range of this species it has been reported from rocky mountains; sandy mountain slopes; rocky canyons; canyon bottoms; crevices in boulders and rocks; ridges; foothills; rocky and gravelly hills; rocky hillsides; rocky slopes; bajadas; rocky outcrops; amongst boulders and rocks; bases of boulders; protected clefts; gravelly and sandy flats; valley floors; along and in bouldery and sandy washes; edges of streams, and riparian areas often in the shade of other plants growing in dry bouldery, rocky, gravelly and sandy ground; gravelly loam ground; clay ground; silty ground, and humusy ground, occurring from 200 to 5,200 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop; it was also noted as having been used as a drug or medication. Birds and rodents feed on the fruits. *Mammillaria grahamii* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Mammillaria grahamii* Engelm., Pages 156 & 159-161; *Mammillaria grahamii* Engelm. var. *grahamii*, Pages 159-160; *Mammillaria grahamii* Engelm. var. *oliviae* (Orcutt) L. Benson, Pages 160-161, and *Mammillaria microcarpa* Engelm., Pages 152-153 & 156, color photographs), 15 (recorded as *Mammillaria grahamii* Engelm. var. *grahamii*; *Mammillaria grahamii* Engelm. var. *oliviae* (Orcutt) L. Benson, and *Mammillaria microcarpa* Engelm.), 16 (recorded as *Mammillaria microcarpa* Engelm.), 18 (genus), 27 (recorded as *Mammillaria grahamii*, Page 172, color photograph: Plate 94, Page 113; *Mammillaria grahamii* Engelm. var. *oliviae* (Orcutt) L. Benson, Pages 173, color photograph: Plate 95, Page 113, and *Mammillaria microcarpa* Engelm., Page 179, color photograph: Plate 99, Page 114), 28 (recorded as *Mammillaria microcarpa*, color photograph), 43 (012210), 45 (color photograph), 46 (recorded as *Mammillaria microcarpa* Engelm., Page 578 and *Mammillaria oliviae* Orcutt, Page 578), 48 (genus), 58 (recorded as *Mammillaria microcarpa* Engelm.), 63 (012210 - color presentation), 77 (color photograph #11), 85 (012210 - restricted distribution information, color presentation), 86 (recorded as *Mammillaria microcarpa*, color photograph), 115 (color presentation), 119 (recorded as *Neomammillaria microcarpa* (Engelm.) B. & R., *Neomammillaria milleri* B. & R.), 127, **WTK** (March 9, 2007)*

Mammillaria grahamii var. *grahamii* (see *Mammillaria grahamii*)

Mammillaria grahamii var. *oliviae* (see *Mammillaria grahamii*)

Mammillaria microcarpa (see *Mammillaria grahamii*)

Mammillaria oliviae (see *Mammillaria grahamii*)

Neomammillaria microcarpa (see *Mammillaria grahamii*)

Neomammillaria milleri (see *Mammillaria grahamii*)

Neomammillaria oliviae (see *Mammillaria grahamii*)

Opuntia bigelovii (see *Cylindropuntia bigelovii*)

Opuntia discata (see *Opuntia engelmannii* var. *engelmannii*)

***Opuntia engelmannii* J.F. Salm-Reifferscheid-Dyck ex G. Engelmann var. *engelmannii*: Cactus Apple**

SYNONYMY: *Opuntia discata* D. Griffiths, *Opuntia phaeacantha* G. Engelmann var. *discata* (D. Griffiths) L.D. Benson & D.L. Walkington. COMMON NAMES: Abrojo, Cactus Apple, Desert Pricklypear Cactus, Engelmann Prickly Pear, Engelmann's Prickly-pear, Engelmann Pricklypear, Flaming Pricklypear, Joconostle, Nopal, Prickly Pear, Vela de Coyote. DESCRIPTION: Terrestrial perennial stem-

succulent shrub (forms clumps 20 inches to 8 feet in height and 20 inches to 10 feet or more in width, one plant was reported as being 20 inches in height and 8¼ feet in width, one plant was reported as being 3 feet in height and 4½ feet in width, one plant was reported as being 3 feet in height and 6 to 12 feet in width, one plant was reported as being 3 feet in height and 8 feet in width, one plant was reported as being 40 inches in height and 79 inches in width, one plant was reported to be 40 inches in height and 10 feet in width); the paddle-shaped stems (8 to 16 inches in length and 6½ to 12 inches in width) are bluish-green, gray-green, green, dark green or yellow-green; the spines are brown-red, chalky-white, pale straw or pale yellow-brown usually with red or red-brown bases aging to black or gray; the glochids are yellow; the flowers (2¼ to 3½ in diameter) may be lemon-yellow, pink, pink-red, red-pink, rose-red, salmon, tannish-yellow, yellow, light yellow-orange, yellow-orange or yellow-peach turning to orange, orange-yellow or pink-orange with age; the anthers are yellow; the stigma lobes are lime green; flowering generally takes place between mid-March and late June (additional records: one for early January, two for mid-February, one for mid-July, two for mid-August, one for early September, six for mid-September, three for early October and one for late December); the mature fruits (also known as tunas are 2½ to 3¼ in length and 1¼ inches in diameter) are magenta-rose, purple, red or reddish-purple. HABITAT: Within the range of this species it has been reported from mountains; mountaintops; sandy mountainsides; bedrock mesas; edges of cliffs; canyons; canyon bottoms; talus slopes; ledges; ridges; rocky ridgetops; rocky hills; bouldery, rocky and gravelly hillsides; bouldery, rocky, rocky-gravelly and sandy slopes; bajadas; rocky outcrops; amongst boulders and rocks; lava beds; breaks; steppes; plains; rocky, gravelly and sandy and silty flats; basins; valley floors; along roadsides; along and in gravelly and gravelly-humusy arroyos; gullies; along streams; along creeks; creekbeds; along and in washes; along and in gravelly-sandy drainages; banks of rivers; benches; shelves; terraces; sandy floodplains; amongst mesquites; ditches, and gravelly-sandy and sandy riparian areas growing in dry bouldery, rocky, rocky-gravelly, rocky-sandy, shaley, gravelly, gravelly-sandy and sandy ground; clayey ground; silty ground, and gravelly humusy ground, occurring from 1,000 to 7,500 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. The species, *Opuntia engelmannii*, was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, beverage, cooking agent or paint crop; it was also noted as having been used as a tool, as a lubricant (var. *engelmannii*) and as a drug or medication. The flowers open around 8 AM and remaining open for one or two days, and may live to be 30 or more years of age. The juicy fruits (tunas) with edible pulp are fed on by many browsing animals, including Black Bear (*Ursus americanus amblyceps*), Coyote (*Canis latrans mearnsi*), Javelina (*Peccari tajacu sonoriensis*) and Desert Tortoise (*Gopherus agassizi*) among others, and birds. The plant provides cover for many desert animals. *Opuntia engelmannii* var. *engelmannii* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Opuntia phaeacantha* Engelm. var. *discata* (Griffiths) Benson & Walkington “This is the largest and, in especially southern Arizona, one of the best-known native prickly pears of the Southwestern Deserts of the United States. It is variable in habit of growth, shape and size of joints, and size and distribution of spines. It is almost always found growing with var. *major*, which has longer brown spines restricted largely to the upper part of the narrower joint. Almost everywhere there are intergrading forms with many character recombinations. Var. *discata* is rarely stable but apparently a fringe-population extreme tied in closely with the more abundant and wide-ranging var. *major*.”), Pages 99 & 101-103, color photograph), 15 (recorded as *Opuntia phaeacantha* Engelm. var. *discata* (Griffiths) Benson & Walkington), 16 (recorded as *Opuntia phaeacantha* Engelm. var. *discata* (Griffiths) L. Benson - “Rocky slopes and gravelly flats; common; intergrading with *O. p.* var. *major*.”), 26 (species), 27 (recorded as *Opuntia phaeacantha* Engelm. var. *discata* (Griffiths) L. Benson, Pages 53 & 99-100, color photographs: Plates 30 & 30A, Pages 99 & 100), 28 (color photograph), 43 (063009), 45 (species, color photograph), 46 (species, Page 583), 48 (genus), 58 (recorded as *Opuntia phaeacantha* Engelm. var. *discata* (Griffiths) Benson & Walk.), 63 (0123110 - color presentation), 77 (recorded as *Opuntia phaeacantha* var. *discata* (Griffiths) Benson & Walkington, color photograph #14 labeled as *Opuntia phaeacantha*), 85 (012310 - color presentation), 91 (recorded together with *Opuntia engelmannii* Salm-

Dyck. *Opuntia phaeacantha* var. *discata* (Griffiths) Benson & Walkington / *Opuntia phaeacantha* var. *major* Engelm.: “Both species are sympatric throughout much of their range and often can be found together.”), 115 (color presentation of the species), 119 (recorded as *Opuntia discata* Griffiths), 127 (variety *engelmannii* and species), **WTK** (March 9, 2007)*

Opuntia fulgida (see *Cylindropuntia fulgida* var. *fulgida*)

Opuntia fulgida var. *fulgida* (see *Cylindropuntia fulgida* var. *fulgida*)

Opuntia gilvescens (see *Opuntia phaeacantha*)

Opuntia leptocaulis (see *Cylindropuntia leptocaulis*)

***Opuntia macrocentra* G. Engelm. var. *macrocentra*: Purple Pricklypear**

SYNONYMY: *Opuntia violacea* G. Engelm. ex B.D. Jackson var. *macrocentra* (G. Engelm.) L.D. Benson; *Opuntia violacea* G. Engelm. ex B.D. Jackson var. *violacea*. COMMON NAMES: Black-spined Pricklypear, Duranzilla, Long-spined Pricklypear, Purple Pricklypear. DESCRIPTION: Terrestrial perennial stem-succulent shrub or tree (forms clumps to 2 to 5 feet in height and about as tall as wide); the stems (4 to 8 inches in length and 3 to 5 inches in width) are light blue, blue-green tinged with red, dark purple or red; the spines are black to reddish-brown; the flowers (2 to 3½ inches in diameter) are yellow with a red throat; flowering generally takes place between May and June; the fleshy fruits (1 to 2½ inches in length and ¾ to 1 inch in diameter) are purple, purplish-red or red. HABITAT: Within the range of this species it has been reported from rocky hills; rocky hillsides; slopes; bajadas; sand hills; plains; sandy flats; valley floors, and along washes growing in dry rocky, gravelly and sandy ground, occurring from 2,000 to 5,500 feet in elevation in the woodland, grassland and desert scrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. *Opuntia macrocentra* var. *macrocentra* is native to southwest-central and southern North America. *5, 6, 12 (*Opuntia violacea* var. *macrocentra* and *Opuntia violacea* var. *violacea*, Pages 91-93, color photograph, Page 94), 18 (species), 26 (genus), 27 (recorded as *Opuntia violacea* Engelm. var. *macrocentra* (Engelm.) L. Benson, Page 59 and *Opuntia violacea* Engelm. var. *violacea* L. Benson. Page 57, color photographs: Plates 33 & 33A, Page 100), 43 (062710), 45 (color photograph), 46 (Page 583), 48 (genus), 63 (062710 - color presentation), 85 (062710 - unable to access species information), 91, **WTK** (March 9, 2007, also recorded as *Opuntia gosseliniana*)*

***Opuntia phaeacantha* G. Engelm.: Tulip Pricklypear**

SYNONYMY: *Opuntia gilvescens* D. Griffiths, *Opuntia phaeacantha* G. Engelm. var. *major* G. Engelm., *Opuntia phaeacantha* G. Engelm. var. *phaeacantha*, *Opuntia phaeacantha* G. Engelm. var. *superbospina* (D. Griffiths) L.D. Benson. COMMON NAMES: Abrojo, Brown-spine Pricklypear, Brownspine Pricklypear, Brown-spined Prickly-pear, Desert Prickly-pear, Joconostle, Major Prickly-pear, Major Pricklypear, Mojave Prickly-pear, Mojave Pricklypear, New Mexico Prickly-pear, Nopal, Purple-fruit Prickly-pear, Sprawling Prickly Pear, Tulip Pricklypear, Vela de Coyote, Yellow Pricklypear, Yellow-spine Prickly-pear. DESCRIPTION: Terrestrial perennial stem-succulent shrub (10 inches to 7 feet in height and 3 to 10 feet in width sometimes forming clumps up to 75 feet in width, sometimes developing a definite trunk, one plant was reported to be 10 inches in height and 40 inches in width, one plant was reported to be 1 foot in height and 3 feet in width, one plant was reported to be 14 inches in height and 52 inches in width, plants were reported that were 16 inches in height and 40 inches in width, one plant was reported to be 16 inches in height and 48 inches in width, one plant was reported to be 16 inches in height and 60 inches in width, one plant was reported to be 18 inches in height and 8 to 10 feet in width, one plant was reported to be 20 inches in height and 13 feet in width, one plant was reported to be 2 feet in height and 5 to 6 feet in width, one plant was reported to be 30 inches in height and 5 feet in width, plants were reported to be 3 feet in height and 4 to 10 feet in width); the paddle-

shaped stems (4 to 10 inches in length and 3 to 8 inches in width) may be bluish-green, gray-brown, gray-green, dull green, green, dark green, greenish-yellow, purple, reddish or yellow-gray-green; the spines are blackish, brown, charcoal, gray, reddish, red-brown, white or yellow; the glochids are golden, reddish-brown or tan; the flowers (1½ to 3 inches in diameter) may be golden-apricot (with yellow-green mid-stripes), orange, orange-yellow, pink, pink-purple, red, red-pink, pale yellow, yellow (with an orange or red center or brown, greenish, greenish-brown or red mid-stripes) or yellow-orange aging to red-orange; the anthers are yellow; the stigma lobes are green or yellow-green; flowering generally takes place between mid-March to early July (additional records: one for early January, one for late January, one for early February, one for late July, three for mid-August, two for late August, one for late September and one for early October); the mature pear-shaped fruits (1¼ to 3½ inches in length and 1 to 1¼ inches in width) are maroon, purple, purple-red, red, dark red, red-brown or wine-red. HABITAT: Within the range of this species it has been reported from mountains; mesas; plateaus; cliffs; canyons; canyon bottoms; rocky rincons; talus slopes; bases of cliffs; bluffs; rocky-gravelly-sandy buttes; knolls; rocky ledges; ridges; ridgetops; foothills; rocky and gravelly hills; cobbly and sandy hilltops; bouldery, rocky, gravelly and gravelly-sandy-loamy hillsides; bouldery, rocky, rocky-gravelly, gravelly, sandy and silty slopes; gravelly bajadas; rocky outcrops, amongst rocks; on boulders and rocks; lava beds; blow-sand; prairies; sandy llanos; plains; rocky, cindery and sandy flats; valleys; along sandy roadsides; in rocky and sandy arroyos; bottoms of arroyos; draws; springs; along creeks; along and in sandy riverbeds; along gravelly washes; sandy drainages; silty-loamy and silty-clayey-loamy dry lakebeds; along (sandy) banks of rivers; cobbly-sandy-silty and gravelly-sandy terraces; sandy-loamy bottomlands; sandy floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry bouldery; rocky, rocky-gravelly, rocky-gravelly-sandy, shaley, cobbly, cindery, gravelly, gravelly-sandy and sandy ground; gravelly-sandy loam, sandy loam, sandy-clayey loam and silty loam and silty-clayey loam ground; gravelly-sandy clay ground; cobbly-sandy silty and silty ground, and humusy ground, occurring from 800 to 7,900 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, beverage and/or cooking agent crop; it was also noted as having been used for tools, in pottery making and as a drug or medication. This plant provides cover for many desert animals. Deer, Javelina (*Peccari tajacu sonoriensis*) and rodents feed on the stems, and the fruits are eaten by deer, grasshoppers, Javelina and other desert animals (including grasshoppers). Cristate forms have been reported. The change in nomenclature in USDA NRCS has not been recognized in BONAP, varieties remain as varieties of *Opuntia phaeacantha* (accessed 041806). *Opuntia phaeacantha* is native to southwest-central and southern North America. *5, 6, 12 (recorded as *Opuntia phaeacantha* Engelm., Pages 95-101; *Opuntia phaeacantha* Engelm. var. *major* Engelm., Pages 99-101, and *Opuntia phaeacantha* Engelm. var. *phaeacantha*, Pages 97-98), 15 (recorded as *Opuntia phaeacantha* var. *major* Engelm., color photograph on Page 77 includes habitat and associated species), 16 (recorded as *Opuntia phaeacantha* Engelm. var. *major* Engelm.), 26 (color photograph), 27 (recorded as *Opuntia phaeacantha* Engelm., Pages 50, color photograph: Plate 28, Page 99; *Opuntia phaeacantha* Engelm. var. *major* Engelm., Pages 51, color photograph: Plate 29, Page 99, and *Opuntia phaeacantha* Engelm. var. *superbospina* (Griffith) L. Benson, Pages 54, color photograph: Plate 31, Page 100), 43 (070109), 45 (color photograph), 46 (recorded as *Opuntia phaeacantha* Engelm., Page 583 and *Opuntia gilvescens* Griffiths, Page 583), 48 (genus - recorded as *Opuntia*), 58 (recorded as *Opuntia phaeacantha* Engelm. var. *major* Engelm.), 63 (012310 - color presentation), 77 (recorded as *Opuntia phaeacantha* Engelm. var. *major* Engelm., color photograph #14 labeled as *Opuntia phaeacantha*), 85 (012310 - color presentation), 91 (recorded together with *Opuntia engelmannii* Salm-Dyck. (*Opuntia phaeacantha* var. *discata* (Griffiths) Benson & Walkington) / *Opuntia phaeacantha* var. *major* Engelm. - “Both species are sympatric throughout much of their range and often can be found together.”), 119, 127, **WTK** (March 9, 2007, recorded as *Opuntia phaeacantha* var. *major*)*

Opuntia phaeacantha var. *discata* (see *Opuntia engelmannii* var. *engelmannii*)

Opuntia phaeacantha var. *major* (see *Opuntia phaeacantha*)

Opuntia phaeacantha var. *phaeacantha* (see *Opuntia phaeacantha*)

Opuntia phaeacantha var. *superbospina* (see *Opuntia phaeacantha*)

Opuntia spinosior (see *Cylindropuntia spinosior*)

Opuntia versicolor (see *Cylindropuntia versicolor*)

Opuntia violacea var. *macrocentra* (see *Opuntia macrocentra* var. *macrocentra*)

Opuntia violacea var. *violacea* (see *Opuntia macrocentra* var. *macrocentra*)

***Stenocereus thurberi* (G. Engelmann) F. Buxbaum: Organpipe Cactus**

SYNONYMY: *Cereus thurberi* G. Engelmann, *Lemaireocereus thurberi* (G. Engelmann) N.L. Britton & J.N. Rose. COMMON NAMES: Marismena, Mehuelé (Spanish), Organ Pipe Cactus, Organpipe Cactus, Organo, Órgano Marismeña (Spanish), Organpipe Cactus, Pitahaya, Pitahaya Dulce (Spanish for Sweet Cactus Fruit), Pitayo Dulce (Spanish). DESCRIPTION: Terrestrial perennial stem-succulent shrub or tree (erect stems 5 to over 40 feet in height and 5 to 12 or more inches in width with 10 to 20 stems with larger specimens having 40 to 45 stems clustering to 6 to 18 feet in diameter); the stems are gray-green, green, yellow-green or yellowish; the spines are black, brownish, gray or reddish-brown aging to gray; the flowers (1½ to 2½ inches in width) may be brownish-green, greenish-white, pale lavender, lavender with white margins, pink, purple, white or white with a pale pink center; the anthers are yellowish; the stigma lobes are white; flowering generally takes place between mid-May and mid-June (additional records: one for early March, one for mid-April, one for mid-July and one for late July), the ripe fruits are red or reddish with green scales. HABITAT: Within the range of this species it has been reported from mountains; mesas; rocky canyons; rocky canyonsides; ledges; ridges; foothills; rocky and sandy hills; rocky and rocky-gravelly hillsides; rocky slopes; gravelly bajadas; amongst rocks; sand dunes; sandy plains; flats; valley floors; coastal bluffs; sandy and powdery coastal plains, and rocky benches growing in dry rocky, rocky-gravelly, stony, gravelly, sandy ground and powdery ground, occurring from sea level to 3,700 feet elevation in the scrub, grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, but it is sensitive to frosts. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, fodder, beverage, cooking agent and/or fiber crop; it was also noted as having been used as a ceremonial item, for tools (in the making of torches) and in making paint. It has been estimated that plants 10 to 15 feet in height may be between 50 and 75 years of age, Cristate forms have been reported. The flowers reportedly open after sunset and close the following morning. The Broad-billed Hummingbird (*Cyanthus latirostris*), Costa's Hummingbird (*Calypte costae*) and Rufous Hummingbird (*Selasphorus rufus*) have been observed visiting the flowers. The flowers are pollinated by bats, including Southern Long-nosed Bat (*Leptonycteris curasoae yerbabuena*), and bees. The ripe fruits are eaten by ants, bats, Coyotes (*Canis latrans mearnsi*), Desert Bighorn Sheep (*Ovis canadensis mexicana*) and White-winged Doves (*Zenaida asiatica*). The candy, Pitahaya Dulce, is made by cooking the fruits of the Organ Pipe Cactus with those of Prickly Pear Cacti. This is a PERIPHERAL - DISJUNCT POPULATION. *Stenocereus thurberi* is native to southwest-central and southern North America. *5, 6, 8, 12 (recorded as *Cereus thurberi* Engelm., Pages 111 & 113-114), 13 (color photograph, in habitat with associated species Plate D.1), 18, 27 (recorded as *Cereus thurberi* Engelm., Page 63, color photographs: Plate 38, Page 102 and Page 93 color photograph of plant in habitat), 28 (recorded as color photograph, recorded as *Cereus thurberi*), 43 (070310), 45 (color photograph, recorded as *Stenocereus thurberi*), 46 (recorded as *Lemaireocereus*

thurberi (Engelm.) Britt. & Rose, Pages 569-570), 48 (recorded as *Cereus thurberi*), 53 (*Cereus thurberi* Engelm.), 63 (070310 - color presentation including habitat), **85** (070310 - color presentation including habitat, unable to access species information), 91 (recorded as *Stenocereus thurberi* (Engelm.) Buxb.), 119 (recorded as *Lemaireocereus thurberi* (Engelm.) B. & R.), 127*

Convolvulaceae: The Morning-glory Family

***Ipomoea* C. Linnaeus: Morning-glory**

COMMON NAME: Morning-glory. *43 (052010), 46 (Pages 675-678), 63 (080409), **WTK** (March 9, 2007)*

Euphorbiaceae: The Spurge Family

***Chamaesyce florida* (G. Engelmann) C.F. Millspaugh: Chiricahua Mountain Sandmat**

SYNONYMY: *Euphorbia florida* G. Engelmann. COMMON NAMES: Chiricahua Mountain Sandmat, Florida Spurge, Golondrina, Spurge. DESCRIPTION: Terrestrial annual forb/herb (ascending stems 1 to 18 inches in height); the stems are pink-tan; the leaves are green; the flower-like cups have green glands (centers) and white (aging rose), white-pink or white with pinkish tips petaloid appendages; flowering generally takes place between mid-July and early November (additional records: two for early January, one for late June and two for mid-December). HABITAT: Within the range of this species it has been reported from mountains; mesas; rocky canyons; canyon walls; sandy canyon bottoms; chasms; sandy ridgetops; rocky foothills; rocky and sandy hills; rocky hillsides; rocky, rocky-gravelly, rocky-sandy, gravelly-loamy and sandy-loamy slopes; bajadas; dunes; plains; gravelly and sandy flats; basins; valley floors; coastal dunes; along rocky-sandy, gravelly-clayey and sandy roadsides; arroyos; along and in streambeds; along and in gravelly and sandy washes; gravelly-clayey depressions; along sandy banks of arroyos, rivers and washes; bottomlands; floodplains; edges of stock tanks; sandy riparian areas, and disturbed areas growing in wet, moist and dry rocky, rocky-gravelly, rocky-sandy, gravelly and sandy ground; gravelly loam, gravelly-clayey loam and sandy loam ground, and gravelly clay ground, occurring from sea level to 5,000 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: The stems have a milky sap. *Chamaesyce florida* is native to southwest-central and southern North America. *5, 6, 15, 16 (recorded as *Euphorbia florida* Engelm.), **18** (“All euphorbias have milky white sap that is irritating on contact or toxic, if ingested, (degree of irritation or toxicity varies, depending on the species).”), 43 (020310), 46 (recorded as *Euphorbia florida* Engelm., Page 518), 58, 63 (020310), **68** (see: Poisonous Properties of Spurges, Page 202), 77 (recorded as *Euphorbia florida* Engelm.), **80** (**Species of the genus *Euphorbia* are considered to be Secondary Poisonous Range Plants.** “The milky juice of Spurge is considered poisonous. Plants may cause skin irritation, diarrhea, photosensitization, and cyanogenetic poisoning. Cattle, horses, sheep, and humans may be affected. The green plants are generally unpalatable but the dried plants in hay are more palatable and remain toxic. ... Poisoning may be prevented by keeping animals off areas heavily infested with spurge when other desirable feed is unavailable, and by not feeding contaminated hay. Range improvement will both reduce spurge infestations through grass competition, and decrease consumption by making more desirable forage available.” See text for additional information.), **85** (020310 - color presentation), **86** (“Most members of the family (Euphorbiaceae) are poisonous, and their milky sap will irritate the membranes of the eyes and mouth.”), 115 (color presentation)*

Euphorbia florida (see *Chamaesyce florida*)

***Jatropha cardiophylla* (J. Torrey) J. Müller Argoviensis: Sangre de Cristo**

COMMON NAMES: Limberbush, Matacora, Nettlespurge, Sangre de Cristo, Sangre-de-Cristo, Sangre-de-drago, Sangregrado, Sangregrado, Sangringada, Torote. DESCRIPTION: Terrestrial perennial deciduous, semi-succulent shrub (1 to 7 feet in height); the flexible stems are basally branches; the bark is reddish; the leaves shiny green; the small bell-shaped flowers may be cream-white, pink, white or yellow; flowering generally takes place between mid-July and late September. HABITAT: Within the range of this species it has been reported from mountains; mesas; rocky canyons; canyon bottoms; foothills; rocky hills; rocky hillsides; rocky slopes; rocky and gravelly bajadas; boulderfields; gravelly plains; gravelly-sandy flats; basins; valley floors; rocky roadsides; within sandy arroyos; bottoms of arroyos; cobbly and cobbly-gravelly-loamy draws; along and in sandy washes; margins of washes; floodplains; riparian areas, and disturbed areas growing in dry bouldery, rocky, cobbly, gravelly, gravelly-sandy and sandy ground and cobbly-gravelly loam and gravelly loam ground, occurring from 100 to 4,800 feet in elevation in the scrub, grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial fiber crop for use in making baskets. The shiny heart-shaped emerald green leaves appear around the time of the first rains and then provide color when the leaves turn gold in the fall. *Jatropha cardiophylla* is native to southwest-central and southern North America. *5, 6, 13 (color photograph), 15, 16, 43 (020510), 45 (color photograph), 46 (Page 509), 48, 58, 63 (020510), 77, 80 (Species of the genus *Jatropha* are considered to be Rarely Poisonous and Suspected Poisonous Range Plants. "Seeds of several species of *Jatropha* are toxic to humans and livestock but no poisoning has been reported from Arizona."), 85 (020510 - color presentation), 91, 115 (color presentation), 127, **WTK** (March 9, 2007)*

Fabaceae (Leguminosae): The Pea Family

***Acacia constricta* G. Bentham: Whitethorn Acacia**

SYNONYMY: *Vachellia constricta* (G. Bentham) D.S. Seigler & J.E. Ebinger. COMMON NAMES: All-thorn Acacia, Chaparo Prieta, Chaparro Prieto, Common Whitethorn, Garabato, Gidag (Tohono O'odham), Gigantillo, Huisache, Largoncillo, Mescat Acacia, Twinthorn Acacia, Vara Prieta, Vinorama, Whitethorn Acacia, White Thorn, Yellow Cat Claw. DESCRIPTION: Terrestrial perennial deciduous (drought and cold) shrub or tree (1 to 20 feet in height with crowns to about the same in width, one plant was described as being 8 feet in height with a crown 8 feet in width); the bark may be light gray, mahogany or nearly black; the stems may be red; the spines on the branches and stems are gray or white; the small pinnate leaves are green; the small flowers have been described as being golden, golden-yellow, orange-yellow, light yellow, yellow or yellowish-orange; flowering generally takes place between late March and late October (additional records: two for early March and one for late December); the seedpods are brown, purple-red, reddish or rusty-brown. HABITAT: Within the range of this species it has been reported from mountains; mesas; cliffs; canyons; canyon sides; sandy canyon bottoms; sandy ridges; foothills; rocky and gravelly hills; bouldery hilltops; rocky and gravelly hillsides; escarpments; rocky, rocky-clayey-loamy and clayey-loamy slopes; gravelly bajadas; rocky outcrops; amongst boulders; sandy-loamy plains; gravelly flats; valley floors; coastal plains; along rocky, rocky-gravelly-loamy, rocky-gravelly-clayey loam, rocky-clayey-loamy, gravelly, gravelly-sandy, gravelly-sandy-loamy, gravelly-sandy-clayey-loamy, gravelly-loamy, gravelly-clayey loam and sandy roadsides; along and in rocky arroyos; bottoms of arroyos; rocky gulches; along streambeds; creeks; along and in sandy creekbeds; rivers; along and in gravelly, gravelly-sandy, sandy and silty-clayey washes; drainages; swales; along (gravelly-sandy and sandy) banks of streams, creeks, rivers and washes; along edges of washes; (rocky) margins of arroyos and washes; mudflats; benches; alluvial terraces; sandy bottomlands; floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry bouldery, rocky, rocky-sandy, gravelly, gravelly-sandy and sandy ground; rocky-gravelly loam, rocky-gravelly-clayey loam, rocky-clayey loam, gravelly loam, gravelly-sandy loam, gravelly-sandy-clayey loam, gravelly-clayey loam, sandy loam, sandy-clayey loam, clayey loam and loam ground, and silty clay ground, occurring

from 1,100 to 6,500 feet (infrequently as low as 500 feet and as high as 9,200 feet) in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, plants may live to be more than 72 years of age and the flowers may be fragrant. Whitethorn Acacia is used for food (but not extensively) by the Desert Mule Deer (*Odocoileus hemionus*) and Scaled Quail (*Callipepla squamata*), Merriam's Kangaroo Rats (*Dipodomys merriami*), Bailey's Pocket Mice (*Chaetodipus baileyi*) and Rock Pocket Mice (*Chaetodipus intermedius*) as well as a variety of other birds and mammals feed on the seeds. *Acacia constricta* is native to southwest-central and southern North America. *5, 6, 13 (color photograph), 15, 16, 18, 26 (color photograph), 28 (color photograph), 43 (080409), 46 (Page 399), 48, 53 (note under *Acacia farnesiana*), 63 (020710 - color presentation), 68, 77, 80 (This species is listed as a Major Poisonous Range Plant. "The plants are high in cyanide forming-compounds and have been reported to cause death of cattle in Arizona. In general, the plants are not palatable to livestock although the pods are grazed. However, in the fall of the year at or near frost time, when the range grasses become less palatable, cattle may eat heavily of these plants and death is likely to result. ... Animals should be removed from heavily infested areas during the early frost period or considerable death losses may occur." See text for additional information.), 85 (020710 - color presentation), 91, 115 (color presentation), 134, **WTK** (March 9, 2007)*

***Acacia constricta* G. Bentham var. *constricta*: Whitethorn Acacia**

SYNONYMY: (for *Acacia constricta*: *Vachellia constricta* (G. Bentham) D.S. Seigler & J.E. Ebinger). COMMON NAMES: All-thorn Acacia, Chaparo Prieta, Chaparro Prieto, Common Whitethorn, Garabato, Gidag (species, Tohono O'odham), Gigantillo, Huisache, Largoncillo, Mescat Acacia, Twinthorn Acacia, Vara Prieta, Vinorama, Whitethorn Acacia, White Thorn, Yellow Cat Claw. DESCRIPTION: Terrestrial perennial deciduous (drought and cold) shrub or tree (1 to 20 feet in height with crowns to about the same in width, one plant was described as being 8 feet in height with a crown 8 feet in width); the bark may be light gray, mahogany or nearly black; the stems may be red; the spines on the branches and stems are gray or white; the small pinnate leaves are green; the spines are gray or white; the small flowers are golden, golden-yellow, orange-yellow or yellow; flowering generally takes place between late March and late October (additional records: two for early March and one for late December); the seedpods are brown, purple-red or reddish. HABITAT: Within the range of this species it has been reported from mountains; mesas; canyons; sandy canyon bottoms; sandy ridges; foothills; rocky and gravelly hills; bouldery hilltops; rocky and gravelly hillsides; rocky and rocky-clayey-loamy slopes; gravelly bajadas; rock outcrops; amongst boulders; sandy-loamy plains; gravelly flats; valleys; coastal plains; along rocky, rocky-gravelly-loamy, rocky-clayey-loamy, gravelly, gravelly-sandy, gravelly-sandy-loamy, gravelly-sandy-clayey-loamy, gravelly-loamy, gravelly-clayey-loamy and sandy roadsides; along and in rocky arroyos; bottoms of arroyos; rocky gulches; along streambeds; creeks; along and in sandy creekbeds; gravelly-sandy rivers; along and in gravelly, gravelly-sandy, sandy and silty-clayey washes; drainage ways; (gravelly-sandy and sandy) banks of streams, creeks, rivers and washes; along edges of washes; (rocky) margins of arroyos and washes; benches; terraces; sandy bottomlands; floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry bouldery, rocky, rocky-sandy, gravelly, gravelly-sandy and sandy ground; rocky-gravelly loam, rocky-clayey loam, gravelly loam, gravelly-sandy loam, gravelly-sandy-clayey loam, gravelly clayey loam, sandy loam and loam ground, and silty clay ground, occurring from 1,200 to 6,500 feet (infrequently to as low as 500 feet and to as high as 9,200 feet) in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, plants may live to be more than 72 years of age and the flowers may be fragrant. Whitethorn Acacia is used for food (but not extensively) by the Desert Mule Deer (*Odocoileus hemionus*) and Scaled Quail (*Callipepla squamata*), Merriam's Kangaroo Rats (*Dipodomys merriami*), Bailey's Pocket Mice (*Chaetodipus baileyi*) and Rock Pocket Mice (*Chaetodipus intermedius*) as well as a variety of other birds and mammals feed on the seeds. *Acacia constricta* var. *constricta* is native to southwest-central and southern North America. *5, 6, 13, 26 (color photograph of species), 28 (color photograph of species), 43

(080409), 46 (species, Page 399), 48 (species), 53 (species note under *Acacia farnesiana*), 63 (020710), 68, 80 (The species, *Acacia constricta*, is listed as a Major Poisonous Range Plant. “The plants are high in cyanide forming-compounds and have been reported to cause death of cattle in Arizona. In general, the plants are not palatable to livestock although the pods are grazed. However, in the fall of the year at or near frost time, when the range grasses become less palatable, cattle may eat heavily of these plants and death is likely to result. ... Animals should be removed from heavily infested areas during the early frost period or considerable death losses may occur.” See text for additional information.), 85 (020710), 91, 115 (color presentation of the species), 134, **WTK** (March 9, 2007)*

***Acacia constricta* G. Bentham var. *paucispina* E.O. Wooton & P.C. Standley: Whitethorn Acacia**

SYNONYMY: (for *Acacia constricta*: *Vachellia constricta* (G. Bentham) D.S. Seigler & J.E. Ebinger). COMMON NAMES: Chaparro Prieto, Common Whitethorn, Garabato, Gidag (species, Tohono O’odham), Gigantillo, Huisache, Largoncillo, Mescat Acacia, Vara Prieta, Vinorama, Whitethorn Acacia, White Thorn. DESCRIPTION: Terrestrial perennial deciduous (drought and cold) shrub or tree (2 to 18 feet in height with a crown to about the same in width); the bark may be light gray, mahogany or nearly black; the stems may be red; the small fragrant flowers are orange-yellow or yellow; flowering generally takes place between late March and late October (flowering records for early June, mid-June and late July); the seedpods are brown, purple-red or reddish. HABITAT: Within the range of this species it has been reported from mesas; canyons; ridges; gravelly hills; bouldery hilltops; rocky slopes; bajadas; amongst boulders; gravelly flats; rocky-gravelly-loamy and gravelly-loamy roadsides; along arroyos; along washes; swales; floodplains, and riparian areas growing in dry bouldery, rocky and gravelly ground and rocky-gravelly loam and gravelly loam ground, occurring from 2,000 to 3,700 feet (infrequently to 9,000 feet) in elevation in the forest, grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, plants may live to be more than 72 years of age and the flowers may be fragrant. This variety is spineless. *Acacia constricta* var. *paucispina* is native to southwest-central and southern North America. *5, 6, 13, 26 (color photograph of species), 28 (color photograph of species), 43 (080409), 46 (Page 399), 48 (species), 53 (species note under *Acacia farnesiana*), 63 (020710), 68, 80 (The species, *Acacia constricta*, is listed as a Major Poisonous Range Plant. “The plants are high in cyanide forming-compounds and have been reported to cause death of cattle in Arizona. In general, the plants are not palatable to livestock although the pods are grazed. However, in the fall of the year at or near frost time, when the range grasses become less palatable, cattle may eat heavily of these plants and death is likely to result. ... Animals should be removed from heavily infested areas during the early frost period or considerable death losses may occur.” See text for additional information.), 85 (020710), 91, 115 (color presentation of the species), 134, **WTK** (March 9, 2007)*

***Acacia greggii* A. Gray (var. *greggii* is the variety reported as occurring in Arizona): Catclaw Acacia**

SYNONYMY: (for *A.g.* var. *greggii*: *Acacia greggii* A. Gray var. *arizonica* D. Isely). COMMON NAMES: Acacia, Algarroba, Arizona Acacia (for var. *greggii*), Cat Claw, Cat Claw Acacia, Catclaw, Catclaw Acacia, Cat’s-claw, Devil’s Catclaw, Devil’s Claw, Devil’s-claw, Devilsclaw, Devil’s-claw Acacia, Di:s (Seri), Gatuno, Gregg Catclaw, Gregg’s Acacia, Tearblanket, Tepame, Tesota, Texas Catclaw, Texas Mimosa, Texas-mimosa, Una de Gato, Wait-a-minute, Wait-a-minute Bush, Wright Acacia (for var. *wrightii*). DESCRIPTION: Terrestrial perennial winter-deciduous shrub or tree (40 inches to 35 feet in height with a broad crown, one plant was reported as being 6½ feet in height with a crown 10 feet in width, one plant was reported as being 13 feet in height with a crown 16½ feet in width); the bark is gray-black or red-brown; the leaves are gray-green or green; the flowers may be cream, cream-white, cream-yellow, green, greenish-yellow, lemon-yellow, dull white, white, pale yellow, yellow, yellow-cream or yellow-green in catkins; flowering generally takes place between early March and early August (additional records: two for late August, one for mid-September, two for late September, one for early October, three for mid-October, one for early November, one for mid-November, one for early December and one for late December); the mature fruits (straight or twisted pods) are brown or brownish-red. HABITAT: Within the range of this species it has been reported from rocky mountains; mountainsides;

gravelly mesas; rocky canyons; rocky and sandy canyon bottoms; gorges; rocky bluffs; rocky and sandy ridges; ridgetops; foothills; rocky hills; gravelly hilltops; rocky, gravelly and gravelly-loamy hillsides; bedrock, rocky, rocky-gravelly-loamy, gravelly, gravelly-sandy and sandy slopes; alluvial fans; bajadas; amongst boulders; debris flows; plains; sandy flats; basins; valley floors; loamy valley bottoms; coastal plains; along gravelly-sandy, gravelly-sandy-clayey-loamy and sandy roadsides; along and in arroyos; bottoms of arroyos; draws; ravines; seeps; springs; along streams; along creeks; along sandy and sandy-silty creekbeds; along rivers; along and in rocky, rocky-sandy, gravelly, gravelly-sandy and sandy washes; within drainages; along drainage ways; along (rocky, gravelly-sandy, gravelly-silty, sandy and sandy-silty) banks of arroyos, streams, creeks, rivers and washes; along (sandy) edges of arroyos, creeks and washes; margins of washes; shorelines; sand bars; shelves; gravelly-sandy and sandy terraces; sandy bottomlands; lowlands; sandy-loamy floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry bouldery, rocky, rocky-sandy, gravelly, gravelly-sandy and sandy ground; rocky-gravelly loam, gravelly-sandy-clayey loam, gravelly loam, sandy loam, clayey loam and loam ground; gravelly clay, sandy clay and clay ground, and gravelly silty and sandy silty ground, occurring from slightly above sea level to 6,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat; the flowers are fragrant, it may live to be up to 120 years of age. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, fodder and/or fiber crop; it was also noted as having been used as a fuel, tool and for making perfumed sachets. Catclaw Acacia provides food, shelter, protection, shade, nesting sites, roosting sites and nesting material to a wide variety of species of wildlife. *Acacia greggii* is native to southwest-central and southern North America. *5, 6, 13 (color photograph), 15, 18, 26 (color photograph), 28 (color photograph), 43 (020710), 46 (“This is probably the most heartily disliked plant in the state, the sharp, strong prickles tearing the cloths and lacerating the flesh.”, Page 398), 48 (“A good honey plant but a poisonous weed on range lands.”), 52, 53, 58, 63 (020710 - color presentation), 77, 80 (**This species is listed as a Secondary Poisonous Range Plant.** “Plants contain cyanide-forming compounds and symptoms are typical of cyanide poisoning. The new foliage is relished by cattle in the early spring. It also may be grazed considerably during dry seasons or drouth periods when other feed is short. Plants are most dangerous in the fall during first frosts. Cattle are most often poisoned, but losses in Arizona are not heavy. Poisoning may be prevented by deferring heavily infested areas during the early frost periods.” See text for additional information.), 85 (020710 - color presentation), 91, 115 (color presentation), 127*

***Acacia greggii* A. Gray var. *greggii*: Catclaw Acacia**

SYNONYMY: *Acacia greggii* A. Gray var. *arizonica* D. Isely. COMMON NAMES: Acacia, Algarroba, Arizona Acacia (applied to var. *greggii*), Cat Claw, Cat Claw Acacia, Catclaw, Catclaw Acacia, Cat's-claw, Devil's Catclaw, Devil's Claw, Devil's-claw, Devilsclaw, Di:s (Seri), Gatuno, Gregg Catclaw, Gregg's Acacia, Tearblanket, Tepame, Tesota, Texas Catclaw, Texas Mimosa, Una de Gato, Wait-a-minute, Wait-a-minute Bush. DESCRIPTION: Terrestrial perennial winter-deciduous shrub or tree (40 inches to 25 feet in height with a broad crown); the bark is gray-black or red-brown; the leaves are gray-green or green; the flowers may be cream, cream-white, cream-yellow, green, greenish-yellow, lemon-yellow, white, yellow, yellow-cream or yellow-green in catkins; flowering generally takes place between early March and mid-July (additional record: one for mid-October); the mature fruits (straight or twisted pods) are brown or brownish-red. HABITAT: Within the range of this species it has been reported from mountains; mountainsides; mesas; canyons; rocky and sandy canyon bottoms; rocky bluffs; rocky and sandy ridges; ridgetops; hillsides; rocky, rocky-clayey-loamy, sandy and loamy slopes; amongst boulders; alluvial fans; sandy flats; valley floors; gravelly-sandy-clayey-loamy and sandy roadsides; sandy edges of arroyos; draws; ravines; along streams; along creeks; along rivers; along gravelly and sandy washes; within drainages; along banks of rivers and washes; along edges of washes; margins of arroyos; floodplains; mesquite bosques, and riparian areas growing in dry bouldery, rocky, rocky-sandy, gravelly, gravelly-sandy and sandy ground; rocky-gravelly loam, rocky-clayey loam, gravelly-sandy-clayey loam, sandy loam and clayey loam ground, and gravelly clay ground, occurring from slightly above sea level to

5,300 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat; the flowers are fragrant, it may live to be up to 120 years of age. The species, *Acacia greggii*, was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, fodder and/or fiber crop; it was also noted as having been used as a fuel, tool and for making perfumed sachets. Catclaw Acacia provides food, shelter, protection, shade, nesting sites, roosting sites and nesting material to a wide variety of species of wildlife. *Acacia greggii* var. *greggii* is native to southwest-central and southern North America. *5, 6, 13 (color photograph of species, species), 16 (recorded as *Acacia greggii* Gray var. *arizonica* Isely), 18 (species), 26 (species, color photograph of species), 28 (species, color photograph of species), 43 (020710), 46 (species, “This is probably the most heartily disliked plant in the state, the sharp, strong prickles tearing the clothes and lacerating the flesh.”), 48 (species - “A good honey plant but a poisonous weed on range lands.”, Page 398), 52 (species) 53, (species), 63 (020710), 80 (The species is listed as a Secondary Poisonous Range Plant. “Plants contain cyanide-forming compounds and symptoms are typical of cyanide poisoning. The new foliage is relished by cattle in the early spring. It also may be grazed considerably during dry seasons or drouth periods when other feed is short. Plants are most dangerous in the fall during first frosts. Cattle are most often poisoned, but losses in Arizona are not heavy. Poisoning may be prevented by deferring heavily infested areas during the early frost periods.” See text for additional information.), 85 (020710), 91 (species), 115 (color presentation of the species), 127 (species), **WTK** (March 9, 2007)*

Acacia greggii var. *arizonica* (see *Acacia greggii* var. *greggii*)

***Calliandra eriophylla* G. Bentham: Fairyduster**

SYNONYMY: *Calliandra eriophylla* G. Bentham var. *erriophylla*. COMMON NAMES: Cabelleto de Angel, Cabeza Angel, Desert Fairy-duster, Fairy Duster, Fairy-duster, Fairyduster, False Mesquite, False Mesquite Calliandra, Guajillo, Hairy-leaved Calliandra, Huajillo, Mesquitella (Spanish), Mesquitilla, Mock Mesquite. DESCRIPTION: Terrestrial perennial deciduous subshrub or shrub (4 inches to 5 feet in height, one plant was described as being 40 inches in height with a crown 80 inches in width); the stems are bluish, light gray, whitish or white-gray; the leaves may be leaves grayish, dark green or red; the flowers are cream-white, pink, pink-red, pink-white, purple, red, red and white, reddish-purple, rose or violet-red; flowering generally takes place between early February and mid-June (additional records: two for mid-January, four for mid-August, two for late August, one for early September, one for mid-September, one for early October, three for mid-October, four for late October, two for early November, one for mid-November, two for late November, one for early December, one for mid-December and two for late December). HABITAT: Within the range of this species it has been reported from mountains; rocky peaks; mesas; plateaus; rocky canyons; along canyon bottoms; buttes; knolls; sandy ridges; rocky ridgetops; rocky, shaley-sandy and gravelly-clayey-loamy foothills; rocky hills; hilltops; rocky hillsides; along bedrock, bouldery, rocky, rocky-clayey, gravelly and gravelly-sandy-loamy slopes; gravelly bajadas; rocky outcrops; amongst boulders and rocks; boulderfields; interior dunes; plains; rocky, gravelly and sandy flats; basins; valley floors; along rocky and sandy roadsides; along rocky-sandy arroyos; within gullies; around seeps; around springs; around seeping streams; along and in gravelly and sandy washes; within bouldery drainage ways; along water courses; (rocky) banks of arroyos and lakes; edges of washes and drainage ways; shores of lakes; gravelly terraces; ditches; riparian areas, and disturbed areas growing in dry desert pavement; bouldery, rocky, rocky-sandy, shaley-sandy, gravelly and sandy ground; pebbly-clayey loam, gravelly-sandy loam, gravelly-clayey loam and sandy loam ground, and rocky clay ground, occurring from sea level to 6,900 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, and is a soil binder. This plant was reported to have been utilized by native peoples of North America; it was noted as having been used as a drug or medication. Fairy Duster is browsed by wildlife and found to be highly palatable by Mule Deer (*Odocoileus hemionus*) and White-tailed Deer (*Odocoileus virginianus*), and hummingbirds have been

observed visiting the flowers. *Calliandra eriophylla* is native to southwest-central and southern North America. *5, 6, 13, 15, 16, 18, 28 (color photograph), 43 (080409), 46 (Page 397), 48, 58, 63 (020910 - color presentation), 77 (color photograph #32), 85 (020910 - color presentation), 86 (color photograph), 91, 115 (color presentation), 127, **WTK** (March 9, 2007)*

Calliandra eriophylla var. *erriophylla* (see *Calliandra eriophylla*)

Cercidium floridum (see *Parkinsonia florida*)

Cercidium floridum subsp. *floridum* (see *Parkinsonia florida*)

Cercidium microphyllum (see *Parkinsonia microphylla*)

***Dalea pulchra* H.S. Gentry: Santa Catalina Prairie Clover**

COMMON NAMES: Bush Dalea, Santa Catalina Prairie-clover, Gregg Dalea, Indigo Bush, Indigo Dalea, Pea Bush, Santa Catalina Prairie Clover, Santa Catalina Prairie-clover, Santa Catalina Prairie-clover, Santa Catalina Prairie-clover. DESCRIPTION: Terrestrial perennial subshrub or shrub (reported as being 20 inches to 8 feet in height); the leaves are grayish-green or silvery; the flowers ($\frac{1}{4}$ inch in diameter) are pale lavender with a pink-purple tinge, lavender, lavender-pink, pink-purple, pink-purple & white, pale pinkish-white, light purple, purple, purple & yellow, rose-purple & whitish or violet; based on few records located, flowering generally takes place between late January and late May (flowering records: three for early January, one for late January, two for early February, one for mid-February, early March thru early May, five for late May, one for mid-June, one for early November and one for mid-December). HABITAT: Within the range of this species it has been reported from mountains; mountainsides; rocky mesas; rocky canyons; canyon bottoms; rocky knolls; rocky ridges; ridgetops; gravelly-sandy foothills; rocky hills; hillsides; bedrock, rocky and gravelly slopes; along roadsides; along streams; along and in washes, and drainages growing in dry rocky, gravelly, pebbly and gravelly-sandy ground, occurring from 100 to 4,900 feet in elevation in the woodland, scrub, grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant is a larval food plant for the Southern Dogface (*Colias cesonia*). *Dalea pulchra* is native to southwest-central and southern North America. *5, 6, 13, 15, 18, 28 (color photograph), 43 (082510), 46 (note, Page 439), 58, 63 (082510), 82, **85** (082610 - color presentation), 91, 115 (color presentation)*

***Parkinsonia florida* (G. Bentham ex A. Gray) S. Watson: Blue Paloverde**

SYNONYMY: *Cercidium floridum* G. Bentham, *Cercidium floridum* G. Bentham var. *floridum*. COMMON NAMES: Blue Palo Verde, Blue Palo-verde, Blue Paloverde, Caro (Mayo), Palo Verde (Spanish for Green Pole, Green Stick or Green Tree), Paloverde, Stedak U'us (Pima), Studuk U'us (Bajo Pima). DESCRIPTION: Terrestrial perennial deciduous shrub or tree (40 inches to 40 feet in height); the bark may be blue-green, green, yellow or yellow-green, and gray on the older trunks; the leaves are blue-green; the flowers ($\frac{3}{4}$ to 1 inch in width) are yellow or seldom white; flowering generally takes place between early March and mid-June (additional records: two for early February, two for mid-August, two for early September, one for late September, one for early October, two for mid-October, one for late October, two for early November, one for mid-November and one for early December); the mature fruits ($1\frac{1}{2}$ to 4 inches in length) are light brown. HABITAT: Within the range of this species it has been reported from mountains; mountainsides; mesas; along canyons; canyon walls; sandy canyon bottoms; buttes; gravelly-clayey ridges; foothills; rocky, rocky-sandy, gravelly-loamy and sandy hills; bajadas; rocky, rocky-sandy and sandy slopes; sand hills; sand dunes; rocky-sandy, cindery, sandy and sandy-silty flats; valley floors; valley bottoms; coastal slopes; along rocky-gravelly-sandy, gravelly-sandy and sandy roadsides; along gravelly arroyos; along sandy bottoms of arroyos; rocky draws; seeps; streambeds; creekbeds; along rivers; along riverbeds; along and in rocky, rocky-sandy, gravelly, gravelly-sandy and sandy washes; drainages; watercourses; playas; along (rocky and sandy) banks of arroyos, rivers and

washes; edges of draws and washes; margins of rivers and washes; gravelly sand bars; benches; gravelly terraces; loamy bottomlands; clayey lowlands; sandy-loamy floodplains; mesquite bosques; fencerows; catchments; stock tanks; along canals; along canal banks; gravelly-sandy riparian areas, and disturbed areas growing in dry rocky, rocky-gravelly-sandy, rocky-sandy, cindery, gravelly, gravelly-sandy and sandy ground; gravelly loam, sandy loam and loam ground; gravelly clay and clay ground, and sandy silty ground, occurring from sea level to 5,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, it has a very showy display of yellow flowers in very showy in late March and April. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food crop; it was also noted as having been used for shelter and for tools. The Blue Paloverde may be useful in controlling erosion. Bighorn Sheep (*Ovis canadensis*), Mule Deer (*Odocoileus hemionus*) and other wildlife browse the fruits, leaves and twigs and the seeds are eaten by birds and rodents and used by Bruchid Beetles. *Parkinsonia florida* is native to southwest-central and southern North America. *5, 6, 13 (recorded as *Cercidium floridum* Bentham, color photograph of habitat Plate S.2), 15, 16 (recorded as *Cercidium floridum* Benth.), 18, 26 (recorded as *Cercidium floridum*, color photograph), 28 (recorded as *Cercidium floridum*, color photograph), 43 (021310 - *Cercidium floridum* Benth. ex A. Gray, *Parkinsonia florida* S. Watson), 46 (recorded as *Cercidium floridum* Benth., Page 407), 48, 52 (recorded as *Cercidium floridum* Benth. ex Gray, color photograph), 53 (recorded as *Cercidium floridum* Benth.), 58, 63 (021310 - color presentation), 77 (recorded as *Cercidium floridum* Benth.), 85 (021410 - color presentation), 86 (recorded as *Cercidium floridum*, color photograph), 91 (recorded as *Cercidium floridum* Benth.), 115 (color presentation), 127, **WTK** (March 9, 2007)*

***Parkinsonia microphylla* J. Torrey: Yellow Paloverde**

SYNONYMY: *Cercidium microphyllum* (J. Torrey) J.N. Rose & I.M. Johnston. COMMON NAMES: Dipua, Foothill Palo Verde, Foothill Paloverde, Hillside Paloverde, Horsebean, Kuk Cehedagi (Tohono O'odham), Little Horsebean, Littleleaf Horsebean, Little Leaf Paloverde, Little-leaf Palo Verde, Little-leaf Palo-verde, Little-leaf Paloverde, Littleleaf Palo Verde, Littleleaf Paloverde, Palo Verde (Spanish for Green Pole, Green Stick or Green Tree), Palo-verde, Paloverde, Yellow-Palo-verde, Yellow Paloverde. DESCRIPTION: Terrestrial perennial drought deciduous shrub or tree (40 inches to 26 feet in height with a crown diameter of 12 to 18, one plant was described as being 6 feet in height and 7 feet in width, one plant was described as being 9 feet in height and width, one plant was described as being 16 feet in height and width); the bark is green, olive-green or yellow-green, and gray on older trunks; the ends of the leafy branchlets are spine-like; the small leaflets are green, greenish-gray or yellow-green; the flowers (½ inch in width) are lemon-yellow, whitish & yellow, yellow, yellow-green or yellow & white; the styles are pale yellow or pale yellow-green; the filaments are pale yellow or pale yellow-green; the anthers are orange; flowering generally takes place between mid-March and mid-June (additional records: one for mid-August and one for mid-October); the mature seedpods (2 to 3 inches in length) are light brown or tan. HABITAT: Within the range of this species it has been reported from mountains; rocky mountaintops; mesas; cliffs; rocky walls; rocky canyons; canyon walls; canyon bottoms; bluffs; buttes; ledges; ridges; bouldery and rocky foothills; bases of foothills; rocky hills; rocky hillsides; rocky and sandy slopes; alluvial fans; rocky, gravelly and gravelly-silty bajadas; boulder fields; bouldery and rocky outcrops; plains; gravelly and sandy flats; valley floors; sandy valley bottoms; along rocky and gravelly roadsides; within gravelly-sandy arroyos; along and in rocky, gravelly, gravelly-sandy and sandy washes; drainages; (rocky-sandy) banks of arroyos and rivers; along edges of washes; margins of arroyos and washes; rocky sand bars; coves; gravelly terraces; floodplains; ditches; riparian areas, and disturbed areas growing in dry desert pavement; bouldery, rocky, gravelly, gravelly-sandy and sandy ground; sandy loam, clay loam and loam ground; clay ground, and gravelly silty ground, occurring from sea level to 4,000 feet in elevation in the grassland and desertscrub ecological formations. NOTES: This plant may be an attractive component of a restored native habitat and may live to be more than 400 years of age. This plant was reported to have been utilized by native peoples of North America and could be investigated to

determine its value as a home garden or commercial food crop. To reduce water loss during extended periods of drought a tree may undergo a natural drought-pruning process where entire branches die back. The Foothill Paloverde is a common “nurse plant” of the Saguaro or Giant Cactus (*Carnegiea gigantea*) and provides a sheltered microhabitat in which other desert plants are able to become established. Bighorn Sheep (*Ovis canadensis*), Mule Deer (*Odocoileus hemionus*), jackrabbits and other small mammals browse the fruits, leaves and twigs; the Collard Peccary (*Peccari tajacu*) feed on the fruit, and the seeds are used by Bruchid Beetles. The Foothill Paloverde is considered a significant foraging site for birds; it is used as a nesting site by the Black-tailed Gnatcatcher (*Polioptila melanura*) and Verdins, and as a roosting site by Gambel’s Quail (*Callipepla gambelii* subsp. *gambelii*). The Costa’s Hummingbird (*Calypte costae*) has been observed visiting the flowers. *Parkinsonia microphylla* is native to southwest-central and southern North America. *5, 6, 10, 13 (recorded as *Cercidium microphyllum*, color photograph in habitat Plate T.1), 15, 16 (recorded as *Cercidium microphyllum* (Torr.) Rose & Johnst.), 18, 26 (recorded as *Cercidium microphyllum*, color photograph), 28 (recorded as *Cercidium microphyllum*, color photograph), 43 (021410 - *Cercidium microphyllum* Rose & I.M. Johnst.), 46 (recorded as *Cercidium microphyllum* (Torr.) Rose & Johnston, Page 407), 48, 52 (recorded as *Cercidium microphyllum* (Torr.) Rose & I.M. Johnst., color photograph), 53 (recorded as *Cercidium microphyllum* (Torr.) Rose & Johnst.), 63 (021410 - color presentation), 77 (recorded as *Cercidium microphyllum* (Torr.) Rose & Johnst.), 85 (021410 - color presentation), 86 (note under *Cercidium floridum*), 91 (recorded as *Cercidium microphyllum* (Torr.) Rose & I.M. Johnston), 115 (color presentation), 127, 134, **WTK** (March 9, 2007)*

Prosopis juliflora var. *velutina* (see *Prosopis velutina*)

***Prosopis velutina* E.O. Wooton: Velvet Mesquite**

SYNONYMY: *Prosopis juliflora* (O. Swartz) A.P. de Candolle var. *velutina* (E.O. Wooton) C.S. Sargent. COMMON NAMES: Algarroba, Chachaca, Fluweelprosopis (Afrikaans), Kvi (or possibly Kui - Tohono O’odham), Mesquite, Mezquite, Mizquitl, Velvet Mesquite. DESCRIPTION: Terrestrial perennial deciduous shrub or tree (2 to 56 feet in height, one plant was reported to be 6½ feet in height with a canopy 6½ feet in width, one plant was reported to be 13 feet in height with a canopy 16½ feet in width, one tree was reported to be 20 feet in height and 40 feet in width); the bark on the trunk and older branches is dark brown, dark brownish-green or dark gray; the leaves are gray-green; the flowers (cylindrical spikes 2 to 5 inches in length) are cream, cream-yellow, green-yellow, greenish-white, pale yellow, yellow, yellow-green or yellowish-green; flowering generally takes place between mid-March and late August (additional records: one for early October and one for early November); the mature seedpods (3 to 8 inches in length) are red, tan, yellow or mottled. HABITAT: Within the range of this species it has been reported from mountains; mesas; plateaus; canyons; along sandy canyon bottoms; rocky bases of cliffs; buttes; rocky and sandy ridges; foothills; rocky hills; rocky hillsides; rocky and rocky-loamy slopes; alluvial fans; gravelly bajadas; rocky outcrops; rocky plains; gravelly and sandy flats; sandy valley floors; valley bottoms; along rocky-gravelly-loamy, gravelly-clayey-sandy-loamy and silty-clayey roadsides; along and in sandy arroyos; rocky-gravelly-loamy draws; seeps; springs; around seeping streams; along streams; along rocky streambeds; along creeks; creekbeds; along rivers; along rocky-sandy riverbeds; along and in rocky, gravelly-sandy and sandy washes; along drainages; within drainage ways; playas; cienegas; banks of streams, creeks and rivers; (gravelly and sandy) edges of rivers, washes and ponds; sandy-loamy benches; gravelly and gravelly-sandy terraces; bottomlands; floodplains; mesquite bosques; along fencelines; around stock tanks; around reservoirs; along canals; canal banks; ditches; along ditch banks; riparian areas, and disturbed areas growing in dry rocky, rocky-gravelly, rocky-sandy, gravelly, gravelly-sandy and sandy ground; rocky loam, rocky-gravelly loam, gravelly-clayey-sandy loam, sandy loam, clayey loam and loam ground; silty clay ground, and sandy silty, clayey silty and silty ground, occurring from 100 to 6,300 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, it may live to be more than several hundred years of age. This plant was reported to have been

utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, beverage, fiber and/or dye or paint (boiled resin used as a pottery paint) crop; it was also noted as having been used as fuel, as a tool, as toys, as a drug or medication and as a guide for determining a planting season. The Velvet Mesquite is a common “nurse plant” of the Saguaro or Giant Cactus (*Carnegiea gigantea*). Much of the mesquite forest (bosques) originally found along the desert water courses have been lost to fuel wood cutting and clearing for agricultural fields and commercial and residential development. Velvet Mesquite Bosques were small, open, park-like woodlands with the Velvet Mesquite often occurring in nearly pure stands and interspersed with other common species such as the Netleaf Hackberry (*Celtis laevigata* var. *reticulata*), Catclaw Acacia (*Acacia greggii* var. *greggii*), Mexican Elder (*Sambucus nigra* subsp. *canadensis*), Desert Hackberry (*Celtis ehrenbergiana*), Greythorn (*Ziziphus obtusifolia* var. *canescens*), Wolfberry (*Lycium* spp.), Four-wing Salt-bush (*Atriplex canescens*) and Vine Mesquite Grass (*Panicum obtusum*). The Velvet Mesquite provides food and shelter for many species of wildlife. The plant is a food source for quail, Desert Mule Deer (*Odocoileus hemionus crooki*) and Desert Bighorn Sheep (*Ovis canadensis mexicana*). Coyotes (*Canis latrans*), Round-tailed Ground Squirrels (*Spermophilus tereticaudus*), Desert Cottontails (*Sylvilagus audubonii*) and many other wild animals feed on the seed pods. Velvet Mesquite is the host for a Drywood Termite (*Incisitermes banksi*). Bruchid Beetles feed on the fruits and seeds. *Prosopis velutina* is native to southwest-central and southern North America. *5, 6, 13 (recorded as *Prosopis juliflora* (Swartz) DC. var. *velutina* (Wootton) Sarg., color photograph), 15, 16, 18, 26 (color photograph), 28 (color photograph), 43 (071609), 46 (recorded as *Prosopis juliflora* (Swartz) DC. var. *velutina* (Wootton) Sarg., Page 402), 48, 52 (color photograph), 53 (species: recorded as *Prosopis juliflora* (Sw.) DC.), 58, 63 (021610), 68, 77, 80 (This species is listed as a Rarely Poisonous and Suspected Poisonous Range Plant. “Heavy, long-continued consumption of pods and leaves of these common desert shrubs may cause rumen impaction and poisoning.”), 85 (021610 - color presentation), 91), 115 (color presentation), 127, 134, ADS (Arizona Daily Star, Sunday, July 26, 2009, Tucson & Region, B1: Mesquite Pods are of Consuming Interest), **WTK** (March 9, 2007)*

Vachellia constricta (see *Acacia constricta*)

Fouquieriaceae: The Ocotillo Family

***Fouquieria splendens* G. Engelmann: Ocotillo**

SYNONYMY: *Fouquieria splendens* G. Engelmann subsp. *splendens* G. Engelmann.
COMMON NAMES: Albarda, Barda, Barda, Candle Bush, Candlewood, Coach Whip, Coach-whip, Coachwhip, Coachwhip Cactus, Flamingsword, Jacob’s Staff, Monkey-tail, Ocotillo, Ocotillo del Corral, Slimwood, Vine-cactus, Vine Cactus. DESCRIPTION: Terrestrial perennial cold- and drought-deciduous semi- and stem-succulent shrub (5 to 33 feet in height with a crown width of 5 to 15 feet); the stems (cluster of 5 to 100 wand-like stems branching from the base) are gray, gray & dark gray, gray-green or green; the leaves are green; the flowers (2 to 10 inch long clusters at the tips of the stems) may be coral-red, cream, cream-white, orange, orange-red, pinkish-purple, red, reddish-orange, red & yellow, salmon, scarlet, scarlet-coral, white or yellow; flowering generally takes place over a period of 50 to 60 days between early February and early June (additional records: two for late June, two for early July, one for mid-July, one for late July, one for early August, one for late August, two for mid-September, one for late September, one for mid-October, two for late October, two for early November and two for early December); the mature fruits are capsules containing winged seeds. HABITAT: Within the range of this species it has been reported from mountains; mountainsides; gravelly-sandy and sandy mesas; crags; canyon rims; cliffs; bouldery and rocky canyons; crevices in rocks; gravelly ridges; rocky ridgetops; ridgelines; foothills; rocky and rocky-sandy hills; rocky hilltops; rocky and gravelly hillsides; bedrock, bouldery-cobbly, rocky, rocky-gravelly, shaley-sandy, stony, gravelly, gravelly-sandy and gravelly-loamy slopes; alluvial fans; rocky and sandy bajadas; rocky outcrops; amongst boulders; lava flows; sand hills; sand dunes; dune swales; gravelly outwash fans; gravelly and sandy plains; gravelly and gravelly-sandy

flats; basins; rocky and sandy valley floors; valley bottoms; along gravelly roadsides; rocky arroyos; gullies; along rivers; along sandy washes; (bedrock, bouldery-cobbly and sandy) banks of rivers and washes; (rocky-sandy) shores of lakes; benches; along floodplains and riparian areas growing in dry desert pavement; bouldery, bouldery-cobbly, rocky, rocky-gravelly, rocky-sandy, shaley-sandy, stony, gravelly, gravelly-sandy and sandy ground; gravelly loam ground, and clay ground, occurring from sea level to 7,500 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, beverage and/or fiber crop; it was also noted as having been used as a fuel, tool, drug or medication, ceremonial item and as an ornamental landscape plant. Older plants may be 150 to 200 years of age. This “vase-shaped” plant has been described by Benson and Darrow as being “one of the most distinctive shrubs in the Southwestern Deserts, and it is one of the plants giving outstanding character to the flora of the region”. Desert Bighorn Sheep (*Ovis canadensis* subsp. *mexicana*), Mule Deer (*Odocoileus hemionus*) and Whitetailed Deer (*Odocoileus virginianus* subsp. *couesi*) browse this plant. The Broad-billed Hummingbird (*Cynanthus latirostris*), Butterflies, Carpenter Bee (*Xylocopa californica*), Costa’s Hummingbird (*Calypte costae*), Finches, Orioles, Rufous Hummingbird (*Selasphorus rufus*), Solitary Bees, Syrphid Flies, Verdins, and Warblers have been observed visiting the flowers. The Ocotillo is a preferred food plant of the Costa’s Hummingbird. *Fouquieria splendens* is native to southwest-central and southern North America. *5, 6, 10, 13 (color photograph: Plate N), 15, 16, 18, 26 (color photograph), 28 (color photograph), 43 (080309), 45 (color photograph), 46 (Page 640), 48, 58, 63 (021810 - color presentation), 77 (color photograph #27), 85 (021810 - color presentation), 86 (color photograph), 91, 106 (021810 - color presentation), 107, 115 (color presentation), 127, **WTK** (March 9, 2007)*

Fouquieria splendens subsp. *splendens* (see *Fouquieria splendens*)

Krameriaceae: The Ratany Family

***Krameria* C. Linnaeus: Ratany**

COMMON NAMES: Ratany *43 (062710 - *Krameria* Loefl.), 46 (Pages 403-404), 63 (062710 - color presentation), 85 (062710 - color presentation), **WTK** (March 9, 2007)*

Malpighiaceae: The Barbados-cherry Family

***Janusia gracilis* A. Gray: Slender Janusia**

COMMON NAMES: Desert Vine, Fermina, Slender Janusia. DESCRIPTION: Terrestrial perennial deciduous forb/herb or vine (clambering, climbing, scrambling or twining stems 16 inches to 10 feet in length, one plant was reported to have been 16 inches in height with a crown 10 inches in diameter); the leaves are grayish-green or reddish; the flowers (to ½ inch in width) are orange-yellow or yellow; flowering generally takes place between early March and mid-November (additional records: two for early January, one for late January, one for early December, one for mid-December and one for late December); the winged fruits (paired samaras) are pink, purple-red, red, red-green or reddish. HABITAT: Within the range of this species it has been reported from mountains; rocky mountaintops; rocky mountainsides; mesas; cliffs; rocky canyons; sandy canyon bottoms; gravelly-sandy bases of cliffs; amongst crevices; rocky buttes; rocky knolls; rocky and gravelly ridges; rocky ridgetops; foothills; rocky hills; rocky hillsides; along bouldery-rocky, rocky, rocky-gravelly, rocky-clayey-loamy and gravelly slopes; alluvial fans; gravelly bajadas; volcanic plugs; bouldery and rocky outcrops; amongst rocks; plains; gravelly flats; basins; valley floors; rocky-gravelly roadsides; along rocky arroyos; bottoms of arroyos; draws; within gullies; ravines; along streams; along rocky streambeds; along creeks; bouldery-

rocky-sandy creekbeds; along and in gravelly and sandy washes; along drainages; waterholes; palm oases; (rocky) banks of streams; edges of washes; benches; floodplains, and riparian areas growing in dry bouldery, bouldery-rocky, bouldery-rocky-sandy, rocky, rocky-gravelly, gravelly, gravelly-sandy and sandy ground and rocky-clayey loam and clayey loam ground, occurring from sea level to 7,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. It is small woody vine often reported as scrambling over rocks, twining among shrubs or forming small tangled shrublets. Slender Janusia is browsed by the Sonoran Desert Tortoise (*Gopherus agassizi*), Desert Mule Deer (*Odocoileus hemionus* subsp. *crooki*) and Whitetail Deer (*Odocoileus virginianus* subsp. *couesi*). *Janusia gracilis* is native to southwest-central and southern North America. *5, 6, 13, 15, 16, 28 (color photograph), 43 (030310), 46 (Page 497), 48, 58, 63 (030310 - color presentation), 77 (color photograph #83), 85 (030310 - color presentation), 115 (color presentation), **WTK** (March 9, 2007)*

Rhamnaceae: The Buckthorn Family

Condalia lycioides var. *canescens* (see *Ziziphus obtusifolia* var. *canescens*)

Condalia spathulata (see footnote 46 under *Condalia warnockii* var. *kearneyana*)

***Condalia warnockii* M.C. Johnston var. *kearneyana* M.C. Johnston: Kearney's Snakewood**

COMMON NAMES: Crucillo, Guichutilla, Kearney Condalia, Kearney Snakewood, Kearney's Snakewood, Mexican Buckthorn, Mexican Crucillo, Squaw-bush, Squawbush. DESCRIPTION: Terrestrial perennial deciduous (considered evergreen except during periods of severe drought) shrub (20 inches to 13 feet in height, one plant was reported to be 6½ feet in height with a crown 10 feet in width, one plant was reported to be 10 feet in height with a crown 10 feet width); the minute flowers are yellowish; flowering generally takes place between February and November (flowering records: one for mid-February, one for mid-August, one for late August and one for mid-September; however, flowering taking place throughout the year has also been reported); the fruits are black, dark purple or reddish black. HABITAT: Within the range of this species it has been reported from mountains; gravelly and sandy mesas; cliff faces; canyons; canyon bottoms; rocky ledges; ridges; edges of meadows; foothills; hills; rocky, gravelly and sandy slopes; rocky and gravelly bajadas; amongst boulders; gravelly and sandy flats; basins; valley floors; rocky arroyos; gulches; rocky washes; along and in drainages; banks of creeks; (gravelly) edges of washes and drainages; terraces; floodplains, and around gravelly-sandy stock tanks growing in dry bouldery, rocky, gravelly, gravelly-sandy and sandy ground, occurring from 1,600 to 5,600 feet in elevation in the grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. *Condalia warnockii* var. *kearneyana* is native to southwest-central and southern North America. *5, 6, 13, 15, 16, 28 (color photograph), 43 (042210), 46 (recorded as *Condalia spathulata* A. Gray, Page 530), 58, 63 (042210), 77, 85 (042210 - color presentation), 91, **WTK** (March 9, 2007)*

***Ziziphus obtusifolia* (W.J. Hooker ex J. Torrey & A. Gray) A. Gray var. *canescens* (A. Gray) M.C. Johnston: Lotebush**

SYNONYMY: *Condalia lycioides* (A. Gray) A. Weberbauer var. *canescens* (A. Gray) W. Trelease. COMMON NAMES: Abrojo, Bachata, Barbachatas, Buchthorn, Clepe, Garrapata, Garumbullo, Gray-leafed Abrojo, Gray-leaved Abrojo, Gray-thorn, Greythorn, Gumdrop Tree, Lotebush, Oschuvapat (Pima), Palo Blanco, Southwestern Condalia, White Crucillo. DESCRIPTION: Terrestrial perennial drought deciduous shrub or tree (3 to 13 feet in height, one plant was reported to be 40 inches in height with a crown 18 inches in width, one was reported to be plant 7 feet in height with a crown 7 feet in width, one plant was reported to be 10 feet in height with a crown 10 feet in width, one plant was reported to be 13 feet in height with a crown 13 feet in width); the stems are bluish, gray, gray-green, green or

whitish with the twigs ending in stout thorns; the leaves are gray-green, green or yellow-green, the inconspicuous flowers are cream, green, greenish-white, yellow, yellow-green or whitish-green; flowering generally takes place between mid-May and late November (additional records: one for late January, one for mid-March, one for late March, one for mid-April and one for late April); the ripe fruits are black, blue-purple, dark blue or purple. HABITAT: Within range of this species it has been reported from mountains; mesas; rocky canyons; along canyon bottoms; scree; talus slopes; bases of cliffs; crevices in rocks; buttes; ridges; ridgelines; foothills; rocky hills; hilltops; rocky hillsides; rocky and gravelly slopes; rocky alluvial fans; gravelly bajadas; amongst boulders, rocks and gravels; sandy-silty plains; rocky and gravelly flats; basins; rocky valley floors; gravelly and gravelly-loamy roadsides; arroyos; bottoms of arroyos; gulches; ravines; bouldery bottoms of ravines; seeps; in clay around springs; rivulets; along streams; along rocky streambeds; along creeks; along gravelly-sandy creekbeds; along gravelly and gravelly-sandy rivers; riverbeds; along and in rocky and sandy washes; drainages; marshes; along (rocky) banks of streams, creeks, rivers and washes; (gravelly-sandy) edges of arroyos and creeks; beaches; sandy benches; terraces; bottomlands; floodplains; mesquite bosques; along fencerows; along canals; gravelly-sandy riparian areas, and disturbed areas growing in dry desert pavement; bouldery, bouldery-sandy, rocky, rocky-gravelly, rocky-gravelly-sandy, gravelly, gravelly-sandy and sandy ground; cobbly-gravelly loam, gravelly loam and gravelly-clayey loam ground; sandy clay and clay ground, and sandy silty ground, occurring from sea level to 5,500 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, fodder and/or beverage (*Ziziphus obtusifolia*) crop; it was also noted as having been used as a tool, as a drug or medication and as a commodity used in personal hygiene. The flowers are visited by orange-winged Spider Wasps. Gray Foxes (*Urocyon cinereoargenteus*), Raccoons (*Procyon lotor*), Ringtails (*Bassariscus astutus*), Gambel's Quail (*Callipepla gambelii*), Scaled Quail (*Callipepla squamata*), Mockingbirds (*Mimus polyglottos*), Northern Orioles (*Icterus bullockii*), Phainopeplas (*Phainopepla nitens*), Band-tailed Pigeons (*Columba fasciata*), White-necked Ravens (*Corvus cryptoleucus*), Curved-billed Thrashers (*Toxostoma curvirostre*), Golden-fronted Woodpeckers (*Melanerpes aurifrons*), White-winged Doves (*Zenaida asiatica*) and other birds feed on the fruit. The plants numerous spines provide an impenetrable refuge for birds and many species of birds make use of the Lotebush as a preferred nesting site. *Ziziphus obtusifolia* var. *canescens* is native to southwest-central and southern North America. *5, 6, 13 (color photograph), 15, 16, 28 (species, color photograph of species), 43 (042210), 46 (recorded as *Condalia lycioides* (Gray) Weberb. var. *canescens* (Gray) Trel., Page 530), 58, 63 (042210), 77, 85 (042310 - color presentation of dried material), 91, 127, **WTK** (March 9, 2007)*

Solanaceae: The Potato Family

***Lycium* C. Linnaeus: Desert-thorn**

COMMON NAMES: Desert-thorn, Lycium, Thornbush, Wolfberry. *43 (052010), 46 (Pages 749-752), 63 (040207), **WTK** (March 9, 2007)*

***Nicotiana obtusifolia* M. Martens & H.G. Galeotti var. *obtusifolia*: Desert Tobacco**

SYNONYMY: *Nicotiana trigonophylla* M.F. Dunal. COMMON NAMES: Coyote Tobacco, Desert Tobacco, Punche (a Punch), Tabaquillo (Little Tobacco), Tabaquillo de Coyote, Wo'i Viva (Yaqui). DESCRIPTION: Terrestrial annual, biennial or perennial forb/herb or subshrub (1 to 3½ feet in height); the leaves are gray-green or dark green; the flowers may be cream, cream & pale green, cream-green, cream-white, cream-yellow, greenish, greenish-white, greenish-yellow, deep purple, lemon-yellow, pale white, white, yellow, yellow-cream, yellow-green, yellow-white or yellowish-greenish; flowering generally takes place between late February and early November (additional records: one for mid-January, one for late November, one for mid-December and one for late December, flowering probably takes place

throughout the rest of the year). HABITAT: Within the range of this species it has been reported from mountains; bouldery and rocky-gravelly mountaintops; plateaus; along rims; cliffs; rocky and gravelly-loamy canyons; canyon walls; along canyon bottoms; gorges; talus slopes; bases of cliffs; along crevices in boulders and rocks; rocky bluffs; rocky buttes; rocky ledges; bouldery ridges; bouldery ridgetops; edges of meadows; craters; cinder cones; foothills; rocky hills; hilltops; bouldery-rocky and rocky hillsides; bouldery escarpments; bouldery, bouldery-rocky, bouldery-gravelly, rocky, rocky-gravelly-sandy-clayey, cindery, gravelly-sandy, sandy-loam and sandy-clayey slopes; bajadas; rocky outcrops; amongst boulders, rocks and stones; bases of boulders and rocks; sandy lava flows; dunes; debris fans; rocky plains; sandy flats; basins; valley floors; valley bottoms; rocky-sandy coastal shores; along railroad right-of-ways; along rocky, rocky-gravelly-sandy-clayey-loamy, gravelly, gravelly-sandy, gravelly-sandy-clayey-loamy and sandy roadsides; along sandy-loamy arroyos; arroyo walls; arroyo bottoms; in sand and loam around springs; along streams; along gravelly-sandy and sandy streambeds; rocky creeks; sandy creekbeds; bouldery-sandy and sandy riverbeds; along and in bedrock, rocky, rocky-sandy, gravelly-sandy and sandy washes; drainages; bouldery drainage ways; sandy waterholes; marshy areas; (rocky, cobbly, sandy and silty) banks of creeks, rivers and washes; edges of lakes; rocky-sandy shores of lakes; mudflats; gravelly and sandy terraces; bottomlands; floodplains; ditches; ditch banks; sandy riparian areas; waste places, and disturbed areas growing in moist, damp and dry bouldery, bouldery-rocky, bouldery-gravelly, bouldery-sandy, rocky, rocky-gravelly, rocky-sandy, cobbly, cindery, gravelly, gravelly-sandy and sandy ground; rocky-gravelly-sandy-clayey loam, gravelly loam, gravelly-sandy-clayey loam, sandy loam, clayey loam and loam ground; rocky-gravelly-sandy clay and sandy clay ground, and silty ground, occurring from sea level to 6,900 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant tends to be weedy; however, consideration could be given to using some plants in your project because the flowers are used by hummingbirds when other nectar-rich sources are not available. This plant was reported to have been utilized by native peoples of North America; it was noted as having been used for food, beverage and/as a as a drug or medication. *Nicotiana obtusifolia* var. *obtusifolia* is native to southwest-central and southern North America. *5, 6, 15 (recorded as *Nicotiana trigonophylla* Dunal), 16 (recorded as *Nicotiana trigonophylla* Dunal), 28 (recorded as *Nicotiana trigonophylla*, color photograph), 43 (050310), 46 (recorded as *Nicotiana trigonophylla* Dunal, Page 761), 58 (recorded as *Nicotiana trigonophylla* Dunal), 63 (050310 - color presentation), 68, 77 (recorded as *Nicotiana trigonophylla* Dunal), 80 (This species is listed as a Secondary Poisonous Range Plant. “The poisonous principle is the highly toxic nicotine and other alkaloids which are poisonous to all classes of livestock and to humans. The plants are generally unpalatable to range livestock but frequent losses have been reported. ... Since wild tobaccos are generally unpalatable and grow predominantly in waste places, range improvement to reduce waste areas and to provide ample forage is the best means of preventing losses.”), 85 (050310 - color presentation), 86 (recorded as *Nicotiana trigonophylla*, color photograph), 115 (color presentation of the species), 127, **WTK** (March 9, 2007)*

Nicotiana trigonophylla (see *Nicotiana obtusifolia* var. *obtusifolia*)

Ulmaceae: The Elm Family

***Celtis ehrenbergiana* (J.F. Klotzsch) F.M. Liebmann: Spiny Hackberry**

SYNONYMY: *Celtis pallida* J. Torrey, *Celtis tala* J. Gillies ex J. É. Planchon var. *pallida* (J. Torrey) J. É. Planchon. COMMON NAMES: Acebuche, Bainoro, Capul, Desert Hackberry, Garabato, Garambullo, Granjeno (Spanish), Huasteco, Kunwo (Yaqui), Palo de Aguila, Rompecapa, Shiny Hackberry, Spiny Hackberry. DESCRIPTION: Terrestrial perennial evergreen shrub or tree (3 to 20 feet in height, one plant was reported to be 7 feet in height with a crown 7 feet in width); the bark is gray; the thorny branches are whitish-gray; the leaves are dark green; the inconspicuous flowers may be green, greenish-yellow, white-green or yellow, flowering generally takes place between early March and late

October (possibly flowering into November); the ripe fruits are orange, bright red, reddish-orange or yellow. HABITAT: Within the range of this species it has been reported from mountains; mesas; rocky and rocky-gravelly canyons; canyon bottoms; rocky bases of cliffs; ridges; rocky ridgetops; foothills; rocky hills; rocky hillsides; bedrock, bouldery, rocky and gravelly slopes; bajadas; rocky outcrops; amongst boulders; coves; plains; gravelly-sandy and sandy flats; rocky-gravelly basins; along roadsides; rocky arroyos; rocky bottoms of arroyos; draws; gullies; seeps; springs; along seeping streams; along streams; along and in streambeds; in sand along creeks; along rivers; bouldery-cobbly-sandy riverbeds; along and in gravelly and sandy washes; within drainages; banks of arroyos, rivers, washes and drainages; along margins of arroyos and washes; benches; gravelly terraces; gravelly-clayey floodplains; mesquite bosques; around stock tanks; riparian areas, and disturbed areas growing in dry desert pavement; bouldery, bouldery-rocky, bouldery-cobbly-sandy, rocky, rocky-gravelly, gravelly, gravelly-sandy and sandy ground, and gravelly clay ground, occurring from sea level to 5,600 feet in elevation in the scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. The small fruits are reportedly juicy and sweet. The Desert Hackberry may live to be more than 88 years of age and may be useful in controlling erosion. The Desert Hackberry is a larval food plant for the American Snout (*Libytheana carinenta*) and Empress Leilia (*Asterocampa leilia*) and is browsed by deer; it provides a nesting site for the White-wing Dove (*Zenaida asiatica*), and cover for Gambel's Quail (*Callipepla gambelii gambelii*) and other birds and mammals. The fruits are eaten by many birds, small desert mammals, coyotes (*Canis latrans*), foxes and javelinas (*Peccari tajacu*). *Celtis ehrenbergiana* is native to south-central and southern North America; Central America and coastal islands in the Caribbean Sea, and western, eastern and southern South America. *5, 6, 13 (recorded as *Celtis tala* Gillies var. *pallida* (Torrey) Planch.), 15 (recorded as *Celtis pallida* Torr.), 16 (recorded as *Celtis pallida* Torr.), 18, 26 (recorded as *Celtis pallida*, color photograph), 28 (recorded as *Celtis pallida*, color photograph), 43 (050810), 46 (recorded as *Celtis pallida* Torr., Page 220), 48, 58 (recorded as *Celtis pallida* Torr.), 63 (050810), 77 (recorded as *Celtis pallida* Torr.), 85 (050810, also recorded as *Celtis pallida* var. *pallida* Torrey), 91 (recorded as *Celtis pallida* Torr.), 115 (color presentation), **WTK** (March 9, 2007)*

***Celtis laevigata* C.L. von Willdenow var. *reticulata* (J. Torrey) L.D. Benson: Netleaf Hackberry**

SYNONYMY: *Celtis reticulata* J. Torrey. COMMON NAMES: Canyon Hackberry, False Elm, Hack Berry, Hackberry, Netleaf Hackberry, Palo Blanco, Sugar-berry, Sugarberry, Western Hackberry. DESCRIPTION: Terrestrial perennial deciduous shrub or tree (6 to 60 feet in height, one plant was described as being 30 feet in height and width, stunted trees or shrubs of up to 2 feet in height were reported from forests at higher elevations); the bark is gray, dark gray or reddish-brown becoming "warty" with age; the twigs are reddish-brown; the leaves are gray-green, dark green or yellow-green appearing in early April to late May developing fully in June, they turn yellow in the fall; the inconspicuous flowers are green or yellow-green; the anthers are green; the stigmas are whitish-green; flowering generally takes place between mid-March and mid-September; the fruits are black, purplish, pale orange, orange, orange-red-brown, reddish or reddish-black. HABITAT: Within the range of this species it has been reported from mountains; rocky edges of mesas; plateaus; rocky cliffs; along bouldery, rocky, rocky-gravelly and gravelly-loamy canyons; bouldery, rocky, gravelly and gravelly-sandy-clayey canyon bottoms; chasms; gorges; bases of cliffs; bouldery talus; crevices in rocks; bluffs; ledges; rocky ridges; rocky ridgetops; foothills; sandy hills; rocky hillsides; bouldery, bouldery-sandy, rocky and loamy slopes; alluvial fans; rocky outcrops; amongst boulders and rocks; rocky and sandy alcoves; sandy lava flows; lava beds; sand dunes; shell banks; breaks; prairies; plains; sandy flats; basins; sandy valley floors; valley bottoms; along gravelly-loamy roadsides; along arroyos; sandy bottoms of arroyos; bottoms of draws; gulches; rocky gullies; ravines; sandy seeps; springs; along streams; along and in gravelly-sandy and sandy streambeds; in sand along creeks; along and in bouldery and sandy creekbeds; along rivers; riverbeds; along and in rocky, rocky-gravelly, gravelly, sandy and sandy-clayey-loamy washes; drainages; loamy drainage ways; watercourses; oases; among and in pools; ponds; lakes; tanks; rocky banks of arroyos, ravines, streams, streambeds, creeks, rivers, washes and drainages; sandy edges of arroyos,

springs, streams and washes; margins of ponds; shores of lakes; rocky-sandy and gravelly-sandy benches; gravelly, sandy and silty-loamy terraces; silty bottomlands; along floodplains; mesquite bosques; fencerows; gravelly canal banks; along ditches; rocky-gravelly and sandy riparian areas, and disturbed areas growing in wet, moist, damp and dry bouldery, rocky, rocky-gravelly, rocky-sandy, gravelly, gravelly-sandy and sandy ground; gravelly loam, sandy-clayey loam, silty loam and loam ground; gravelly-sandy clay ground, and silty ground, occurring from 400 to 7,700 feet in elevation in the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial food, fiber and/or dye crop; it was also noted as having been used for tools, as a drug or medication or as a fuel. The Netleaf Hackberry may be useful in the rehabilitation of disturbed sites and suitable for planting in patios, yards and along streets in urban areas and may live to be 100 to 200 years in age. The Netleaf Hackberry provides cover and food for many species of birds and mammals; the American Beaver (*Castor canadensis*) feeds on the wood; the plant is browsed by Pronghorn (*Antilocapra americana*), Mule Deer (*Odocoileus hemionus*) and White-tailed Deer (*Odocoileus virginianus*); the fruit is eaten by wildlife; and Scrub Jays (*Aphelocoma californica*) feed on the leaf galls that form on the foliage. *Celtis laevigata* var. *reticulata* is native to south-central and southern North America. *5, 6, 13, 15 (recorded as *Celtis reticulata* Torr.), 18, 26 (recorded as *Celtis reticulata*, color photograph), 28 (recorded as *Celtis reticulata*, color photograph), 43 (050910 - *Celtis laevigata* var. *reticulata* (Torr.) L.D. Benson), 46 (Page 220), 48, 52 (recorded as *Celtis reticulata*, color photograph), 53, 58 (recorded as *Celtis reticulata* Torr.), 63 (050910 - color presentation), 85 (050910 - color presentation), 115 (color presentation), 127, **WTK** (March 9, 2007)*

Celtis pallida (see *Celtis ehrenbergiana*)

Celtis pallida var. *pallida* (see footnote 85 under *Celtis ehrenbergiana*)

Celtis reticulata (see *Celtis laevigata* var. *reticulata*)

Celtis tala var. *pallida* (see *Celtis ehrenbergiana*)

Verbenaceae: The Verbena Family

***Aloysia wrightii* (A. Gray) A.A. Heller: Wright's Beebrush**

SYNONYMY: *Lippia wrightii* A. Gray. COMMON NAMES: Altamisa, Bee Brush, Beebrush, Lemon Verbena, Mexican Oregano (a common name which is also applied to *Aloysia lycioides* which is the Mexican Oregano of commerce), Mintbush Lippia, Oreganillo, Vara Dulce, Wild Lemon Verbena, Wright Aloysia, Wright Beebrush, Wright's Beebrush, Wright Aloysia, Wright Lippia. DESCRIPTION: Terrestrial perennial drought-deciduous or semi-evergreen shrub (20 inches to 6½ feet in height and about the same in width); the small flowers, located in dense elongate spikes (¾ to 2¾ inches in length and ½ inch in width) are cream-white or white; flowering generally takes place between early March and early May and between mid-July and mid-December (additional record: one for early January). HABITAT: Within the range of this species it has been reported from mountains; rocky and clayey mesas; cliffs; rims of gorges; bedrock, bouldery-sandy, rocky, gravelly and gravelly-loamy canyons; along rocky canyon bottoms; gorges; rocky and gravelly bases of cliffs; talus slopes; crevices in rocks; buttes; rocky ledges; ridges; rocky ridgetops; clearings in woodlands; rocky foothills; rocky hills; rocky and rocky-clayey hillsides; bedrock, bouldery, rocky, rocky-gravelly-loamy, rocky-sandy-clayey-loamy, stony, gravelly and gravelly-sandy-loamy slopes; bajadas; rocky outcrops; amongst rocks; sandy lava flows; lava beds; debris fans; breaks; plains; rocky flats; basins; rocky valley floors; along roadsides; along rocky arroyos; bottoms of arroyos; within rocky draws; within ravines; along streams; creekbeds; along rivers; along and

in bouldery, rocky, cobbly, gravelly and sandy washes; within drainages; marshy areas; (rocky) banks of rivers and washes; edges of arroyos and washes; (rocky) margins of arroyos; (rocky-sandy) shores of lakes; gravel bars; terraces; floodplains; along ditches, and riparian areas growing in damp and dry rocky desert pavement; bouldery, bouldery-rocky, bouldery-sandy, rocky, rocky-gravelly, rocky-sandy, cobbly, cindery-sandy, gravelly and sandy ground; rocky-gravelly loam, rocky-sandy-clayey loam, gravelly loam and gravelly-sandy loam ground, and rocky clay and clay ground, occurring from 1,000 to 7,500 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat, and has been suggested for use as an informal hedge; in herb gardens, and natural landscapes. This plant was reported to have been utilized by native peoples of North America and could be investigated to determine its value as a home garden or commercial beverage crop; it was also noted as having been used as a drug or medication. The Wright Beebrush may live to be more than 72 years of age. is native to southwest-central and southern North America. *5, 6, 13 (*Lippia wrightii* A. Gray), 15, 16, 18, 43 (050910 - *Aloysia wrightii* A. Heller), 46 (Page 729), 58, 63 (050910 - color presentation), 77, 85 (050910 - color presentation), 91, 127, **WTK** (March 9, 2007)*

Lippia wrightii (see *Aloysia wrightii*)

Viscaceae (Loranthaceae): The Christmas Mistletoe Family

***Phoradendron californicum* T. Nuttall: Mesquite Mistletoe**

SYNONYMY: *Phoradendron californicum* T. Nuttall var. *distans* W. Trelease. COMMON NAMES: American Mistletoe, Desert Mistletoe, Mesquite American Mistletoe, Mesquite Mistletoe, Toji, Western Dwarf Mistletoe. DESCRIPTION: Terrestrial perennial subshrub or shrub (8 inches to 5 feet in height, one clump was described as being 16 inches in length and 36 inches in width); the stems (16 to 40 inches in length) may be brown, green, green-reddish, dark olive-green, reddish, red-brown, yellow-green or yellowish; the fragrant flowers are greenish-yellow; flowering generally takes place between late July and early June (additional records: one record for late June and one record for early July); the fruits may be orange, orange-pink, pink, pink-red, pale red, reddish, red-orange, white, white-pink or white-reddish with the older berries turning brown-red or red. HABITAT: Partial parasite observed growing on Catclaw Acacia, Kearney Condalia, Blue Paloverde and Velvet Mesquite, and commonly reported as growing on: *Acacia* spp. (*Acacia constricta*, Whitethorn Acacia; *Acacia farnesiana*, Sweet Acacia, and *Acacia greggii*, Catclaw Acacia); *Condalia* spp. (*Condalia globosa*, Bitter Snakewood and *Condalia warnockii*, Kearney Snakewood); *Larrea tridentata*, Creosote Bush; *Olneya tesota*, Desert Ironwood; *Parkinsonia* spp. (*Parkinsonia aculeata*, Jerusalem Thorn; *Parkinsonia florida*, Blue Palo Verde; *Parkinsonia microphylla*, Yellow Palo Verde, and *Parkinsonia praecox*, Sonoran Palo Verde); *Prosopis* spp. (*Prosopis glandulosa*, Honey Mesquite; *Prosopis pubescens*, Screwbean Mesquite, and *Prosopis velutina*, Velvet Mesquite); *Simmondsia chinensis*, Jojoba, and *Ziziphus obtusifolia*, Lotebush, occurring from sea level to 5,100 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: When removing the Mesquite Mistletoe from the trees and shrubs on your property consider leaving some of the plants for wildlife, Verdins nest in the stems and the Phainopepla (*Phainopepla nitens*) feeds on the berries. This plant was reported to have been utilized by native peoples of North America; it was noted as having been used as a food (berries) and as a drug or medication. *Phoradendron californicum* is native to southwest-central and southern North America. *5, 6, 13 (color photograph), 15, 16, 28 (color photograph), 43 (051710 - *Phoradendron californicum* var. *distans* Trel. in Trel.), 46 (recorded as *Phoradendron californicum* Nutt., Page 224 and *Phoradendron californicum* Nutt. var. *distans* Trelease, Page 224), 58, 63 (051410 - color presentation), 77, **80** (Species of the genus *Phoradendron* are considered to be Rarely Poisonous and Suspected Poisonous Range Plants. "Cattle may be killed by browsing these parasitic forbs, but plants are unpalatable and poisoning is rare. Also children

may be poisoned by eating the berries.”), 85 (051410 - color presentation), 97, 115 (color presentation), 127, **WTK** (March 9, 2007)*

Phoradendron californicum var. *distans* (see *Phoradendron californicum*)

Zygophyllaceae: The Creosote-bush Family

Larrea divaricata (see *Larrea tridentata* var. *tridentata*)

Larrea divaricata subsp. *tridentata* (see *Larrea tridentata* var. *tridentata*)

***Larrea tridentata* (A.P. de Candolle) F.V. Coville var. *tridentata*: Creosote Bush**

SYNONYMY: *Larrea divaricata* auct. non A.J. Cavanilles, *Larrea divaricata* A.J. Cavanilles subsp. *tridentata* (A.P. de Candolle) R.S. Felger & C.H. Lowe. COMMON NAMES: Chaparral, Coville Creosotebush, Creosote Bush, Creosote-bush, Creosotebush, Gobernadora, Greasewood (erroneously called), Guamis, Hediondilla (Spanish - for Little Bad Smeller). DESCRIPTION: Terrestrial perennial evergreen shrub (20 inches to 13 feet in height and about the same in width); the bark is gray; the leaves are bright glossy green or yellow-green; the flowers ($\frac{1}{2}$ to 1 inch in diameter) are yellow or yellow-white; flowering takes place throughout the year with the peak blooming periods occurring in the spring, between March and April, and then again between November and December; the round, fuzzy fruits ($\frac{1}{4}$ inch in diameter) are gray, reddish, white or rust colored. HABITAT: Within the range of this species it has been reported from mountains; rocky, gravelly and sandy mesas; plateaus; rims of canyons; sandy canyons; canyon bottoms; talus slopes; sandy pockets of soil; rocky ridges; foothills; hills; hillsides; rocky and gravelly slopes; alluvial fans; gravelly and sandy bajadas; rocky outcrops; amongst boulders and rocks; sand dunes; sandy plains; cindery-gravelly, gravelly and sandy flats; valley floors; sandy roadsides; arroyos; bottoms of arroyos; riverbeds; along and in gravelly-sandy and sandy washes; (sandy) banks of streams, creeks and rivers; edges of washes; gravelly and sandy terraces; floodplains; mesquite bosques; riparian areas, and disturbed areas growing in dry bouldery, rocky, rocky-sandy, stony, gravelly, gravelly-sandy and sandy ground; rocky-clayey loam and clayey loam ground; sandy clay ground, and rocky-sandy silty and silty ground, occurring from below sea level to 5,000 feet in elevation in the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: This plant may be an attractive component of a restored native habitat. This plant was reported to have been utilized by native peoples of North America; it was noted as having been used as a building material, as tools, in the making of brooms, brushes and musical instruments, as a drug or medication and in body art. Older stems of the Creosote Bush may be 40 to 90 years of age. Using Creosote Bush in the restoration of disturbed sites may increase water infiltration and storage, transplants recommended over spot-seeding and rodent protection for the transplanted seedlings is necessary. When planting a Creosote Bush consider planting a small Desert Night-blooming Cereus (*Peniocereus greggii* var. *transmontanus*) at the base of the plant. The branches will provide support and the roots will protect the tuber of the cereus from hungry Javelinas. The Creosote Bush is the characteristic plant of the southwestern deserts in North America with its distribution very closely delineating the desert regions. As the Creosote Bush ages the older central stems of the plant die off and new stems form at the outer edge of the crown. New stems are not created at the center of the plant. As the crown of the plant expands a “clonal ring”, made up of genetically identical individual shrublets, develops which continues the outward expansion of the ring eventually reaching several yards in diameter. It has been estimated that some of the older rings approach from 9,400 to 11,700 years of age. The Creosote Bush provides cover for many animals; Lac Scale insects (*Tachardiella larreae*), jackrabbits, desert woodrats and other small mammals feed on this plant; stem galls are produced in response to the Creosote Gall midge (*Asphondylia* sp.), and the Desert Tortoise (*Gopherus agassizi*) often digs its shelter under the base of the plant where the roots help to stabilize the soil. *Larrea tridentata* var. *tridentata* is native to southwest-central and southern North America. *5, 6, 13 (color

photograph), 16, 18, 26 (species, recorded as *Larrea tridentata*, color photograph of species), 28 (species, recorded as *Larrea tridentata*, color photograph of species), 43 (051710 - *Larrea tridentata* Coville, *Larrea divaricata* Cav. subsp. *tridentata* (Sessé & Moc. ex DC.) Felger), 46 (species, recorded as *Larrea tridentata* (DC.) Coville: “An outstanding xerophyte and a very important element of the perennial desert flora in southern and western Arizona. ... Creosote-bush has a strong characteristic odor, especially noticeable when the foliage is wet. The plant is ordinarily not touched by livestock, although it is reported that sheep, especially pregnant ewes, have been killed by partaking of it. This plant is reported to cause dermatitis in exceptional persons who are allergic to it.”, Page 491), 48, 63 (051610 - color presentation), 77 (color photograph #101), 80 (This species is listed under Rarely Poisonous and Suspected Poisonous Range Plants. “Early reports accusing this common desert shrub of being poisonous have been proven wrong.”), 85 (051610 - color presentation), 91, 101 (species, color photograph of species), 107, 115 (color presentation), 127, **WTK** (March 9, 2007)*

LISTING OF ANIMALS

STRICTLY ENFORCED LAWS PROTECT MANY OF ARIZONA'S NATIVE
ANIMALS FROM COLLECTION AND FROM BEING DISTURBED OR KILLED

Operation GAME THIEF: 602-942-3000

Kingdom Animalia: The Animal Kingdom

Subkingdom Metazoa: The Multicellular Animals

Section Deuterostomia: The Deuterostomes

Phylum Chordata: The Chordates

Subphylum Vertebrata: The Vertebrates

CLASS MAMMALIA: The MAMMALS

Antilocapridae: The Pronghorn Family

***Antilocapra americana* G. Ord: Pronghorn**

COMMON NAMES: American Pronghorn, “Antelope”, Chihuahuan Pronghorn, Chihuahuan Pronghorn Antelope, Prong-horn, Pronghorn, Pronghorn Antelope, Prong-horned Antelope, Sonoran Pronghorn, Sonoran Pronghorn Antelope. HABITS: Feeds on cacti, forbs, grasses and shrubs. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (113006 - subsp. *americana* (Ord), *mexicana* Merriam and *sonoriensis* Goldman), 55 (recorded as *Antilocapra americana* Ord. Prong-horned Antelope. Formerly widely distributed in grassland areas throughout the state; presently restricted to areas of favorable habitat.), 65, 73, 106 (052806), 100 (color photograph), 110 (Historic Range: Southwest Arizona, south of the Bill Williams River and east to the Santa Cruz River. In Mexico, the northern part of the State of Sonora.), 118 (recorded as *Antilocapra americana americana* (Ord) - Distribution: mapping

and records for northeastern and northwestern Arizona; *Antilocapra americana mexicana* Merriam - Distribution: Southeastern Arizona, and *Antilocapra americana sonoriensis* Goldman - Distribution: Southwestern Arizona. Figure 111, Page 255)*

***Antilocapra americana* subsp. *mexicana* C.H. Merriam: Chihuahuan Pronghorn**

COMMON NAMES: “Antelope”, Chihuahuan Pronghorn, Chihuahuan Pronghorn Antelope, Prong-horn, Pronghorn, Pronghorn Antelope, Prong-horned Antelope. HABITS: The species feeds on cacti, forbs, grasses and shrubs. HABITAT: Within the range of this species it has been reported from the woodland, grassland, desertscrub and wetland ecological formations. NOTES: EXTIRPATED from southeastern Arizona, several reintroductions have taken place. *8 (Historically throughout south-eastern and south-central Arizona.), 14 (113006 - historically occurred in grass-shrub valleys and grasslands of southeastern and south-central Arizona), 55 (species: recorded as *Antilocapra americana* Ord. Prong-horned Antelope. Formerly widely distributed in grassland areas throughout the state; presently restricted to areas of favorable habitat.), 65 (species), 73 (species), 100 (color photograph of species), 106 (052806 - species), **118** (recorded as *Antilocapra americana mexicana* Merriam - Distribution: Southeastern Arizona. Figure 111, Page 255)*

Bovidae: The Cow, Sheep and Allies Family

***Ovis canadensis* G. Shaw: Rocky Mountain Bighorn Sheep**

COMMON NAMES: Berrego Cimarron (Hispanic), Bighorn, Bighorn Sheep, Desert Bighorn, Desert Bighorn Sheep, Mountain Sheep, Rocky Mountain Bighorn Sheep. HABITS: Feeds on agave, brittle bush, bursage, bush muhly, cacti, catclaw, cholla, coffeeberry, desert fluffgrass, desert ironwood, desert thorn, fairy duster, filaree, galleta, grama, jojoba, mesquite, mallow, Nevada joint fir, plantain, prickly-pear, ratany, ricegrass, saguaro, saltbush, threeawn and turpentine broom. Young are dropped in small scraped out depressions located in protected places on inaccessible peaks. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, **55** (recorded as *Ovis canadensis* Shaw. Bighorn. Probably formerly statewide in mountainous or rocky situations; presently restricted to scattered low desert mountains.), 65, 73, 100 (color photograph), 106 (052906), 118 (recorded as *Ovis canadensis mexicana* Merriam - Distribution: Probably formerly statewide in mountainous situations. Figure 112, Page 257)*

***Ovis canadensis* subsp. *mexicana* C.H. Merriam: Desert Bighorn Sheep**

COMMON NAMES: Berrego Cimarron (Hispanic), Berrego Cimarron del Desierto (Hispanic), Bighorn, Bighorn Sheep, Desert Bighorn, Desert Bighorn Sheep, Mountain Sheep, Rocky Mountain Bighorn Sheep. HABITS: The species feeds on agave, brittle bush, bursage, bush muhly, cacti, catclaw, cholla, coffeeberry, desert fluffgrass, desert ironwood, desert thorn, fairy duster, filaree, galleta, grama, jojoba, mesquite, mallow, Nevada joint fir, plantain, prickly-pear, ratany, ricegrass, saguaro, saltbush, threeawn and turpentine broom; young are dropped in small scraped out depressions located in protected places on inaccessible peaks. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (species: recorded as *Ovis canadensis* Shaw. Bighorn. Probably formerly statewide in mountainous or rocky situations; presently restricted to scattered low desert mountains.”), 65 (species), 73 (species), 100 (color photograph of species, species record), 106 (072306), **118** (recorded as *Ovis canadensis mexicana* Merriam - Distribution: Probably formerly statewide in mountainous situations. Figure 112, Page 257)*

Canidae: The Dog and Allies Family

***Canis latrans* T. Say: Coyote**

COMMON NAME: Coyote, Prairie Wolf. HABITS: Feeds on amphibians, berries, birds, carrion, fruits, gophers, insects, mice, rabbits, reptiles and squirrels. The young are born in dens that may be dug in the ground or located in caves. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, **55** (recorded as *Canis latrans* Say. Coyote. Statewide (120 - 9,100 feet.), 65 (color photograph), 73, 100 (color photograph), 106 (052906), 118 (recorded as *Canis latrans mearnsi* Merriam - Distribution: Statewide. Figure 87, Page 217)*

***Canis latrans* subsp. *mearnsi* Merriam: Coyote**

COMMON NAME: Coyote. HABITS: The species feeds on amphibians, berries, birds, carrion, fruits, gophers, insects, mice, rabbits, reptiles and squirrels. The young are born in dens that may be dug in the ground or located in caves. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (species), 55 (species: recorded as *Canis latrans* Say. Coyote. Statewide (120 - 9,100 feet.), 65 (color photograph of species, species record), 73 (species), 100 (color photograph of species, species record), 106 (052906 - species), **118** (recorded as *Canis latrans mearnsi* Merriam - Distribution: Statewide. Figure 87, Page 217)*

***Canis lupus* subsp. *baileyi* Nelson and Goldman: Mexican Gray Wolf**

COMMON NAMES: Lobo, Lobo Mexicano (Hispanic), Mexican Gray Wolf, Mexican Wolf, Timber Wolf. HABITS: Feeds on berries, birds, fish, fruits, insects, deer, elk, javelina, livestock, small mammals, bighorn sheep, pronghorn and rabbits. Maternity dens are chambers without nests usually located in the ground on high ground, under rock ledges, slopes of canyon walls or hills near water. HABITAT: Within the range of this species it has been reported from forest, woodland, grassland and wetland ecological formations. NOTES: The Mexican Gray Wolf is the smallest subspecies of gray wolf in North America. This wolf generally avoids desert areas. At one time the Mexican Gray Wolf was extirpated from Arizona; however, successful re-introduction efforts are bringing it back from near extinction. *8, 14 (082608), 55 (species: recorded as *Canis lupus* Frisch. Gray Wolf. Formerly throughout the eastern portions of the state, at present rare or approximately extinct.), 73 (species), 100 (species, color photograph of species), 106 (082608), 110 (recorded as *Canis lupus baileyi* - shows the historic range as being roughly that portion of Pima County east of the Tohono O'odham Nation), 118 (*Canis lupus baileyi* Nelson and Goldman - Distribution: Southeastern Arizona. Figure 88, Page 219)*

***Urocyon cinereoargenteus* (J.C. von Schreber): Common Gray Fox**

COMMON NAMES: Common Gray Fox, Gray Fox, Zorra Gris (Hispanic). HABITS: The species feeds on birds, desert cottontails, hackberry and prickly-pear fruits, grasses, insects (crickets and grasshoppers), juniper berries, lizards, manzanita berries, nuts, small rodents and snakes. Nests are made of bark, grasses and leaves and located in underground burrows, small caves, piles of rock, amongst boulders, crevices in cliffs and in hollows in trees. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: The Gray Fox climbs trees. *14 (082608 - subsp. *scottii* (Mearns)), **55** (recorded as *Urocyon cinereoargenteus* (Schreber). Gray Fox. Statewide with the possible exception of the northeast portion (120 - 5,800 feet.), 65 (species, color photograph), 73, 100 (color photograph), 106 (052906 - species with a listing of 16 subspecies), 118 (recorded as *Urocyon cinereoargenteus scottii* Mearns - Distribution: Probably statewide. Figure 90, Page 222)*

***Urocyon cinereoargenteus* subsp. *scottii* Mearns: Common Gray Fox**

COMMON NAMES: Common Gray Fox, Gray Fox, Zorra Gris (Hispanic). HABITS: The species feeds on birds, desert cottontails, hackberry and prickly-pear fruits, grasses, insects (crickets and grasshoppers), juniper berries, lizards, manzanita berries, nuts, small rodents and snakes. Nests are made of bark, grasses and leaves and located in underground burrows, small caves, piles of rock, amongst

boulders, crevices in cliffs and in hollows in trees. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: The Gray Fox climbs trees. *14 (082608 - subsp. *scottii* (Mearns)), 55 (species: recorded as *Urocyon cinereoargenteus* (Schreber). Gray Fox. Statewide with the possible exception of the northeast portion (120 - 5,800 feet).), 65 (species, color photograph of species), 73 (species), 100 (species, color photograph of species), 106 (052906 - species with a listing of 16 subspecies), 118 (recorded as *Urocyon cinereoargenteus scottii* Mearns - Distribution: Probably statewide. Figure 90, Page 222)*

***Vulpes macrotis* C.H. Merriam: Kit Fox**

COMMON NAMES: Kit Fox, Zorra del Desierto (Hispanic). HABITS: Feeds on berries, birds, cottontail rabbits, crickets, grasses, grasshoppers, ground squirrels, jack rabbits, kangaroo rats, lizards and pocket mice. The young are born in dens in underground burrows that have been excavated in soft soils. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: *Vulpes velox* (T. Say): The Swift Fox is generally considered a separate species by most authors. *14 (050907 - subspp. *macrotis* Merriam and *neomexicanus* Merriam), 55 (recorded as *Vulpes macrotis* Merriam. Kit Fox. Widely distributed at lower elevations throughout the southern part of the state (120 - 5,000 feet).), 65, 73, 78, 100 (color photograph), 106 (052906), 118 (recorded as *Vulpes macrotis arispus* Elliot - Distribution: Lower elevations in western and southern part of the state. *Vulpes macrotis neomexicana* Merriam - Distribution: Extreme southeastern Arizona. Figure 89, Page 220)*

***Vulpes macrotis* subsp. *arispus* Elliot: Kit Fox**

COMMON NAMES: Kit Fox, Zorra del Desierto (Hispanic). HABITS: The species feeds on berries, birds, cottontail rabbits, crickets, grasses, grasshoppers, ground squirrels, jack rabbits, kangaroo rats, lizards and pocket mice. The young are born in dens in underground burrows that have been excavated in soft soils. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: *Vulpes velox* (T. Say): The Swift Fox is generally considered a separate species by most authors. *14 (050907 - subspp. *macrotis* Merriam and *neomexicanus* Merriam), 55 (species: recorded as *Vulpes macrotis* Merriam. Kit Fox. Widely distributed at lower elevations throughout the southern part of the state (120 - 5,800 feet).) 65 (species), 73 (species), (100 - color photograph), 106 (052906), 118 (recorded as *Vulpes macrotis arispus* Elliot - Distribution: Lower elevations in western and southern part of the state. Figure 89, Page 220)*

Vulpes velox (see Note under *Vulpes macrotis*)

Cervidae: The Deer and Allies Family

***Odocoileus hemionus* (C.S. Rafinesque-Schmaltz): Mule Deer**

COMMON NAMES: Black-tailed Deer, Burro, Desert Mule Deer, Mule Deer, Venado Pardo (Hispanic). HABITS: Feeds on acorns, beans, branches, fruits, leaves or needles, nuts, seeds and/or twigs of aspen, barberry, bitterbrush, blackberry, buckbrush, buckwheat, calliandra, ceanothus, catclaw, cedar, cliffrose, dogwood, Douglas fir, huckleberry, joint fir, jojoba, juniper, mountain mahogany, mountainlover, oak, pinyon, ponderosa pine, poplar, sagebrush, saltbush, serviceberry, thimbleberry, white fir, wild cherry, willow and yew, and grasses lupines, mistletoe, moss, mushrooms, salal, sedges and spurges. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (recorded as *Odocoileus hemionus* (Rafinesque). Black-tailed or Mule Deer. Statewide, but not of uniform distribution (250 - 9,000 feet).), 65, 73, 100 (color photograph), 106 (052906), 118 (recorded as *Odocoileus hemionus crooki* (Mearns) - Distribution: Northeastern, central and southeastern part of the state. Figure 109, Page 252)*

***Odocoileus hemionus* subsp. *crooki* (Mearns): Mule Deer**

COMMON NAMES: Black-tailed Deer, Burro, Desert Mule Deer, Mule Deer, Venado Pardo (Hispanic). HABITS: The species feeds on acorns, beans, branches, fruits, leaves or needles, nuts, seeds and/or twigs of aspen, barberry, bitterbrush, blackberry, buckbrush, buckwheat, calliandra, ceanothus, catclaw, cedar, cliffrose, dogwood, Douglas fir, huckleberry, joint fir, jojoba, juniper, mountain mahogany, mountainlover, oak, pinyon, ponderosa pine, poplar, sagebrush, saltbush, serviceberry, thimbleberry, white fir, wild cherry, willow and yew, and grasses lupines, mistletoe, moss, mushrooms, salal, sedges and spurges. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (species), 55 (species: recorded as *Odocoileus hemionus* (Rafinesque) Black-tailed or Mule Deer. Statewide, but not of uniform distribution (250 - 9,000 feet).), 65, 73 (species), 100 (species, color photograph of species), 106 (052906 - species), **118** (recorded as *Odocoileus hemionus crooki* (Mearns) - Distribution: Northeastern, central and southeastern part of the state. Figure 109, Page 252)*

***Odocoileus virginianus* (Zimmermann): Coues' White-tailed Deer**

COMMON NAMES: Arizona Whitetail, Coues' Deer, Coues' White-tailed Deer, Desert Whitetail, Maso (Yaqui), Fantail, Sonora White-tailed Deer, Sonoran Fantail, Venado Cola Blanca (Hispanic), Virginia Deer, Whitetail, White-tailed Deer, Whitetail Deer. HABITS: The species feeds on fungi, grass and acorns, branches, buds, cones, fruits, leaves, mast, needles and /or twigs of alder, barberry, buckbrush, calliandra, catclaw acacia, Emory and scrub oaks and other evergreen oaks, hackberry, hemlock, holly-leaf buckthorn, juniper, mesquite, mountainlover, Oregon-grape, pinyon, ratany, sagebrush, skunkbush, spiderwort, spruce, willow, yellow-leaf silktassel. Young are generally dropped along ridges and hillsides. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, **55** (recorded as *Odocoileus virginianus* (Zimmermann). White-tailed Deer. Southeastern Arizona (1,200 - 9,000 feet).), 65, 73, 100 (color photograph), 106 (052906), 118 (recorded as *Odocoileus virginianus couesi* (Coues & Yarrow) - Distribution: Southern Arizona. Figure 110, Page 254)*

***Odocoileus virginianus* subsp. *couesi* (E. Coues & Yarrow): Coues' White-tailed Deer**

COMMON NAMES: Arizona Whitetail, Coues' Deer, Coues' White-tailed Deer, Desert Whitetail, Fantail, Maso (Yaqui), Sonora White-tailed Deer, Sonoran Fantail, Venado Cola Blanca (Hispanic), Virginia Deer, Whitetail, White-tailed Deer, Whitetail Deer. HABITS: The species feeds on fungi, grass and acorns, branches, buds, cones, fruits, leaves, mast, needles and /or twigs of alder, barberry, buckbrush, calliandra, catclaw acacia, Emory and scrub oaks and other evergreen oaks, hackberry, hemlock, holly-leaf buckthorn, juniper, mesquite, mountainlover, Oregon-grape, pinyon, ratany, sagebrush, skunkbush, spiderwort, spruce, willow, yellow-leaf silktassel. Young are generally dropped along ridges and hillsides. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (species: recorded as *Odocoileus virginianus* (Zimmermann). White-tailed Deer. Southeastern Arizona (1,200 - 9,000 feet).), 65, 73 (species), 100 (species, color photograph of species), 106 (052906 - species), **118** (recorded as *Odocoileus virginianus couesi* (Coues & Yarrow) - Distribution: Southern Arizona. Figure 110, Page 254)*

Felidae: The Cat Family

***Felis concolor* (C. Linnaeus): Mountain Lion**

SYNONYMY: *Puma concolor* (C. Linnaeus). COMMON NAMES: American Lion, Brown Tiger, California Lion, Cat-a-Mountain, Catamount, Catamount Cat (a mountain Red Tiger), Cougar, Deer Tiger, El Leon (Mexico), Florida Panther, Ghost Cat, Indian Devil, King Cat, Leon de Montana

(Hispanic), Mexican Lion, Mountain Lion, Mountain Screamer, Painted Cat, Painter, Panther; Puma (Indian), Ted Tiger (Belize), Silver Lion, Sneak Cat, Sucuarana (Brazil), Yuma Mountain Lion. HABITS: Feeds on beavers, bighorn sheep, birds, black bears, bobcats, cottontail rabbits, coyotes, deer (its major prey species in Arizona), elk, jackrabbits, javelina, livestock, porcupines, pronghorn, raccoons, skunks and small mammals. Kittens are born in dens located in protected areas such as shallow caves, crevices, downed logs, rock shelters and impenetrable thickets. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: Running should be curtailed in areas where Mountain Lions are known to frequent, a person running may elicit an attack response from a nearby Mountain Lion. Mountain Lions are extremely agile and have great jumping power and have been reported as being able to leap to a height of 18 feet into a tree. *8 (*Puma concolor* (Linnaeus)), 14 (091108 - *Puma concolor* subsp. *azteca* (Merriam); *hippolestes* (Merriam); *kaibabensis* (Nelson and Goldman), and *stanleyana* (Goldman). The Yuma Mountain Lion (*Felis concolor browni*) is included as a separate record.), 55 (recorded as *Felis concolor* Linnaeus. Mountain Lion. Statewide (200 - 8,000 feet.), 65, 73, 100 (color photograph), 106 (052906), 118 (recorded as *Felis concolor azteca* Merriam - Distribution: Statewide except extreme western and northwestern parts; *Felis concolor browni* (Merriam) - Distribution: Southwestern part of the state, and *Felis concolor kaibabensis* Nelson and Goldman - Distribution: Northwestern Arizona, north and west of the Colorado River. Figure 105, Page 245)*

Felis onca subsp. *arizonensis* (see *Panthera onca* subsp. *arizonensis*)

Felis pardalis subsp *sonoriensis* (see *Leopardus pardalis* subsp *sonoriensis*)

Felis rufus (see *Lynx rufus*)

Felis rufus subsp. *baileyi* (see *Lynx rufus* subsp. *baileyi*)

***Leopardus pardalis* subsp *sonoriensis* Goldman: Ocelot**

SYNONYMY: (*Felis pardalis* Linnaeus, *Felis pardalis* subsp. *sonoriensis* Goldman). COMMON NAMES: Jaguatirica (Brazil), Manigordo (Costa Rica), McKenney's Wildcat, Ocelot, Painted Leopard, Tigrillo. HABITS: (Feeds on amphibians, lesser anteaters, armadillos, birds, fish, insects, land crabs, small to medium-sized mammals (including mice, rats and rabbits among others) and reptiles (including lizards, snakes and land tortoises). Kittens are born in a nest lined with grass or other materials located in rocky bluffs, caves, rocky dens, hollow logs or dense thickets. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: EXTIRPATED. *8 (species), 14 (091108 - subsp *sonoriensis*), 55 (species: recorded as *Felis pardalis* Linnaeus. Ocelot. Formerly southeastern Arizona as far north as Fort Verde; no recent records.), 100 (species, color photograph of species), 106 (091108 - includes a listing with location of subspecies), 118 (recorded as *Felis pardalis* subsp. *sonoriensis* Goldman - Distribution: Formerly southeastern Arizona as far north as Ft. Verde. Figure 104, Page 244)*

***Lynx rufus* (J.C. von Schreber): Bobcat**

SYNONYMY: *Felis rufus* (J.C. von Schreber). COMMON NAMES: Bobcat, Gato Montes (Hispanic), Wildcat. HABITS: Feeds on almost any meat source available including ground nesting birds, carrion, domestic cats, cottontail rabbits, deer, foxes, jackrabbits, lizards, small mammals, opossums, porcupines, raccoons, reptiles, rodents, bighorn sheep, skunks and woodchucks. Shelter may be taken in a rock cleft, thickets or on the branches of trees. Young are born in dens located in rocky caves, rock shelters, recesses and protected areas with nests made of leaves and other dry plant material. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (091108 - *Lynx rufus baileyi* Schreber), 55 (recorded as *Lynx rufus* (Schreber). Bobcat. Statewide (120 - 9,300 feet.), 65, 73, 100 (color photograph),

106 (052906), 118 (recorded as *Lynx rufus baileyi* Merriam - Distribution: Statewide. Figure 106, Page 247)*

***Lynx rufus* subsp. *baileyi* Merriam: Bobcat**

SYNONYMY: *Felis rufus* subsp. *baileyi* Elliot. COMMON NAMES: Bobcat, Gato Montes (Hispanic), Wildcat. HABITS: Feeds on almost any meat source available including ground nesting birds, carrion, domestic cats, cottontail rabbits, deer, foxes, jackrabbits, lizards, small mammals, opossums, porcupines, raccoons, reptiles, rodents, bighorn sheep, skunks and woodchucks. Shelter may be taken in a rock cleft, thickets or on the branches of trees. Young are born in dens located in rocky caves, rock shelters, recesses and protected areas with nests made of leaves and other dry plant material. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (091108 - *Lynx rufus baileyi* Schreber), 55 (species: recorded as *Lynx rufus* (Schreber). Bobcat. Statewide (120 - 9,300 feet.), 65, 73 (species), 100 (species, color photograph of species), 106 (052906 - spies), **118** (recorded as *Lynx rufus baileyi* Merriam - Distribution: Statewide. Figure 106, Page 247)*

***Panthera onca* subsp. *arizonensis* Goldman: Jaguar**

SYNONYMY: *Felis onca* subsp. *arizonensis* Goldman. COMMON NAMES: Black Panther, Blank Panther, Jaguar, Jaguar (Hispanic), Jaguarete (Spanish), Yaguar. HABITS: Feeds on armadillos, birds, caiman, capybaras, deer, fish, frogs, livestock, pacas, peccaries (javelina), mice, rabbits, tapirs, turtles and other vertebrates. Young are born in dens located in caves, rocky areas, dense brush and thickets. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: EXTIRPATED from Arizona. *8, 14 (091008 - *Panthera onca* subsp. *arizonensis* Goldman), 55 (species: recorded as *Felis onca* Linnaeus. Jaguar. Probably formerly rare throughout the state. Today an occasional individual is found in the southern part of the state.), 65 (species), 100 (species, color photograph of species), 106 (052906), **118** (recorded as *Felis onca arizonensis* Goldman - Distribution: Probably formerly rare throughout the state. Today an occasional individual found in the southern part of the state. Figure 104, Page 244)*

Puma concolor (see *Felis concolor*)

Puma yaguarondi (see *Herpailurus yaguarondi*)

Geomyidae: The Pocket Gopher Family

***Thomomys bottae* subsp. *modicus* Goldman: Botta's Pocket Gopher**

COMMON NAMES: Botta's Pocket Gopher, Southwestern Pocket Gopher, Tuza de Botta (Hispanic), Valley Pocket Gopher. HABITS: The species feeds on bulbs, grasses, herbaceous plants, roots and tubers. Young are born in nests in underground burrows. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051107 - species, several varieties listed), 55 (species: recorded as *Thomomys bottae* (Eydoux and Gervais). Valley Pocket Gopher. Widely distributed throughout the state at all elevations.), 65 (species), 73 (species), 100 (species, color photograph of species), 106 (052906 - species), **118** (recorded as *Thomomys bottae modicus* Goldman - Distribution: Known from the Santa Cruz and Altar Valleys. Figure 46, Page 107)*

Heteromyidae: The Kangaroo Rat and Pocket Mouse Family

***Chaetodipus baileyi* subsp. *baileyi* C.H. Merriam: Bailey's Pocket Mouse**

SYNONYMY: *Perognathus baileyi* subsp. *baileyi* C.H. Merriam. COMMON NAMES: Bailey's Pocket Mouse, Raton de Bailey (Hispanic). HABITS: The species feeds on vegetation, and fruits and seeds of cacti, grasses and other herbs. Nests are located underground burrows. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, and desertscrub ecological formations. *14 (082508 - subsp. *baileyi* Merriam), 55 (species: recorded as *Perognathus baileyi* Merriam. Bailey's Pocket Mouse. Widely distributed in the southern part of the state (900 - 4,700 feet.), 65 (genus), 73 (species), 100 (species, color photograph of species), 106 (082508), **118** (recorded as *Chaetodipus baileyi baileyi* Merriam - Distribution: Grasslands of southeastern Arizona. Figure 51, Page 133)*

***Chaetodipus hispidus* subsp. *conditi* Allen: Hispid Pocket Mouse**

SYNONYMY: *Perognathus hispidus* subsp. *conditi* Allen S.F. Baird. COMMON NAME: Hispid Pocket Mouse. HABITS: The species feeds on insects (grasshoppers), leaves and seeds. Nests are constructed of grasses and located in underground burrows. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (082508 - subsp. *conditi*), 55 (species: recorded as *Perognathus hispidus* Baird. Hispid Pocket Mouse. Locally common in grasslands of southeastern part of the state; an isolated population occurs near Camp Verde (3,200 - 5,000 feet.), 65 (genus), 73 (species), 100 (species), 106 (082508 - species), **118** (recorded as *Perognathus hispidus conditi* Allen - Distribution: Grasslands of southeastern Arizona. Figure 51, Page 132)*

***Chaetodipus intermedius* subsp. *intermedius* C.H. Merriam: Rock Pocket Mouse**

SYNONYMY: *Perognathus intermedius* subsp. *intermedius* C.H. Merriam. COMMON NAMES: Raton de Rocas de Bosla (Hispanic), Rock Pocket Mouse. HABITS: The species feeds on seeds. Burrows are dug in soil near to or under rocks. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051007), 55 (species: recorded as *Perognathus intermedius* Merriam. Rock Pocket Mouse. Widely distributed in rocky areas in the Colorado River valley, western and southern Arizona (120 - 6,000 feet.), 65 (genus), 73 (species - recorded as *Perognathus intermedius*), 100 (species), 106 (051007 - species), **118** (recorded as *Chaetodipus intermedius intermedius* Merriam - Distribution: Known from Mohave County southward and eastward, across most of the state to Cochise County. Figure 54, Page 141)*

***Chaetodipus penicillatus* subsp. *pricei* (S.W. Woodhouse): Desert Pocket Mouse**

SYNONYMY: *Perognathus penicillatus* subsp. *pricei* S.W. Woodhouse). COMMON NAMES: Desert Pocket Mouse, Raton de Desierto (Hispanic), Sonoran Desert Pocket Mouse. HABITS: The species feeds on seeds of creosote bush, grass, greythorn, herbs and mesquite. The nest is made in underground burrows. HABITAT: Within the range of this species it has been reported from the scrub, grassland, desertscrub and wetland ecological formations. *14 (051007), 55 (species: recorded as *Perognathus penicillatus* Woodhouse. Desert Pocket Mouse. Widely distributed in desert and low grasslands of southern and western Arizona (120 - 5,200 feet.), 65 (genus), 73 (species, *Perognathus penicillatus*), 100 (species, color photograph of species), 106 (051007 - species), **118** (recorded as *Perognathus penicillatus pricei* Allen - Distribution: Known from south-central Arizona. Figure 53, Page 137)*

***Dipodomys merriami* subsp. *merriami* Mearns: Merriam's Kangaroo Rat**

COMMON NAMES: Merriam's Kangaroo Rat, Rata de Nopalera Merriam (Hispanic). HABITS: The species feeds on ants, green plant material and seeds of creosote bush, grama grass, mesquite, ocotillo and purselane. Nests are made in underground burrows often located under bushes. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051007), 55 (species: recorded as *Dipodomys merriami* Mearns. Merriam's Kangaroo Rat. Widely distributed in western and southern parts of the state (120 - 5,000

feet.), 65 (species, color photograph of species), 73 (species), 100 (species, color photograph of species), 106 (genus with a listing of species), **118** (recorded as *Dipodomys merriami merriami* Mearns - Distribution: Occurs throughout most of the western and southern part of the state. Figure 56, Page 145)*

***Dipodomys spectabilis* subsp. *perblandus* Goldman: Banner-tailed Kangaroo Rat**

COMMON NAMES: Banner-tailed Kangaroo Rat, Kangaroo Rat, Rata de Nopalera (Hispanic). HABITS: The species feeds on grasses, forbs, succulent plants, insects, rodents and seeds. Nests are made up of chaff, stems and leaves of grass located in underground burrows in firm soils. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051007), 55 (species: recorded as *Dipodomys spectabilis* Merriam. Banner-tailed Kangaroo Rat. Locally common in grasslands of southeastern Arizona (1,300 - 5,000 feet.), 65 (species, color photograph), 100 (species, color photograph), 106 (genus, listing of species), **118** (recorded as *Dipodomys spectabilis perblandus* Goldman - Distribution: Known from the grasslands of southern Pinal and Pima County. Figure 55, Page 143)*

***Perognathus amplus* subsp. *taylori* Goldman: Arizona Pocket Mouse**

COMMON NAME: Arizona Pocket Mouse. HABITS: The species feeds on green plants, insects and seeds. Nests are located in underground burrows. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *55 (species: recorded as *Perognathus amplus* Osgood. Arizona Pocket Mouse. Locally common in desert areas on south-central, western and north-central parts of the state (500 - 5,100 feet.), 65 (genus), 73 (species), 100 (species, color photograph of species), **118** (recorded as *Perognathus amplus taylori* Goldman. Distribution: Known from south central Arizona. Figure 50, Page 129)*

***Perognathus flavus* subsp. *flavus* S.F. Baird: Silky Pocket Mouse**

COMMON NAME: Silky Pocket Mouse. HABITS: The species feeds on seeds and invertebrates (though very few are taken). Nests are located in underground burrows. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (082508 - subsp. *flavus* (Baird)), 55 (species: recorded as *Perognathus flavus* Baird. Silky Pocket Mouse. Locally common in grasslands throughout the state (2,900 - 6,500 feet.), 65 (genus), 73 (species), 100 (species, color photograph of species), 106 (082508), **118** (recorded as *Perognathus flavus flavus* Baird - Distribution: Southeastern part of the state. Figure 48, Page 124)*

Perognathus baileyi subsp. *baileyi* (see *Chaetodipus baileyi* subsp. *baileyi*)

Perognathus hispidus subsp. *conditi* (see *Chaetodipus hispidus* subsp. *conditi*)

Perognathus intermedius subsp. *intermedius* (see *Chaetodipus intermedius* subsp. *intermedius*)

Perognathus penicillatus subsp. *pricei* (see *Chaetodipus penicillatus* subsp. *pricei*)

Leporidae: The Hare and Rabbit Family

***Lepus alleni* subsp. *alleni* Mearns: Antelope Jack Rabbit**

COMMON NAME: Antelope Jack Rabbit. HABITS: The species feeds on cacti, Catclaw Acacia, grasses, herbs and the bark, buds and leaves of mesquite. Young are born in a nest that is usually located above ground. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (species), 55 (species: recorded as *Lepus alleni* (Mearns). Antelope Jack Rabbit. Occurs in the central third of the southern half of the state.), 65 (species), 73 (species), 100 (species, color photograph of species), 106 (052906 - species), **118** (recorded

as *Lepus alleni* subsp. *alleni* Mearns - Distribution: Occurs in the central third of the southern half of the state. Figure 31, Page 68)*

***Lepus californicus* (J.E. Gray): Black-tailed Jack Rabbit**

COMMON NAMES: Black-tailed Jack Rabbit, “Jackass Rabbit”. HABITS: Feeds on grass, mesquite leaves and prickly-pear cacti. Young are born in nests located either above or below ground in forms that have been lined with breast hair, after birth the young are moved to separate nests and cared for individually by the female. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (recorded as *Lepus californicus* Gray. Black-tailed Jack Rabbit. Statewide.), 65, 73, 100 (color photograph), 106 (052906), 118 (recorded as *Lepus californicus deserticola* Mearns - Distribution: Occurs in the western half of the state; *Lepus californicus eremicus* J.A. Allen - Distribution: Southeastern Arizona, and *Lepus californicus texianus* Waterhouse - Distribution: Occurs in the northeastern quarter of the state. Figure 32, Page 69)*

***Lepus californicus* subsp. *eremicus* J.A. Allen: Black-tailed Jack Rabbit**

COMMON NAMES: Black-tailed Jack Rabbit, “Jackass Rabbit”. HABITS: The species feeds on grass, mesquite leaves and prickly-pear cacti. Young are born in nests located either above or below ground in forms that have been lined with breast hair, after birth the young are moved to separate nests and cared for individually by the female. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (species), 55 (species: recorded as *Lepus californicus* Gray. Black-tailed Jack Rabbit. Statewide.), 65 (species), 73 (species), 100 (species, color photograph of species), 106 (052906 - species), 118 (recorded as *Lepus californicus eremicus* J.A. Allen - Distribution: Southeastern Arizona. Figure 32, Page 69)*
Lepus californicus subsp. *eremicus* J.A. Allen: Black-tailed Jack Rabbit

***Sylvilagus audubonii* (S.F. Baird): Desert Cottontail**

COMMON NAME: Desert Cottontail. HABITS: Feeds on green plants, cacti, bark and twigs. Young are born into nests lined with forbs, grasses and the female’s fur which are located on the ground and in brush piles, piles of rocks, and burrows abandoned by other animals. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (recorded as *Sylvilagus audubonii* (Baird). Desert Cottontail. Common at elevations below 6,000 feet throughout the state.), 65, 73, 100 (color photograph), 106 (052906), 118 (recorded as *Sylvilagus audubonii arizonae* (J.A. Allen) - Distribution: Widely distributed at elevations up to 6,000 feet in the western half of the state; *Sylvilagus audubonii minor* (Mearns) - Distribution: Known only from the southeastern part of the state, and *Sylvilagus audubonii warreni* Nelson - Distribution: Known only from the northeastern part of the state. Figure 34, Page 74), WTK (March 9, 2007)*

***Sylvilagus audubonii* (S.F. Baird) subsp. *arizonae*: Desert Cottontail**

COMMON NAME: Desert Cottontail. HABITS: The species feeds on green plants, cacti, bark and twigs. Young are born into nests lined with forbs, grasses and the female’s fur which are located on the ground and in brush piles, piles of rocks, and burrows abandoned by other animals. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (species), 55 (species: recorded as *Sylvilagus audubonii* (Baird). Desert Cottontail. Common at elevations below 6,000 feet throughout the state.), 65 (species), 73 (species), 100 (species, color photograph of species), 106 (052906 - species), 118 (recorded as *Sylvilagus audubonii arizonae* (J.A. Allen) - Distribution: Widely distributed at elevations up to 6,000 feet in the western half of the state. Figure 34, Page 74)*

***Conepatus leuconotus* subsp. *venaticus* (Goldman): Common Hog-nosed Skunk**

SYNONYMY: *Conepatus mesoleucus* subsp. *venaticus* Goldman. COMMON NAMES: Common Hog-nosed Skunk, Hog-nosed Skunk, Hognose Skunk, Rooter Skunk, Zorrillo Nariz de Puerco (Hispanic). HABITS: The species feeds on arachnids, birds, insects, small mammals, mollusks, plant material, reptiles and worms. These skunks take refuge in caves, crevices in rocks and in the ground. Rocky areas are used for denning with the young born beneath rocks, grasses are used for nesting. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (072306 - subsp. *venaticus* (Goldman)), 55 (species: recorded as *Conepatus mesoleucus* Lichtenstein. Hog-nosed Skunk. Southeastern part of the state (2,000 - 6,000 feet.), 65 (species), 73 (species: *Conepatus mesoleucus*), 100 (species record (*Conepatus mesoleucus*), color photograph of species), 106 (072306 - genus), **118** (recorded as *Conepatus mesoleucus venaticus* Goldman - Distribution: South central and southeastern Arizona. Figure 102, Page 241)*

Conepatus mesoleucus subsp. *venaticus* (see *Conepatus leuconotus* subsp. *venaticus*)

***Mephitis macroura* subsp. *milleri* (Mearns): Hooded Skunk**

COMMON NAMES: Hooded Skunk, Zorrillo (Hispanic). HABITS: The species feeds on small birds, insects and other invertebrates, rodents and plant material. The young are born in a dens located in burrows or among rocks. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050807 - subsp. *milleri* (Mearns)), 55 (species: recorded as *Mephitis macroura* (Lichtenstein). Hooded Skunk. Southeastern part of the state (2,000 - 6,000 feet.), 65 (species), 73 (species), 100 (species, color photograph of species), 106 (053006 - genus), **118** (recorded as *Mephitis macroura milleri* (Mearns) - Distribution: South central and southeastern Arizona. Figure 101, Page 240)*

***Mephitis mephitis* (J.C. von Schreber): Striped Skunk**

COMMON NAMES: Striped Skunk, Zorrillo Rayado (Hispanic). HABITS: Feeds on amphibians, berries, the eggs of ground nesting birds, carrion, crayfish, earthworms, fishes, fruits, insects (beetles, crickets and grasshoppers among others), mollusks, plant material, reptiles, rodents, snails and spiders. The young are born in nests made of dried grasses and leaves located in dirt banks, underground burrows abandoned by other animals, downed logs, pits and rock outcrops. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: The Striped Skunk is most active dusk through dawn. This species is the chief carrier of rabies in the United States and those active during the daylight hours frequently being found to be rabid. *14 (082308 - subsp. *estor* Merriam, *hudsonica* (Richardson) and *varians* (Gray)), **55** (recorded as *Mephitis mephitis* (Schreber). Striped Skunk. Statewide (300 - 9,000 feet.), 65 (color photograph), 73, **100** (color photograph), 106 (053006 - genus), 118 (recorded as *Mephitis mephitis estor* Merriam - Distribution: Statewide. Figure 100, Page 239)*

***Mephitis mephitis* (J.C. von Schreber) subsp. *estor* Merriam: Striped Skunk**

COMMON NAMES: Striped Skunk, Zorrillo Rayado (Hispanic). HABITS: Feeds on amphibians, berries, the eggs of ground nesting birds, carrion, crayfish, earthworms, fishes, fruits, insects (beetles, crickets and grasshoppers among others), mollusks, plant material, reptiles, rodents, snails and spiders. The young are born in nests made of dried grasses and leaves located in dirt banks, underground burrows abandoned by other animals, downed logs, pits and rock outcrops. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: The Striped Skunk is most active dusk through dawn. This species is the chief carrier of rabies in the United States and those active during the daylight hours frequently being found to be rabid. *14 (082308 - subsp. *estor* Merriam), 55 (species: recorded as *Mephitis mephitis* (Schreber). Striped Skunk. Statewide (300 - 9,000 feet.), 65 (species, color photograph

of species), 73 (species), 100 (species, color photograph of species), 106 (053006 - genus), 118 (recorded as *Mephitis mephitis estor* Merriam - Distribution: Statewide. Figure 100, Page 239)*

***Spilogale gracilis* Merriam: Western Spotted Skunk**

SYNONYMY: *Spilogale putorius* subsp. *gracilis* Merriam. COMMON NAMES: Spotted Skunk, Western Spotted Skunk, Zorillo Pinto (Hispanic). HABITS: Feeds on arachnids, berries, birds and bird eggs, carrion, fruits, insects, small mammals, scorpions and seeds. Dens are made in rock crevices and hollow logs. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051107 - *Spilogale putorius* subsp. *gracilis* Merriam is a synonym for *Spilogale gracilis* C. Linnaeus the Western Spotted Skunk. *Spilogale putorius* subsp. *leucoparia* is a synonym for *Spilogale putorius* C. Linnaeus the Eastern Spotted Skunk), 55 (recorded as *Spilogale putorius* (Linnaeus). Spotted Skunk. Probably statewide (120 - 7,000 feet.), 65 (*Spilogale putorius*), 73 (recorded as *Spilogale gracilis*), 100 (recorded as *Spilogale gracilis*, color photograph), 106 (053006 - genus), 118 (recorded as *Spilogale putorius gracilis* Merriam - Distribution: Probably statewide. Figure 99, Page 237)*

Spilogale putorius (see footnotes 14, 55, 65 and 85 under *Spilogale gracilis*)

Spilogale putorius subsp. *gracilis* (see *Spilogale gracilis*)

Molossidae: The Free-tailed Bat Family

***Eumops perotis* subsp. *californicus* (Merriam): Greater Western Mastiff Bat**

COMMON NAMES: Bonnet Bat, Greater Western Bonneted Bat, Greater Mastiff Bat, Greater Western Mastiff Bat, Mastiff Bat, Murcielago Mastiff (Hispanic), Western Mastiff Bat. HABITS: The species feeds on crickets, long-horned grasshoppers, moths and other small insects. Roosts in crevices and shallow caves in cliffs and rock walls at lower elevations. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (051107), 55 (species: recorded as *Eumops perotis* (Schinz). Western Mastiff Bat. Rare; in small colonies in rock crevices at lower elevations in the western and southern part of the state.), 65 (species), 73 (species), 92 (species), 100 (species, color photograph of species), 106 (053006 - family), 118 (recorded as *Eumops perotis californicus* (Merriam) - Distribution: Probably throughout southern Arizona in the Lower Sonoran Life Zone. Figure 29, Page 65)*

***Nyctinomops femorosaccus* (C.H. Merriam): Pocketed Free-tailed Bat**

SYNONYMY: Also recorded as *Nyctinomops femorosacca* (C.H. Merriam), *Tadarida femorosacca* (Miller). COMMON NAMES: Pocketed Free-tailed Bat, Murcielago Cola en Bolsa (Hispanic). HABITS: Feeds on ants, leafhoppers, moths, wasps and other insects. Roosts in rocky crevices. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14, 55 (recorded as *Tadarida femorosacca* (Merriam). Pocketed Free-tailed Bat. Rare; found at lower elevations in the western and southern part of the state.), 100, 106 (053006 - family), 118 (recorded as *Tadarida femorosacca* (Miller) - Distribution: Probably occurs throughout the Lower Sonoran Life Zone of southern Arizona. Figure 27, Page 63)*

***Nyctinomops macrotis* (J.E. Gray): Big Free-tailed Bat**

SYNONYMY: *Tadarida macrotis* (J.E. Gray), *Tadarida molossa* (Pallas). COMMON NAMES: Big Free-tailed Bat, Murcielago Cola Libre (Hispanic), Murcielago Cola Suelta Mayor (Spanish) HABITS: Feeds on insects. Roosts in rocky cliffs, crevices, fissures, caves and holes in trees. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland,

desertscrub and wetland ecological formations ecological formations. *8, 14, 42 (053006), 55 (recorded as *Tadarida molossa* (Pallas). Big Free-tailed Bat. Rare; statewide, mainly at elevations below 5,000 feet.), 73, 100 (color photograph), 106 (053006 - family), 118 (recorded as *Tadarida molossa* (Pallas) - Distribution: Probably occurs throughout the Lower Sonoran Life Zone of Arizona. Figure 28, Page 64)*

***Tadarida brasiliensis* (I.G. Saint-Hilaire) (subsp *mexicana* (Saussure) is the only subspecies reported as occurring in Arizona): Brazilian Free-tailed Bat**

COMMON NAMES: Brazilian Free-tailed Bat, Guano Bat, Mexican Free-tail Bat, Mexican Free-tailed Bat, Murcielago Brasileiro (Hispanic). HABITS: Feeds on ants, beetles, leafhoppers, moths and other small insects. Roosts in caverns; caves; crevices in rocks; fissures in cliffs; buildings; mines, and under bridges. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14, 55 (recorded as *Tadarida brasiliensis* (I.Geof. St.-Hilaire). Mexican Free-tailed Bat. Locally abundant throughout the state, especially at elevations below 5,000 feet.), 65, 73, 92, 100 (color photograph), 106 (053006), 118 (recorded as *Tadarida brasiliensis mexicana* (Saussure) - Distribution: Probably statewide in some part of the year. Figure 26, Page 62)*

***Tadarida brasiliensis* (I.G. Saint-Hilaire) subsp *mexicana* (Saussure): Brazilian Free-tailed Bat**

COMMON NAMES: Brazilian Free-tailed Bat, Guano Bat, Mexican Free-tail Bat, Mexican Free-tailed Bat, Murcielago Brasileiro (Hispanic). HABITS: Feeds on ants, beetles, leafhoppers, moths and other small insects. Roosts in caverns; caves; crevices in rocks; fissures in cliffs; buildings; mines, and under bridges. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14, 55 (species: recorded as *Tadarida brasiliensis* (I.Geof. St.-Hilaire). Mexican Free-tailed Bat. Locally abundant throughout the state, especially at elevations below 5,000 feet.), 65 (species), 73 (species), 92 (species), 100 (species, color photograph of species), 106 (053006), 118 (recorded as *Tadarida brasiliensis mexicana* (Saussure) - Distribution: Probably statewide in some part of the year. Figure 26, Page 62)*

Tadarida femorosacca (see *Nyctinomops femorosacca*)

Tadarida macrotis (see *Nyctinomops macrotis*)

Tadarida molossa (see *Nyctinomops macrotis*)

Muridae: The Mouse and Rat Family

***Neotoma albigula* subsp. *albigula* Hartley: White-throated Wood Rat**

COMMON NAMES: Packrat, White-throated Packrat, Trade Rat, White-throated Wood Rat. HABITS: The species feeds on cacti, forbs, fruits, juniper, leaves, mesquite beans, seeds and yucca. Nests are built under mesquite, cholla and prickly-pear cacti, or in rocky crevices using sticks, pieces of cholla and prickly-pear cacti, and rubbish, sometimes with underground burrows. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051107 - subsp. *albigula*), 55 (species: recorded as *Neotoma albigula* Hartley. White-throated Wood Rat. Widely distributed at elevations below 7,000 feet throughout all of the state south of the Colorado River (120 - 8,000 feet.), 65 (species, color photograph of species), 73 (species), 100 (species, color photograph of species), 106 (053006 - genus), 118 (recorded as *Neotoma albigula albigula* Hartley - Distribution: Occurs commonly south of the Mogollon Rim. Figure 76, Page 193)*

***Onychomys torridus* subsp. *torridus* (E. Coues): Southern Grasshopper Mouse**

COMMON NAMES: Raton Chapulinero del Sur (Hispanic), Scorpion Mouse, Southern Grasshopper Mouse. HABITS: The species feeds on arthropods, beetles, grasshoppers, insects, lizards, other species of mice, scorpions, seeds and small vertebrates. Nests are located in underground burrows. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (020307 - subsp. *torridus*), 55 (species: recorded as *Onychomys torridus* (Coues). Southern Grasshopper Mouse. Widely distributed in the western and southern parts of the state (120 - 5,000 feet.), 65 (genus), 73 (species), 100 (species, color photograph of species), 106 (053006 - genus, listing of species), 118 (recorded as *Onychomys torridus torridus* (Coues) - Distribution: Southeastern quarter of the state. Figure 62, Page 161)*

***Peromyscus eremicus* subsp. *eremicus* (S.F. Baird): Cactus Mouse**

COMMON NAMES: Cactus Mouse, Raton de Cactaceas (Hispanic). HABITS: The species feeds on flowers, small fruits, insects, green plant material and seeds. Nests are made within the abandoned burrows of other animals, clumps of cacti and among rocks. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (species), 55 (species: recorded as *Peromyscus eremicus* (Baird). Cactus Mouse. Widely distributed in western and southern Arizona (120 - 6,000 feet.), 65 (genus), 73 (species), 100 (species, color photograph of species), 106 (053006 - genus), 118 (recorded as *Peromyscus eremicus eremicus* (Baird) - Distribution: Almost all of the western and southern part of the state. Figure 67, Page 171)*

***Peromyscus maniculatus* (Wagner): Deer Mouse**

COMMON NAMES: Deer Mouse, Raton Venado (Hispanic). HABITS: Feeds on bark, berries, bones, centipedes, earthworms, small fruits, fungi, insects, leaves, nuts and snails. Nests are built in buildings, underground burrows, rock crevices debris, in and under logs, and clumps of vegetation. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *blandus* (Osgood) and *rufinus* (Merriam)), 55 (recorded as *Peromyscus maniculatus* (Wagner). Deer Mouse. Statewide (120 - 11,400 feet.), 65 (genus), 73, 100 (color photograph), 106 (053006 - genus), 118 (recorded as *Peromyscus maniculatus blandus* Osgood - Distribution: Extreme southeastern part of the state; *Peromyscus maniculatus rufinus* (Merriam) - Distribution: Higher elevations throughout the state, and *Peromyscus maniculatus sonoriensis* (Le Conte) - Distribution: Grasslands at lower elevations throughout the state. Figure 69, Page 177)*

***Peromyscus maniculatus* subsp. *sonoriensis* (Le Conte): Deer Mouse**

COMMON NAMES: Deer Mouse, Raton Venado (Hispanic). HABITS: The species feeds on bark, berries, bones, centipedes, earthworms, small fruits, fungi, insects, leaves, nuts and snails. Nests are built in buildings, underground burrows, rock crevices debris, in and under logs, and clumps of vegetation. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - species, and subsp. *blandus* (Osgood) and *rufinus* (Merriam)), 55 (species: recorded as *Peromyscus maniculatus* (Wagner). Deer Mouse. Statewide (120 - 11,400 feet.), 65 (genus), 73 (species), 100 (species, color photograph of species), 106 (053006 - genus), 118 (recorded as *Peromyscus maniculatus sonoriensis* (Le Conte) - Distribution: Grasslands at lower elevations throughout the state. Figure 69, Page 177)*

***Peromyscus merriami* subsp. *merriami* Mearns: Merriam's Mouse**

COMMON NAMES: Merriam's Mouse, Mesquite Mouse. HABITS: The species probably feeds on invertebrates and seeds. HABITAT: Within the range of this species it has been reported from the grassland, desertscrub and wetland ecological formations. *14 (082308), 55 (species: recorded as *Peromyscus merriami* Mearns. Merriam's Mouse. Known from scattered localities is Pinal, Pima and Santa Cruz counties (1,600 - 3,600 feet.), 73 (note on species), 100 (species), 106 (072306 - genus,

listing of species), **118** (recorded as *Peromyscus merriami merriami* Mearns - Distribution: Known from mesquite bosque situations in southern Arizona. Figure 68, Page 174)*

***Reithrodontomys megalotis* (Baird): Western Harvest Mouse**

COMMON NAME: Western Harvest Mouse. HABITS: Feeds on arachnids, grasses, insects (larvae and adults) and seeds of grasses, forbs and shrubs. Spherical nests are made of woven plant material and lined with plant fibers and can be located near the ground or above the ground in dense vegetation. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *arizonensis*; *aztecus* J.A. Allen, and *megalotis* (Baird)), **55** (recorded as *Reithrodontomys megalotis* (Baird). Western Harvest Mouse. Statewide (120 - 8,000 feet.), 73, 100 (color photograph), 106 (053006), 118 (recorded as *Reithrodontomys megalotis arizonensis* (Allen) - Distribution: Known only from the region of the type locality (Chiricahua Mountains); *Reithrodontomys megalotis aztecus* (Allen) - Distribution: Extreme northeastern part of state, and *Reithrodontomys megalotis megalotis* (Baird) - Distribution: At medium and low elevations statewide except extreme northeastern part of the state. Figure 64, Page 164)*

***Reithrodontomys megalotis* subsp. *megalotis* (Baird): Western Harvest Mouse**

COMMON NAME: Western Harvest Mouse. HABITS: The species feeds on arachnids, grasses, insects (larvae and adults) and seeds of grasses forbs and shrubs. Spherical nests are made of woven plant material and lined with plant fibers and can be located near the ground or above the ground in dense vegetation. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *megalotis* (Baird)), **55** (species: recorded as *Reithrodontomys megalotis* (Baird). Western Harvest Mouse. Statewide (120 - 8,000 feet.), 73 (species), 100 (species, color photograph of species), 106 (053006), **118** (recorded as *Reithrodontomys megalotis megalotis* (Baird) - Distribution: At medium and low elevations statewide except extreme northeastern part of the state. Figure 64, Page 164)*

Mustelidae: The Weasel and Allies Family

***Taxidea taxus* (J.C. von Schreber): American Badger**

COMMON NAMES: American Badger, Badger, Badger Tejon (Hispanic). HABITS: Feeds on ground dwelling birds (and eggs), carrion, insects, rodents and snakes. Young are born in dens in underground burrows. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *berlandieri* Schreber), **55** (recorded as *Taxidea taxus* (Schreber). Badger. Statewide (120 - 7,000 feet.), 65, 73, 100 (color photograph), 106 (053006), **118** (recorded as *Taxidea taxus* - Distribution: Statewide. Figure 98, Page 235)*

Phyllostomidae: The Leaf-nosed Bat Family

***Leptonycteris curasoae* subsp. *yerbabuena* (Martinez & Villa-R.): Southern Long-nosed Bat**

SYNONYMY: *Leptonycteris nivalis sanborni* D.F. Hoffmeister, *Leptonycteris sanborni* (Saussure). COMMON NAMES: Lesser Long-nosed Bat, Little Long-nosed Bat, Mexican Long-nosed Bat, Murcielago de Sanborn (Hispanic), Sanborn's Long-nosed Bat, Sanborn's Southern Long-nosed Bat, Southern Long-nosed Bat. HABITS: The species feeds on insects, nectar, pollen and the nectar and soft-bodied fruits of agaves and cacti. Roosts are located in caves, rock crevices, abandoned mines and tunnels. HABITAT: Within the range of this species it has been reported from the woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: Long-nosed bats are pollinators of Agaves, Cardons, Organ Pipe Cacti and Saguaros. *8, 14 (050907 - **Populations may be compromised by**

roost-site disturbance, loss of food sources and direct killing by humans.), 35 (This species is vulnerable to disturbances at roosting sites by cave explorers.), 55 (species: recorded as *Leptonycteris nivalis* (Saussure). Long-nosed Bat. Locally common in moist caves. Known from Pinal, Pima, Santa Cruz and Cochise Counties.), 92 (recorded as *Leptonycteris sanborni*), 100 (species, recorded as *Leptonycteris curasoae* and *Leptonycteris nivalis*, color photographs), 106 (053006), 110 (recorded as *Leptonycteris sanborni*), **118** (recorded as *Leptonycteris nivalis nivalis* (Saussure) - Distribution: Known only from the southeastern part of the state. Figure 9, Page 35)*

Leptonycteris nivalis (see footnote 55 under *Leptonycteris curasoae* subsp. *yerbabuena*)

Leptonycteris nivalis nivalis (see footnote 118 under *Leptonycteris curasoae* subsp. *yerbabuena*)

Leptonycteris nivalis sanborni (see *Leptonycteris curasoae* subsp. *yerbabuena*)

Leptonycteris sanborni (see *Leptonycteris curasoae* subsp. *yerbabuena*)

***Macrotus californicus* S.F. Baird: California Leaf-nosed Bat**

COMMON NAMES: California Leaf-nosed Bat, Leaf-nosed Bat, Leafnose Bat, Waterhouse's Leaf-nosed Bat, Murcielago de California (Hispanic). HABITS: Feeds on beetles, butterflies, caterpillars, cicadas, crickets, dragonflies, grasshoppers, leafhoppers, moths and other insects. Roosts are located in caves and abandoned mine tunnels. HABITAT: Within the range of this species it has been reported from the grassland, desertscrub and wetland ecological formations. *8, 14 (050907 - subspp. *californicus* (Audubon & Bachman) and *stephensi* (Dalquest)), 55 (recorded as *Macrotus californicus* Baird. Leaf-nosed Bat. Locally common in shallow caves, mine tunnels and under bridges. Occurs widely at lower elevations in the western and southern parts of the state.), 73, 92, 100 (color photograph), 106 (053006), **118** (recorded as *Macrotus californicus* Baird - Distribution: Known from lower elevations in the southern and western parts of the state. Figure 7, Page 32)*

Procyonidae: The Raccoon and Allies Family

***Bassariscus astutus* (M.H. Lichenstein): Ringtail**

COMMON NAMES: Band-tailed Cat, Cacomistle, Civet Cat, Coon Cat Gato Minero (Hispanic), Miner's Cat, Ringtail, Ringtail Cat, Ring-tailed Cat. HABITS: Feeds on berries, birds, fruits, carrion, crickets, eggs, insects, lizards, small mammals, snakes and spiders. Nests are made of grass located in dens in underground burrows, caves, cliffs, rocky outcrops, cavities in logs, stumps and trees and man-made structures. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8 (subspp. *arizonensis*; *nevadensis*, and *yumanensis*), 14 (050907 - subspp. *arizonensis* Goldman; *flavus* Rhoads; *nevadensis*, and *yumanensis* Huey), **55** (recorded as *Bassariscus astutus* (Lichenstein). Ringtail. Statewide (120 - 6,500 feet).), 65 (color photograph), 73, 100 (color photograph), 106 (051107), 118 (recorded as *Bassariscus astutus arizonensis* Goldman - Distribution: Statewide except extreme southeastern and southwestern parts; *Bassariscus astutus flavus* Rhoads - Distribution: Extreme southeastern part of the state, and *Bassariscus astutus yumanensis* Huey - Distribution: Southwestern Arizona. Figure 93, Page 227)*

***Bassariscus astutus* subsp. *arizonensis* Goldman: Ringtail**

COMMON NAMES: Band-tailed Cat, Cacomistle, Civet Cat, Coon Cat Gato Minero (Hispanic), Miner's Cat, Ringtail, Ringtail Cat, Ring-tailed Cat. HABITS: The species feeds on berries, birds, fruits, carrion, crickets, eggs, insects, lizards, small mammals, snakes and spiders. Nests are made of grass located in dens in underground burrows, caves, cliffs, rocky outcrops, cavities in logs, stumps and trees and man-made structures. HABITAT: Within the range of this species it has been reported from the

forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (050907), 55 (species: recorded as *Bassariscus astutus* (Lichenstein). Ringtail. Statewide (120 - 6,500 feet.), 65 (species, color photograph of species), 73 (species), 100 (species, color photograph of species), 106 (053106 - species), **118** (recorded as *Bassariscus astutus arizonensis* Goldman - Distribution: Statewide except extreme southeastern and southwestern parts. Figure 93, Page 227)*

***Nasua narica* (C.H. Merriam): White-nosed Coati**

COMMON NAMES: Antoon, Chula, Chulo, Coati (Indian Name), Coatimundi, El Gato Solo (Los Gatos en Familia), Pizote, White-nosed Coati. HABITS: Feeds on the berries of juniper and manzanita, birds, carrion, eggs, fruits, insects (including among others crickets and grasshoppers) and other invertebrates, prickly pear fruit, lizards, small mammals, nuts, snakes, tubers, worms and yucca fruits. Young are born in dens located in caves, crevices in rocks, mines shafts and cavities among tree roots. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (091008), 55 (recorded as *Nasua narica* (Linnaeus). Coati. In woodland situations in the Graham, Chiricahua, Huachuca, Patagonia and Pena Blanca mountains (5,000 to 7,500 feet.), 65, 73, 100 (color photograph), 106 (053106), **118** (recorded as *Nasua narica pallida* Allen - Distribution: Mountains of southern and southeastern part of the state. Figure 95, Page 230)*

***Procyon lotor* subsp. *mexicanus* Baird: Common Raccoon**

COMMON NAMES: Common Raccoon, Mexican Raccoon, Northern Raccoon, Raccoon, Racuno (Hispanic). HABITS: Feeds on annelid worms, berries, birds, nestlings and eggs, carrion, crayfishes, small fishes, frogs, fruits, insects, small mammals, nuts, shellfish, turtles and turtle eggs and vegetables. Nests are made of leaves located in dens in small caves, amongst boulders, rocky crevices in cliffs and cavities in trees. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: Raccoons are never very far from permanent water. *14 (090108 - subsp. *fuscipes* (Mearns); *hirus* (Nelson & Goldman); *mexicanus* (Baird), and *pallidus* (Merriam)), 55 (species: recorded as *Procyon lotor* (Linnaeus). Raccoon. Riparian situations along the Colorado, Little Colorado and Gila River systems and in the grasslands of the southeastern portion of the state (120 - 6,900 feet.), 65 (color photograph), 73 (species), 100 (species, color photograph of species), 106 (053106), **118** (recorded as *Procyon lotor mexicanus* Baird - Distribution: Southeastern Arizona. Figure 94, Page 229)*

Sciuridae: The Squirrel and Allies Family

***Ammospermophilus harrisi* (J.J. Audubon & Bachman): Harris' Antelope Squirrel**

SYNONYMY: *Citellus harrisi* (J.J. Audubon & Bachman). COMMON NAMES: Ardilla de Tierra Harris (Hispanic), Harris Antelope Squirrel, Harris' Antelope Squirrel, Yuma Antelope Squirrel. HABITS: Feeds on fruits, insects, plants and seeds. Dens are located in underground burrows. HABITAT: Within the range of this species it has been reported from the grassland, desertscrub and wetland ecological formations. *14 (082308), 55 (recorded as *Citellus harrisi* (Audubon & Bachman). Harris Antelope Squirrel. Southern and western parts of the state at elevations below 6,500 feet.), 65 (color photograph), 73, 100 (color photograph), 106 (053106 - genus), **118** (recorded as *Citellus harrisi harrisi* (Audubon & Bachman) - Distribution: Southern and western Arizona except for most of Yuma County. *Citellus harrisi saxicola* (Mearns) - Distribution: Southwestern Arizona. Figure 38, Page 85), **WTK** (March 9, 2007)*

Citellus harrisi (see *Ammospermophilus harrisi*)

Citellus harrisi subsp. *harrisi* (see footnote 118 under *Ammospermophilus harrisi*)

Citellus harrisi subsp. *saxicola* (see footnote 118 under *Ammospermophilus harrisi*)

Citellus spilosoma subsp. *canescens* (see *Spermophilus spilosoma* subsp. *canescens*)

Citellus tereticaudus (see *Spermophilus tereticaudus*)

Citellus tereticaudus subsp. *neglectus* (see footnote 118 under *Spermophilus tereticaudus*)

Citellus variegatus (see *Spermophilus variegatus*)

Citellus variegatus subsp. *grammurus* (see *Spermophilus variegatus* subsp. *grammurus*)

***Spermophilus tereticaudus* S.F. Baird: Round-tailed Ground Squirrel**

SYNONYMY: *Citellus tereticaudus* S.F. Baird. COMMON NAME: Round-tailed Ground Squirrel. HABITS: Feeds on buds of burroweed and mesquite, cacti, green vegetation, insects, seeds of creosote bush, mesquite, flowers of ocotillo, paloverde, plantain, and saltbush, observed visiting road kill and taking scavenging Gambel's Quail chicks; nests are made of plant fibers and stems and located in dens in underground burrows. HABITAT: Within the range of this species it has been reported from the desertscrub ecological formation. *14, 55 (recorded as *Citellus tereticaudus* Baird. Round-tailed Ground Squirrel. Lower Sonoran Life-zone of the western part of the state (below 3,200 feet.), 65, 73, 100 (color photograph), 106 (053106 - genus), 118 (recorded as *Citellus tereticaudus neglectus* (Merriam) - Distribution: Lower Sonoran Life Zone of southwestern Arizona. Figure 39, Page 90)*

***Spermophilus variegatus* (Erxleben): Rock Squirrel**

SYNONYMY: *Citellus variegatus* (Erxleben). COMMON NAMES: Ardilla Coluda (Hispanic), Rock Squirrel. HABITS: Feeds on acorns, berries, small birds, chicks and eggs, carrion, insects, fruits, small mammals, nuts and seeds. Nests are made of leaves, pine needles and plant fibers and located in dens in underground burrows between boulders, rock crevices and talus. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (recorded as *Citellus variegatus* (Erxleben). Rock Squirrel. Statewide, especially at elevations below 6,000 feet.), 65 (color photograph), 73, 100 (color photograph), 106 (053106 - genus), 118 (recorded as *Citellus variegatus grammurus* (Say) - Distribution: Statewide, especially common below 6000 feet. Figure 37, Page 82)*

***Spermophilus variegatus* subsp. *grammurus* (Erxleben): Rock Squirrel**

SYNONYMY: *Citellus variegatus* subsp. *grammurus* (Say). COMMON NAMES: Ardilla Coluda (Hispanic), Rock Squirrel. HABITS: The species feeds on acorns, berries, small birds, chicks and eggs, carrion, insects, fruits, small mammals, nuts and seeds burrows. Nests are made of leaves, pine needles and plant fibers and located in dens in underground burrows between boulders, rock crevices and talus. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (species: recorded as *Citellus variegatus* (Erxleben). Rock Squirrel. Statewide, especially at elevations below 6,000 feet.), 65 (species, color photograph of species), 73 (species), 100 (species, color photograph of species), 106 (053106 - genus), 118 (recorded as *Citellus variegatus grammurus* (Say) - Distribution: Statewide, especially common below 6000 feet. Figure 37, Page 82)*

Soricidae: The Shrew Family

***Notiosorex crawfordi* (E. Coues): Crawford's Desert Shrew**

COMMON NAMES: Crawford's Desert Shrew, Crawford's Gray Shrew, Desert Shrew, Gray Shrew, Musarana del Deseirto Crawford (Hispanic). HABITS: Feeds on centipedes, insects, lizards, small mice, scorpions, sowbugs and spiders. Nests are made of shredded bark and leaves and located in packrat dens or under dead agaves. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *crawfordi* (Coues)), 55 (recorded as *Notiosorex crawfordi* (Coues). Desert Shrew. Locally common, widely distributed statewide at elevations below 6,000 feet, especially in riparian situations.), 65, 73, 100 (color photograph), 106 (051107), 118 (recorded as *Notiosorex crawfordi crawfordi* (Coues) - Distribution: Probably occurs statewide at elevations below 6000 feet. Figure 5, Page 30)*

***Notiosorex crawfordi* subsp. *crawfordi* (E. Coues): Crawford's Desert Shrew**

COMMON NAMES: Crawford's Desert Shrew, Crawford's Gray Shrew, Desert Shrew, Gray Shrew, Musarana del Deseirto Crawford (Hispanic). HABITS: The species feeds on centipedes, insects, lizards, small mice, scorpions, sowbugs and spiders. Nests are made of shredded bark and leaves and located in packrat dens or under dead agaves. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *crawfordi* (Coues)), 55 (species: recorded as *Notiosorex crawfordi* (Coues). Desert Shrew. Locally common, widely distributed statewide at elevations below 6,000 feet, especially in riparian situations.), 65 (species), 73 (species), 100 (species, color photograph of species), 106 (051107 - species), 118 (recorded as *Notiosorex crawfordi crawfordi* (Coues) - Distribution: Probably occurs statewide at elevations below 6000 feet. Figure 5, Page 30)*

Tayassuidae: The Javelina Family

Dicotyles tajacu subsp. *sonoriensis* (see *Peccari tajacu* subsp. *sonoriensis*)

Pecari angulatus (see footnote 65 under *Peccari tajacu* subsp. *sonoriensis*)

***Peccari tajacu* subsp. *sonoriensis* (Mearns): Collared Peccary**

SYNONYMY: *Dicotyles tajacu* subsp. *sonoriensis* (Mearns), *Tayassu tajacu* subsp. *sonoriensis* (Mearns). COMMON NAMES: Collared Peccary, Jabalina (Hispanic), Javelina, "Musk Hog", Peccary. HABITS: The species feeds on agaves, amphibians, berries, bulbs, fruits, fungi, grasses, insects, mesquite beans, nuts, roots, palm nuts, succulent plants, prickly-pear and other cacti, reptiles, rodents, roots, sotol, tubers and worms. Javelina bed down during the day in thick brush and prickly-pear thickets and at night in burrows usually under the roots of trees. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *sonoriensis* (Mearns)), 55 (species: recorded as *Tayassu tajacu* (Linnaeus). Javelina. Southeastern and central parts of the state (1,200 - 6,000 feet.), 65 (species, recorded as *Pecari angulatus*), 73 (species, recorded as *Dicotyles tajacu*), 100 (species, recorded as *Tayassu tajacu*, color photograph of species), 106 (051107 - species, recorded as *Tayassu tajacu*), 118 (recorded as *Tayassu tajacu sonoriensis* (Mearns) - Distribution: Southern part of the state. Figure 107, Page 249)*

Tayassu tajacu subsp. *sonoriensis* (see see *Peccari tajacu* subsp. *sonoriensis*)

Ursidae: The Bear Family

Euarctos americanus subsp. *amblyceps* (see *Ursus americanus* subsp. *amblyceps*)

***Ursus americanus* subsp. *amblyceps* (Baird): Black Bear**

SYNONYMY: *Euarctos americanus* subsp. *amblyceps* (Baird). COMMON NAMES: American Black Bear, Black Bear, Cinnamon Bear, Oso Negro (Hispanic). HABITS: The species feeds on acorns, ants, beetles, berries, buds, carrion, crickets, currants, fish, fruits, grapes, grubs, insects, leaves, pinyon nuts, prickly-pear fruit, raspberries, sprouts, small to medium-size mammals and other vertebrates and twigs. Shelter is taken in dense cover and they climb trees to escape danger. Nests are made of grasses leaves, mud and sticks located in a den. HABITAT: Within the range of this species it has been reported from the tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (050907 - subsp. *amblyceps* (Baird)), 55 (species: recorded as *Euarctos americanus* (Pallas). Black Bear. Formerly common throughout the mountainous areas of the state, now greatly reduced in numbers and distribution.), 73 (species), 100 (species, color photograph of species), 106 (050907 - includes a listing of subspecies and their distribution), 118 (recorded as *Euarctos americanus amblyceps* (Baird) - Distribution: Probably formerly occurred throughout the state, at least in mountainous areas. Figure 91, Page 224)*

***Ursus arctos* subsp. *horribilus* Ord: Grizzly Bear**

SYNONYMY: *Ursus horribilus* Ord. COMMON NAMES: Apache Grizzly, Arizona Grizzly, Grizzly Bear, Navajo Grizzly, New Mexico Grizzly, Oso Gris (Hispanic), Silvertip Bear, Sonora Grizzly, Texas Grizzly. HABITS: The species feeds on berries, carrion, fish (bass, salmon, trout), fungi, grasses, insects (Army Cutworm moths), leaves, large mammals (Bison, Black Bear, Caribou, Deer, Elk, Moose, Mountain Goats) and small mammals (rodents), nuts (Whitebark Pine nuts), roots and sprouts. The Grizzly Bear beds down in depressions in thickets. Dens are excavated from under rocks or located in caves, crevices or hollow trees. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTES: The last confirmed "kill" in Arizona was made on the slopes of Mount Baldy (Apache County) in the summer of 1939. Grizzly Bears were killed-off by American immigrants because of the risks posed to humans and livestock. The Grizzly Bear has been EXTIRPATED from Arizona. *14 (050907 - *Ursus arctos* subsp. *horriaeus* (Baird) and *perturbans* (Merriam)), 39 (recorded as *Ursus horribilus* - included the following note when referring to Grizzly Bears in the Tucson Area "Jack O'Connor told us of a kill in the Catalinas in 1915. Up until 1912, there were quite a few grizzly bears in the Catalinas and also the Galiuros. The Santa Cruz River bottom was a favorite hangout of these bears, all the way from Nogales to the Tucson area. We have a few authentic reports of desert grizzlies, but Jack talked with some old timers who hunted them in the river bottom." The following dates of last known "kills" were provided: Arizona on September 13, 1935 (however, there was a possible sighting in 1936); California in August 1922; New Mexico has two "last" kills one in the spring of 1923 and the other in 1933; Texas on November 2, 1890, and Utah on August 22, 1923. A grizzly bear was killed in the Sierra del Pinitos in Sonora Mexico, a few miles southeast of Nogales, Arizona, on June 18, 1955. This booklet included the listing of six subspecies taken in Arizona: *Ursus horribilus apache*, the Apache Grizzly; *Ursus horribilus arizonae-merriam*, the Arizona Grizzly; *Ursus horribilus baird*, the New Mexico Grizzly; *Ursus horribilus kennerlyi*, the Sonora Grizzly; *Ursus horribilus navajo*, the Navajo Grizzly, and *Ursus horribilus texensis*, the Texas Grizzly), 40 (recorded as *Ursus arctos* - Grizzly Bears were historically present in the Rincon and Santa Catalina Mountains and along the Santa Cruz River bottom from Nogales to Tucson), 55 (recorded as *Ursus horribilus* Ord. Grizzly Bear. Formerly throughout the mountainous areas of the state, now extinct in Arizona.), 73 (recorded as *Ursus horribilus*), 100 (species: recorded as *Ursus arctos*, color photograph), 106 (051207 - *Ursus arctos* subsp. *horribilus* Ord), 118 (recorded as *Ursus horribilus* - Distribution: Formerly statewide, now extinct in Arizona. Figure 92, Page 225)*

Ursus arctos (see footnotes 14 and 100 under *Ursus arctos* subsp. *horribilus*)

Ursus horribilus (see *Ursus arctos* subsp. *horribilus*)

Ursus horribilus apache (see footnote 39 under *Ursus arctos* subsp. *horribilus*)

Ursus horribilus arizonae-merriam (see footnote 39 under *Ursus arctos* subsp. *horribilus*)

Ursus horribilus baird (see footnote 39 under *Ursus arctos* subsp. *horribilus*)

Ursus horribilus kennerlyi (see footnote 39 under *Ursus arctos* subsp. *horribilus*)

Ursus horribilus navajo (see footnote 39 under *Ursus arctos* subsp. *horribilus*)

Ursus horribilus texensis (see footnote 39 under *Ursus arctos* subsp. *horribilus*)

Vespertilionidae: The Plain-nosed Bat Family

***Antrozous pallidus* (J.L. Le Conte): Pallid Bat**

COMMON NAMES: Murcielago Palid (Hispanic), Pallid Bat. HABITS: Feeds on flightless arthropods on the ground, insects, lizards and nectar. Roosts under bridges, buildings, in caves, crevices in cliffs, rocky outcrops, under slabs of rocks, hollow trees and tunnels. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14, **55** (recorded as *Antrozous pallidus* (Le Conte). Pallid Bat. Locally common throughout the state.), 73, 92 (color photograph), 100 (color photograph), 106 (053106), 118 (recorded as *Antrozous pallidus pallidus* (Le Conte) - Distribution: Statewide. Figure 25, Page 60)*

***Antrozous pallidus* subsp. *pallidus* (J.L. Le Conte): Pallid Bat**

COMMON NAMES: Murcielago Pallid (Hispanic), Pallid Bat. HABITS: The species feeds on flightless arthropods on the ground, insects, lizards and nectar. Roosts under bridges, buildings, in caves, crevices in cliffs, rocky outcrops, under slabs of rocks, hollow trees and tunnels. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14, 55 (species: recorded as *Antrozous pallidus* (Le Conte). Pallid Bat. Locally common throughout the state.), 73 (species), 92 (species, color photograph of species), 100 (species, color photograph of species), 106 (053106 - species), **118** (recorded as *Antrozous pallidus pallidus* (Le Conte) - Distribution: Statewide. Figure 25, Page 60)*

Corynorhinus townsendii (see *Plecotus townsendii*)

Corynorhinus townsendii subsp. *pallescens* (see *Plecotus townsendii* subsp. *pallescens*)

***Eptesicus fuscus* (Palisot de Beauvois): Big Brown Bat**

COMMON NAMES: Big Brown Bat, Murcielago Cafe' Grande (Hispanic). HABITS: The species feeds on insects. Roosts under bridges, in buildings, caves, crevices in cliff faces, mines and holes in saguaros and trees. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, **55** (recorded as *Eptesicus fuscus* (Palisot de Beauvois). Big Brown Bat. Locally common throughout the state.), 73, 92 (color photograph), 100 (color photograph), 106 (053106), 118 (recorded as *Eptesicus fuscus pallidus* (Young) - Distribution: Statewide. Figure 20, Page 52)*

***Eptesicus fuscus* subsp. *pallidus* (Young): Big Brown Bat**

COMMON NAMES: Big Brown Bat, Murcielago Cafe' Grande (Hispanic). HABITS: The species feeds on insects. Roosts under bridges, in buildings, caves, crevices in cliff faces, mines and holes in saguaros and trees. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14, 55 (species: recorded as

Eptesicus fuscus (Palisot de Beauvois). Big Brown Bat. Locally common throughout the state.), 73 (species), 92 (species, color photograph of species), 100 (species, color photograph of species), 106 (053106 - species), **118** (recorded as *Eptesicus fuscus pallidus* (Young) - Distribution: Statewide. Figure 20, Page 52)*

***Euderma maculatum* (J.A. Allen): Spotted Bat**

COMMON NAMES: Death's Head Bat, Jackass Bat, Murcielago Pinto (Hispanic), Pinto Bat, Spotted Bat. HABITS: Feeds on insects. Roosts in cracks and crevices in caves, cliffs and ledges, and under loose rock in rocky situations, possibly in close proximity to water. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: This bat is rarely encountered. Riparian habitats seem to be important. *8, 14 (053007), 55 (recorded as *Euderma maculata* (J.A. Allen). Spotted Bat. Extremely rare; known from four specimens, Maricopa and Yuma counties.), 73, 92, 100 (color photograph), 106 (072306), **118** (recorded as *Euderma maculata* (J.A. Allen) - Distribution: Can be expected almost anywhere in the state although recorded from only four localities. Figure 23, Page 57)*

***Lasionycteris noctivagans* (J.L. Le Conte): Silver-haired Bat**

COMMON NAMES: Murcielago Plateado (Hispanic), Silver-haired Bat. HABITS: Feeds on caddis flies, flies, moths and other insects. Uncommon tree dwelling bat found under bark, in bird nests, dead trees, fissures in rock ledges, tree hollows, and woodpecker holes. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14, 55 (recorded as *Lasionycteris noctivagans* (Le Conte). Silver-haired Bat. Uncommon solitary tree-dwelling bat found throughout the state at elevations above 5,000 feet), 73, 92 (color photograph), 100 (color photograph), 106 (053106 - family), **118** (recorded as *Lasionycteris noctivagans* (Le Conte) - Distribution: Probably statewide, at least during certain seasons of the year. Figure 18, Page 48)*

***Lasiurus cinereus* (Palisot de Beauvois): Hoary Bat**

COMMON NAMES: Hoary Bat, Murcielago (Hispanic). HABITS: Feeds primarily on moths. Roosts in buildings, caves, mines, in dense foliage in shrubs and trees and under leaves on the ground. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (091308 - *Lasiurus cinereus cinereus* (Palisot de Beauvois)), **55** (recorded as *Lasiurus cinereus* (Palisot de Beauvois). Hoary Bat. Uncommon tree dwelling bat found throughout the state in the region of trees.), 73, 92 (color photograph), 100 (color photograph), 106 (genus - 053106), **118** (recorded as *Lasiurus cinereus cinereus* (Beauvois) - Distribution: Statewide. Figure 22, Page 55)*

***Lasiurus cinereus subsp. cinereus* (Palisot de Beauvois): Hoary Bat**

COMMON NAMES: Hoary Bat, Murcielago (Hispanic). HABITS: Feeds primarily on moths. Roosts in buildings; caves; mines; in dense foliage in shrubs and trees, and under leaves on the ground. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (091308 - *Lasiurus cinereus cinereus* (Palisot de Beauvois)), 55 (species: recorded as *Lasiurus cinereus* (Palisot de Beauvois). Hoary Bat. Uncommon tree dwelling bat found throughout the state in the region of trees.), 73, 92 (species, color photograph of species), 100 (species, color photograph of species), 106 (053106 - genus), **118** (recorded as *Lasiurus cinereus cinereus* (Beauvois) - Distribution: Statewide. Figure 22, Page 55)*

***Myotis californicus* (J.J. Audubon & Bachman): California Myotis Bat**

COMMON NAMES: California Bat, California Myotis, California Myotis Bat, Murcielago de California (Hispanic). HABITS: Feeds on arachnids and insects. Roosts in crevices and cracks in cliffs and canyon walls, caves, mine shafts and manmade shelters. HABITAT: Within the range of this species

it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8 (*Myotis californicus* N. Miller), 14 (051007 - subsp. *californicus* (Audubon & Bachman) and *stephensi* (Dalquest)), 55 (recorded as *Myotis californicus* Audubon & Bachman. California Myotis. Locally common throughout the state.), 73, 100 (color photograph), 106 (053106 - genus), 118 (recorded as *Myotis californicus californicus* (Audubon & Bachman) - Distribution: Eastern and southeastern Arizona, and *Myotis californicus stephensi* Dalquest - Distribution: Northern and western part of the state. Figure 16, Page 45)*

***Myotis californicus* (J.J. Audubon & Bachman) subsp. *stephensi*: California Myotis Bat**

COMMON NAMES: California Bat, California Myotis, California Myotis Bat, Murcielago de California (Hispanic). HABITS: The species feeds on arachnids and insects. Roosts in crevices and cracks in cliffs and canyon walls, caves, mine shafts and manmade shelters. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8 (species: recorded as *Myotis californicus* N. Miller), 14 (051007 - subsp. *californicus* (Audubon & Bachman) and *stephensi* (Dalquest)), 55 (species: recorded as *Myotis californicus* Audubon & Bachman. California Myotis. Locally common throughout the state.), 73 (species), 100 (species), 106 (genus - 053106), 118 (recorded as *Myotis californicus stephensi* Dalquest - Distribution: Northern and western part of the state. Figure 16, Page 45)*

***Myotis velifer* subsp. *brevis* Vaughan: Cave Myotis Bat**

COMMON NAMES: Cave Bat, Cave Myotis, Cave Myotis Bat, Mexican Brown Bat, Murcielago de Cueva (Hispanic), Southwestern Cave Myotis. HABITS: The species feeds on small moths and other small insects. Roosts in holes and pockets in caves, crevices, bridges, buildings, abandoned mine shafts, tunnels, and trees. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *14 (051007), 55 (species: recorded as *Myotis velifer* (J.A. Allen). Cave Myotis. Locally abundant in summer months at lower elevations (below 5,000 feet) throughout the southern and western parts of the state.), 73 (species), 92 (species), 100 (species, color photograph of species), 106 (053106 - genus), 118 (recorded as *Myotis velifer brevis* Vaughan - Distribution: Probably statewide. Figure 11, Page 37)*

***Myotis yumanensis* (H. Allen): Yuma Myotis Bat**

COMMON NAMES: Murcielago de Yuma (Hispanic), Yuma Myotis, Yuma Myotis Bat. HABITS: Feeds on small insects. Roosts in caves, crevices and swallow nests in cliffs and rocky walls, tree cavities, under bridges and in buildings in close proximity to water. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (072306 - subsp. *yumanensis*), 55 (recorded as *Myotis yumanensis* (H. Allen). Yuma Myotis. Locally common, statewide in distribution.), 73, 100 (color photograph), 106 (072306 - genus with a listing of species), 118 (recorded as *Myotis yumanensis yumanensis* (H. Allen) - Distribution: Probably statewide at low and medium elevation. Figure 10, Page 36)*

***Myotis yumanensis* subsp. *yumanensis* (H. Allen): Yuma Myotis Bat**

COMMON NAMES: Murcielago de Yuma (Hispanic), Yuma Myotis, Yuma Myotis Bat. HABITS: Feeds on small insects. Roosts in caves, crevices and swallow nests in cliffs and rocky walls, tree cavities, under bridges and in buildings in close proximity to water. HABITAT: Within the range of this species it has been reported from the forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8, 14 (072306 - subsp. *yumanensis*), 55 (species: recorded as *Myotis yumanensis* (H. Allen). Yuma Myotis. Locally common, statewide in distribution.), 73 (species), 100 (species, color photograph of species), 106 (072306 - genus, listing of species), 118 (recorded as *Myotis yumanensis yumanensis* (H. Allen) - Distribution: Probably statewide at low and medium elevation. Figure 10, Page 36)*

***Pipistrellus hesperus* (H. Allen): Western Pipistrelle Bat**

COMMON NAMES: Canyon Bat, Flittermouse, Murcielago del Poniente (Hispanic), Western Pipistrelle, Western Pipistrelle Bat. HABITS: Feeds on insects. Roosts in buildings, crevices in canyon walls, caves, cliffs, rocky outcrops, under rocks and in mine shafts. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8 (recorded as *Pipistrellus hesperus* N. Miller), 14 (051007 - subsp. *hesperus* (H. Allen) and *maximus* (Hatfield)), 55 (recorded as *Pipistrellus hesperus* (H. Allen). Western Pipistrelle. Common throughout the state.), 73, 100 (color photograph), 106 (053106 - genus), 118 (recorded as *Pipistrellus hesperus apus* Elliot - Distribution: Southeastern Arizona, and *Pipistrellus hesperus hesperus* (H. Allen) - Distribution: Northern and western Arizona. Figure 19, Page 49)*

***Pipistrellus hesperus* (H. Allen) subsp. *apus* Elliot: Western Pipistrelle Bat**

COMMON NAMES: Canyon Bat, Flittermouse, Murcielago del Poniente (Hispanic), Western Pipistrelle, Western Pipistrelle Bat. HABITS: Feeds on insects. Roosts in buildings, crevices in canyon walls, caves, cliffs, rocky outcrops, under rocks and in mine shafts. HABITAT: Within the range of this species it has been reported from forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. *8 (species: recorded as *Pipistrellus hesperus* N. Miller), 14 (051007), 55 (species: recorded as *Pipistrellus hesperus* (H. Allen). Western Pipistrelle. Common throughout the state.), 73 (species), 100 (species, color photograph of species), 106 (053106 - genus), 118 (recorded as *Pipistrellus hesperus apus* Elliot - Distribution: Southeastern Arizona. Figure 19, Page 49)*

***Plecotus townsendii* (Cooper) (subsp. *pallescens* is the subspecies reported as occurring in Arizona): Pale Townsend's Big-eared Bat**

SYNONYMY: (for *P.t.* subsp. *pallescens*: *Corynorhinus townsendii* subsp. *pallescens* (Frost)). COMMON NAMES: Lump-nosed Bat, Mule-eared Bat, Murcielago de Townsend (Hispanic), Pale Townsend's Big-eared Bat, Western Big-eared Bat, Western Long-eared Bat, Western Lump-nosed Bat. HABITS: The species feeds on small moths and other small insects; roosts on open ceilings in caves and rock shelters, and under bridges and in water diversion tunnels, abandoned mines, mine tunnels and buildings. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: **The Pale Townsend's Big-eared Bat is a rather sedentary species that is extremely sensitive to human disturbance and the vandalism of roost caves.** *14 (053007), 55 (recorded as *Plecotus townsendii* (Cooper). Lump-nosed Bat. Locally common throughout the state at elevations above 5,000 feet; rare at lower elevations.), 73, 92 (color photograph), 100 (color photograph), 118 (recorded as *Corynorhinus townsendii pallescens* Miller - Distribution: Probably more or less state wide but more abundant in the Upper Sonoran and Transitional Life Zones. Figure 24, Page 58)*

***Plecotus townsendii* subsp. *pallescens* (Miller): Pale Townsend's Big-eared Bat**

SYNONYMY: *Corynorhinus townsendii* subsp. *pallescens* (Frost). COMMON NAMES: Lump-nosed Bat, Mule-eared Bat, Murcielago de Townsend (Hispanic), Pale Townsend's Big-eared Bat, Western Big-eared Bat, Western Long-eared Bat, Western Lump-nosed Bat. HABITS: The species feeds on small moths and other small insects; roosts on open ceilings in caves and rock shelters, and under bridges and in water diversion tunnels, abandoned mines, mine tunnels and buildings. HABITAT: Within the range of this species it has been reported from tundra, forest, woodland, scrub, grassland, desertscrub and wetland ecological formations. NOTE: **The Pale Townsend's Big-eared Bat is a rather sedentary species that is extremely sensitive to human disturbance and the vandalism of roost caves.** *14 (053007), 55 (species: recorded as *Plecotus townsendii* (Cooper). Lump-nosed Bat. Locally common throughout the state at elevations above 5,000 feet; rare at lower elevations.), 73 (species), 92 (species, color photograph of species), 100 (species, color photograph of species), 118 (recorded as *Corynorhinus townsendii pallescens* Miller - Distribution: Probably more or less state wide but more abundant in the Upper Sonoran and Transitional Life Zones. Figure 24, Page 58)*

ACKNOWLEDGEMENTS

I would like to thank Matthew B. Johnson for his review of several of the listings, his input into the layout, his numerous trips into the field to assist in the identification of species and above all for his continued support for this project. I would also like to thank Philip D. Jenkins, Assistant Curator, and the Botanists of the University of Arizona Herbarium for years of assistance with plant identifications. I would also like to thank Neva Connolly and Bill Singleton with the Pima County Department of Transportation and Flood Control District for being willing and able to present the listings on the Sonoran Desert Conservation Plan website. Extensive revisions made to the individual species records were made possible by the Southwest Environmental Information Network (SEINet) and the National Plants Database: USDA, NRCS.

FOOTNOTES and REFERENCES for the Species Distribution Listings compiled for Arizona

(1) General Mapping:

Arizona Atlas & Gazetteer. 2002. DeLorme.
www.delorme.com

National Geographic Arizona Seamless USGS Topographic Maps. Maps created with TOPO! R C 2002
National Geographic.

Cocoraque Butte, Arizona - 7.5 Minute Series Topographic 1979

Cocoraque Butte, Arizona - 15 Minute Series Topographic 1941

La Tortuga Butte - 7.5 Minute Series Topographic 1979

Tucson Metropolitan Street Atlas 2005 Edition. Wide World of Maps, Inc., Phoenix, Arizona.
www.maps4u.com

(2) Physiographic Province Mapping:

Walker, Henry P. and Don Bufkin. 1979. Historical Atlas of Arizona, University of Oklahoma Press, Norman, Page 4A and Map.

(3) Soils Mapping:

Arizona General Soil Map, July 1975, United States Department of Agriculture, Soil Conservation Service and the University of Arizona Agricultural Experiment Station, compiled by J.E. Jay, Y.H. Havens, D.M. Hendricks, D.F. Post and C.W. Guernsey.

Richardson, M.L. and M.L. Miller. March 1974. United States Department of Agriculture - Soil Conservation Service in cooperation with the Pima County Natural Resource Conservation District, Report and Interpretations for the General Soil Map of Pima County, Arizona and General Soil Map Pima County Arizona. Arizona General Soil Map, July 1975, United States Department of Agriculture - Soil Conservation Service and the University of Arizona Agricultural Experiment Station, compiled by J.E. Jay, Y.H. Havens, D.M. Hendricks, D.F. Post and C.W. Guernsey.

(4) Biotic Communities Mapping and Definitions

Ecological formations used in the listings follow those presented in the mapping for the Biotic Communities of the Southwest.

Brown, David E. 1982. Biotic Communities of the American Southwest – United States and Mexico, Desert Plants, Volume 4, Numbers 1-4, Published by the University of Arizona for the Boyce Thompson Southwestern Arboretum, Tucson, Arizona.

Brown, David E. and Charles H. Lowe. Revised June 1983. Biotic Communities of the Southwest, August 1980, General Technical Report RM-78, United States Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station.

Brown, David E., Charles H. Lowe and Charles P. Pase. June 1980. A Digitized Systematic Classification for Ecosystems with an Illustrated Summary of the Natural Vegetation of North America, United States Department of Agriculture, Forest Service, General Technical Report RM-73

(5) Nomenclature:

for Plants:

Generally follows that presented by The Biota of North America Program of the North Carolina Botanical Garden (BONAP) with A Synonymized Checklist of the Vascular Flora of the United States, Puerto Rico and the Virgin Islands, Full Index 1998.

<http://www.bonap.org/>

<http://www.csd.tamu.edu/FLORA/b98/check98.htm>

The International Plant Names Index (2004, 2005)

Published on the Internet:

<http://www.ipni.org> [accessed 2004, 2005, 2006]

<http://plants.usda.gov>. National Plant Data Center, Baton Rouge, LA 70874-4490 USA

for Vertebrate Animals:

Section on Arizona Habitats, The University of Arizona Press, Tucson, Arizona and E. Lendell Cockrum. 1960. The Recent Mammals of Arizona: Their Taxonomy and Distribution, The University of Arizona Press, Tucson, Arizona.

Biota Information System of New Mexico (BISON-M), New Mexico Game and Fish, New Mexico Natural Heritage Program

<http://nmnhp.unm.edu/bisonm/bisonquery.php>

for Invertebrate Animals:

Arizona Game and Fish Department. Unpublished Abstracts Compiled and Edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ.

http://www.gf.state.az.us/w_c/edits/species_concern.shtml

Biota Information System of New Mexico (BISON-M), New Mexico Game and Fish, New Mexico Natural Heritage Program

<http://nmnhp.unm.edu/bisonm/bisonquery.php>

(6) Growth Habits of Plants:

Generally coincides with that presented by the National Plants Database. USDA, NRCS. 2004. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA

Common names identified in the USDA NRCS database have been printed in bold lettering: A few of the plants were not provided with a common name in the USDA NRCS database and additional resources were used, including:

Arizona Game and Fish Department. Unpublished Abstracts Compiled and Edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. *8*

Sunset Western Garden Book Kathleen N. Brenzel, 2001, Sunset Publishing Corporation, Menlo Park, California. *18*

(7) Arid Zone Trees, A Resource for Landscape Professionals, dedicated to providing quality trees to the Landscape Industries that are appropriate to the Desert Southwest

<http://www.aridzonetrees.com/index.htm>

(8) Arizona Game and Fish Department. Unpublished abstracts compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ.

http://www.gf.state.az.us/w_c/edits/species_concern.shtml

Amphibians: 2002. *Bufo microscaphus*, Arizona Toad; 2005. *Bufo retiformis*, Sonoran Green Toad; 2001. *Eleutherodactylus augusti* subsp. *cactorum*, Western Barking Frog; 2003. *Gastrophryne olivacea*, Great Plains Narrow-mouthed Toad; 2002. *Hyla arenicolor*, Canyon Treefrog; 2003. *Pternohyla fodiens*, Lowland Burrowing Treefrog; 2001. *Rana chiricahuensis*, Chiricahua Leopard Frog, and 2001. *Rana yavapaiensis*, Lowland Leopard Frog.

Arachnids: 2004. *Albiorix anophthalmus*, a cave obligate Pseudoscorpion.

Birds: 2003. *Accipiter gentilis*, American Goshawk; 2003. *Aimophila quinquestriata*, Five-striped Sparrow; 2002. *Aimophila ruficeps* subsp. *rupicola*: Yuma Rufous-crowned Sparrow; 2001. *Ammodramus bairdii*, Baird's Sparrow; 2001. *Ammodramus savannarum* subsp. *ammolequs*, Arizona Grasshopper Sparrow; 2001. *Anthus spragueii*, Sprague's Pipit; 2002. *Aquila chrysaetos*, Golden Eagle; 2000. *Asturina nitida*, Northern Grey Hawk; 2001. *Athene cunicularia* subsp. *hypugaea*, Western Burrowing

Owl; 2001. *Buteo regalis*, Ferruginous Hawk; 2001. *Buteo swainsoni*, Swainson's Hawk; 2005. *Buteogallus anthracinus*, Common Black-hawk; 2003. *Caracara cheriway*, Crested Caracara; 2002. *Ceryle alcyon*, Belted Kingfisher; 2001. *Chloroceryle americana*, Green Kingfisher; 2002. *Coccyzus americanus* subsp. *occidentalis*, Western Yellow-billed Cuckoo; 2001-08-27. *Colinus virginianus* subsp. *ridgwayi*, Masked Bobwhite; 2002. *Dendrocygna autumnalis*, Black-bellied Whistling-duck; 2001. *Dendrocygna bicolor*, Fulvous Whistling-duck; 2002. *Dolichonyx oryzivorus*, Bobolink; 2002. *Egretta thula*, Snowy Egret; 2002. *Elanus leucurus*, White-tailed Kite; 2003. *Empidonax fulvifrons* subsp. *pygmaeus*, Northern Buff-breasted Flycatcher; 2003. *Empidonax hammondi*, Hammond's Flycatcher; 2002. *Empidonax traillii* subsp. *extimus*, Southwestern Willow Flycatcher; 1998. *Falco peregrinus* subsp. *anatum*, American Peregrine Falcon; 2001. *Glaucidium brasilianum* subsp. *cactorum*, Cactus Ferruginous Pigmy-owl; 2002. *Haliaeetus leucocephalus*, Bald Eagle; 2004. *Lanius ludovicianus*, Loggerhead Shrike; 2005. *Otus flammeolus*, Flammulated Owl; 2002. *Pandion haliaetus*, Osprey; 2002. *Plegadis chihi*, White-faced Ibis; 2002. *Polioptila nigriceps*, Black-capped Gnatcatcher; 2001. *Rallus longirostris* P. Boddaert subsp. *yumanensis*, Yuma Clapper Rail; 2002. *Setophaga ruticilla*, American Redstart; 2005. *Strix occidentalis* subsp. *lucida*, Mexican Spotted Owl; 2001. *Trogon elegans*, Elegant Trogon; 2003. *Tyrannus melancholicus*, Tropical Kingbird, and 2002. *Vireo bellii* subsp. *arizonae*, Arizona Bell's Vireo.

Dicots: 2000. *Abutilon parishii*, Pima Indian Mallow; 2004. *Ammoselinum giganteum*, Sand Parsley; 2003. *Amoreuxia gonzalezii*, Saiya; 2003. *Amsonia kearneyana*, Kearney's Blue Star; 2004. *Arenaria aberrans*, Mt. Dellenbaugh Sandwort; 1995. *Aster potosinus*, Lemmon's Aster; 2004. *Berberis harrisoniana*, Kofa Barberry; 2000. *Boerhavia megaptera*, Tucson Mountain Spiderling; 2004. *Bursera fagaroides*, Torch Wood Copal; 2003. *Capsicum annuum* var. *glabriusculum*, Chiltepin; 2005. *Castela emoryi*, Crucifixion Thorn; 2004. *Cirsium mohavense*, Mohave Thistle; 2001. *Cleome multicaulis*, Playa Spider Plant; 2001. *Colubrina californica*, California Snakewood; 2001. *Coryphantha scheeri* var. *robustispina*, Pima Pineapple Cactus; 2005. *Coryphantha scheeri* var. *valida*, Slender Needle Corycactus; 2004. *Croton wigginsii*, Dune Croton; 2005. *Cryptantha ganderi*, Gander's Cryptantha; 2001. *Dalea tentaculoides*, Gentry Indigo Bush; 2005. *Desmanthus covillei*, Coville Bundleflower; 2004. *Echinocactus horizonthalonius* var. *nicholii*, Nichol Turk's Head Cactus; 2005. *Echinocactus polycephalus*, Cotton-top Cactus; 2005. *Echinocereus fasciculatus*, Magenta-flower Hedgehog Cactus; 2003. *Echinocereus triglochidiatus* var. *arizonicus*, Arizona Hedgehog Cactus; 2004. *Echinomastus erectocentrus* var. *acumensis*, Acuna Cactus; 2003. *Echinomastus erectocentrus* var. *erectocentrus*, Needle-spined Pineapple Cactus; 2001. *Erigeron arisolius*, Arid Throne Fleabane; 2003. *Eriogonum capillare*, San Carlos Wild-buckwheat; 2005. *Eriogonum ericifolium* var. *ericifolium*, Heathleaf Wild-buckwheat; 2004. *Euphorbia gracillima*, Mexican Broomspurge; 2005. *Euphorbia platysperma*, Dune Spurge; 2005. *Ferocactus cylindraceus* var. *cylindraceus*, California Barrel Cactus; 2001. *Graptopetalum bartramii*, Bartram Stonecrop; 2000. *Hackelia ursina*, Chihuahuan Stickseed; 2000. *Hedeoma dentata*, Mock-pennyroyal; 2000. *Hermannia pauciflora*, Sparseleaf Hermannia; 2001. *Heterotheca rutteri*, Huachuca Golden Aster; 2005. *Ibervillea tenuisecta*, Texas Globe Berry; 2000. *Ipomoea tenuiloba*, Trumpet Morning-glory; 2003. *Lilaeopsis schaffneriana* var. *recurva*, Huachuca Water Umbel; 2000. *Lupinus huachucanus*, Huachuca Mountain Lupine; 2004. *Mammillaria mainiae*, Counter Clockwise Fishhook Cactus; 2004. *Matelea cordifolia*, Sonoran Milkweed Vine; 2006. *Passiflora arizonica*, Arizona Passionflower; 2003. *Pectis imberbis*, Beardless Chinch Weed; 2005. *Peniocereus striatus*, Dahlia Rooted Cereus; 2004. *Penstemon superbus*, Superb Beardtongue; 2005. *Perityle ajoensis*, Ajo Rock Daisy; 2005. *Petalonyx linearis*, Longleaf Sandpaper-plant; 2004. *Pholisma sonorae*, Sand Food; 2004. *Plagiobothrys pringlei*, Pringle Popcorn-flower; 2005. *Rhus kearneyi*, Kearney Sumac; 2005. *Stenocereus thurberi*, Organ Pipe Cactus; 2005. *Stephanomeria schottii*, Schott Wire Lettuce; 2004. *Stevia lemmonii*, Lemmon's Stevia; 2004. *Tragia laciniata*, Sonoran Noseburn; 2004. *Tumamoca macdougalii*, Tumamoc Globeberry; 2005. *Vauquelinia californica* subsp. *sonorensis*, Sonoran Mountain Rosewood, and 2004. *Viola umbraticola*, Shade Violet.

Ferns: 1997. *Cheilanthes pringlei*, Pringle Lip Fern and 2003. *Notholaena lemmonii*, Lemmon Cloak Fern.

Fishes: 2002. *Agosia chrysogaster*, Longfin Dace; 2002. *Catostomus clarki*, Desert Sucker; 2002. *Catostomus insignis*, Sonora Sucker; 2001. *Cyprinodon eremus*, Quitobaquito Pupfish; 2001. *Cyprinodon macularius*, Desert Pupfish; 2002. *Gila intermedia*, Gila Chub; 2002. *Gila robusta*, Roundtail Chub; 2001. *Poeciliopsis occidentalis* subsp. *occidentalis*, Gila Topminnow, and 2001. *Poeciliopsis occidentalis* subsp. *sonorensis*, Yaqui Topminnow.

Gastropods: 2003. *Tryonia quitobaquिताe*, Quitobaquito Tryonia.

Insects: 2001. *Agathymus aryxna*, Arizona Giant Skipper; 2001. *Agathymus polingi*, Poling's Giant Skipper; 2004. *Anthocharis cethura*, Desert Orangetip; 2001. *Calephelis rawsoni* subsp. *arizonensis*, Arizona Metalmark; 2002. *Heterelmis stephani*, Stephan's Heterelmis Riffle Beetle; 2001. *Limenitis archippus* subsp. *obsoleta*, Obsolete Viceroy Butterfly, and 2001. and *Neophasia terlootii*, Chiricahua Pine White.

Mammals: 2002. *Antrozous pallidus*, Pallid Bat; 2002. *Antilocapra americana* subsp. *mexicana*, Chihuahuan Pronghorn Antelope; 2002. *Antilocapra americana* subsp. *sonoriensis*, Sonoran Pronghorn Antelope; 2004. *Bassariscus astutus*, Ringtail; 2001. *Canis lupus baileyi*, Mexican Gray Wolf; 2003. *Choeronycteris mexicana*, Mexican Long-tongued Bat; 2004. *Eptesicus fuscus*, Big Brown Bat; 2003. *Euderma maculatum*, Spotted Bat; 2002. *Eumops perotis* subsp. *californicus*, Greater Western Bonneted Bat; 2003. *Eumops underwoodi*, Underwood's Mastiff Bat; 2004. *Herpailurus yaguarondi*, Jaguarundi; 2004. *Lasionycteris noctivagans*, Silver-haired Bat; 2003. *Lasiurus blossevillii*, Western Red Bat; 2004. *Lasiurus cinereus*, Hoary Bat; 2004. *Leopardus pardalis* subsp. *sonoriensis*, Ocelot; 2003. *Leptonycteris curasoae* subsp. *yerbabuena*, Lesser Long-nosed Bat; 2002. *Lontra canadensis* subsp. *sonora*, Southwestern River Otter; 2001. *Macrotus californicus*, California Leaf-nosed Bat; 2003. *Myotis auriculus*, Southwestern Myotis; 2004. *Myotis californicus*, California Myotis; 2003. *Myotis ciliolabrum*, Western Small-footed Myotis; 2003. *Myotis occultus*, Fringed Myotis; 2003. *Myotis yumanensis*, Yuma Myotis; 2003. *Nyctinomops femorosacca*, Pocketed Free-tailed Bat; 2003. *Nyctinomops macrotis*, Big Free-tailed Bat; 2003. *Myotis thysanodes*, Fringed Myotis; 2002. *Myotis velifer*, Cave Myotis; 2004. *Panthera onca*, Jaguar; 2004. *Pipistrellus hesperus*, Western Pipistrelle; 2007. *Puma concolor*, Mountain Lion; 2005. *Sciurus arizonensis*, Arizona Gray Squirrel; 2003. *Sigmodon ochrognathus*, Yellow-nosed Cotton Rat, and 2004. *Tadarida brasiliensis*, Brazilian Free-tailed Bat.

Monocots: 2005. *Agave x ajoensis*, Ajo Agave; 2003. *Agave murpheyi*, Hohokam Agave; 1994. *Agave parviflora* subsp. *parviflora*, Santa Cruz Striped Agave; 2005. *Agave schottii* var. *treleasei*, Trelease Agave; 2005. *Agave utahensis* var. *kaibabensis*, Kaibab Agave; 2005. *Allium bigelovii*, Bigelow Onion; 1999. *Allium gooddingii*, Goodding Onion; 2005. *Allium parishii*, Parish Onion; 2004. *Carex chihuahuensis*, Chihuahuan Sedge; 2000. *Carex ultra*, Arizona Giant Sedge; 2004. *Cathestecum erectum*, False Grama; 2004. *Hexalectris revoluta*, Chisos Coral-root; 2005. *Hexalectris spicata*, Crested Coral Root; 2001. *Lilium parryi*, Lemon Lily; 2005. *Listera convallarioides*, Broadleaf Twayblade; 2000. *Muhlenbergia xerophila*, Weeping Muhly, and 2005. *Schiedeella arizonica*, Fallen Ladies'-tresses.

Reptiles: 2001. *Aspidoscelis burti* subsp. *stictogrammus*, Giant Spotted Whiptail; 2003. *Aspidoscelis burti* subsp. *xanthonotus*, Redback Whiptail; 2002. *Chionactis occipitalis* subsp. *klauberi*, Tucson Shovel-nosed Snake; 2003. *Chionactis palarostris* subsp. *organica*, Organ Pipe Shovel-nosed Snake; 2001. *Crotalus lepidus* subsp. *klauberi*, Banded Rock Rattlesnake; 2001. *Gopherus agassizi*, Desert Tortoise; 2002. *Heloderma suspectum* subsp. *cinctum*, Banded Gila Monster; 2002. *Heterodon nasicus* subsp. *kennerlyi*, Mexican Hog-nosed Snake; 2005. *Kinosternon sonoriense*, subsp. *longifemorale*, Sonoyta Mud Turtle; 2003. *Lichanura trivirgata* subsp. *gracia*, Desert Rosy Boa; 2003.

Phrynosoma mcallii, Flat-tailed Horned Lizard; 2005. *Sauromalus ater*, Common Chuckwalla; 2001. *Thamnophis eques* subsp. *megalops*, Mexican Garter Snake; 2003. *Uma rufopunctata*, Yuma Desert Fringe-toed Lizard, and 2003. *Xantusia arizonae*, Arizona Night Lizard.

(9) Arizona Rare Plant Committee. Arizona Rare Plant Field Guide, A Collaboration of Agencies and Organizations.

(10) Arizona Sonora Desert Museum, Migratory Pollinators Program, Spring 2003 Update, Table 3. Plants Visited by Hummingbirds in Sonora
http://desertmuseum.org/pollination/table_3.html

(11) Barnes, Will C. 1988. Arizona Place Names, The University of Arizona Press, Tucson, Arizona.

(12) Benson, Lyman. 1981. The Cacti of Arizona, The University of Arizona Press, Tucson, Arizona.

(13) Benson, Lyman and Robert A. Darrow. 1981. Trees and Shrubs of the Southwestern Deserts, The University of Arizona Press, Tucson, Arizona.

(14) Biota Information System of New Mexico (BISON-M), New Mexico Game and Fish, New Mexico Natural Heritage Program
<http://nrmnhp.unm.edu/bisonm/bisonquery.php>

(15) Bowers, Janice E. and Steven P. McLaughlin. 1987. Flora and Vegetation of the Rincon Mountains, Pima County, Arizona. Desert Plants, Vol. 8, No. 2, pp. 50-95, 1987.

(16) Bowers, J.E., and R.M. Turner. 1985. A Revised Vascular Flora of Tumamoc Hill, Tucson, Arizona. Madrono, Vol.32, No.4, pp. 225-252, 20 December 1985.

(17) Breitung, August J., The Agaves, The Cactus and Succulent Journal 1968 Yearbook, Abbey Garden Press, Reseda, California.

(18) Brenzel, Kathleen N. 2001. Sunset Western Garden Book, Sunset Publishing Corporation, Menlo Park, California.

(19) Brown, David E. 1982. Biotic Communities of the American Southwest – United States and Mexico, Desert Plants, Volume 4, Numbers 1-4, Published by the University of Arizona for the Boyce Thompson Southwestern Arboretum, and associated map: Brown, David E. and Lowe, Charles H., Biotic Communities of the Southwest, August 1980, General Technical Report RM-78, United States Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station Revised June 1983.

(20) Bull, John and John Farrand, Jr. 1977. The Audubon Society Field Guide to North American Birds: Eastern Region, Alfred A. Knopf, Inc., New York, New York.

(21) Catalogue of New World Grasses
<http://mobot.mobot.org/W3T/Search/index/nwgcA.html>

(22) Chambers, Nina – Sonoran Institute & Hawkins, Trica Oshant - Environmental Education Exchange. Invasive Plants of the Sonoran Desert, A Field Guide.

(23) Checklist of North American Butterflies Occurring North of Mexico

<http://www.naba.org/pubs/enames2.html>

- (24) Checklist of Plants, Organ Pipe Cactus National Monument, August 2005.
- (25) Dollar, Derrick; Scott Richardson and Erin Deely. 2000. Mammal Survey for the Mason Audubon Center, Tucson, Arizona USA.
- (26) Duffield, Mary Rose and Warren D. Jones. 1981. Plants for Dry Climates, HP Books, Los Angeles, California.
- (27) Earle, W. Hubert. 1963. Cacti of the Southwest, Rancho Arroyo book distributors, Tempe, Arizona.
- (28) Epple, Anne Orth. 1995. A Field Guide to the Plants of Arizona, Falcon Press Publishing Co., Inc., Helena, Montana.
- (29) Erickson, Jim. 1998. 2 Areas Near Santa Ritas Sought for Conservation, Park, the Arizona Daily Star, Tuesday, 17 November 1998.
- (30) Especies con Usos No Maderables en Bosques de Encino, Pino y Pino-Encino en los Estados de Chihuahua, Durango, Jalisco, Michoacan, Guerrero y Oaxaca.
<http://www.semarnat.gob.mx/pfnm/indices.htm>
- (31) Felger, Richard S. 1997. Checklist of the Vascular Plants of Cabeza Prieta National Wildlife Refuge, Arizona, Drylands Institute, Tucson, Arizona.
- (32) Florida Nature
<http://www.floridanature.org/>
<http://www.floridanature.org/copyright.asp>
- (33) Gould, Frank W. 1951. Grasses of Southwestern United States, University of Arizona Press, Tucson, Arizona.
- (34) Hawksworth, Frank G. and Delbert Wiens. March 1996. United States Department of Agriculture, Forest Service. Agricultural Handbook 709 - Dwarf Mistletoes: Biology, Pathology, and Systematics.
http://www.rmrs.nau.edu/publications/ah_709/index.html
- (35) Haynes, Lisa and Susan Schuetze. 1997. Pamphlet: A Sampler of Arizona's Threatened and Endangered Wildlife, Arizona Game and Fish Department and Arizona Department of Agriculture.
- (36) The Hermannia Pages: American Species
<http://www.meden.demon.co.uk/Malvaceae/Hermannia/American.html>
- (37) Heymann, M.M. 1975. Reptiles and Amphibians of the American Southwest, Doubleshoe Publishers, Scottsdale, Arizona.
- (38) Hodge, Carle. 1991. All About Saguaros, Arizona Highways Magazine, Arizona Department of Transportation, Phoenix, Arizona.
- (39) Hoffmeister. 1980. *Ursus arctos*, Specimens in Collections
- (40) Housholder, Bob. 1966. The Grizzly Bear in Arizona

(41) Howery, Larry D. and Gina Ramos. Arizona's Invasive Weeds, The University of Arizona, Cooperative Extension Service and United States Department of the Interior, Bureau of Land Management.

(42) Integrated Taxonomic Information System (ITIS) on-line database:
<http://www.itis.usda.gov>.

(43) The International Plant Names Index (2004), accessed 2005 and 2005, published on the Internet:
<http://www.ipni.org>

(44) Jepson Flora Project
<http://ucjeps.berkeley.edu/>
<http://ucjeps.berkeley.edu/copyright.html>

(45) Johnson, Matthew Brian. 2004. Cacti, other Succulents, and Unusual Xerophytes of Southern Arizona, Boyce Thompson Southwestern Arboretum / Arizona Lithographers, Tucson, Arizona.

(46) Kearney, Thomas K., Robert H. Peebles and collaborators. 1960. Arizona Flora. Second Edition with Supplement by John Thomas Howell and Elizabeth McClintock and collaborators, 4th printing 1973, University of California Press, Berkeley, Los Angeles, California.

(47) Krausman, Paul R. and Michael L. Morrison. 2003. Wildlife Ecology and Management, Santa Rita Experimental Range (1903 to 2002), USDA Forest Service Proceedings RMRS-P-30.2003 Pages 59 thru 67.

(48) Landscaping with Native Arizona Plants. 1973. Natural Vegetation Committee, Arizona Chapter, Soil Conservation Society of America, The University of Arizona Press, Tucson, Arizona.

(49) Las Cienegas National Conservation Area - Records and Reports.

(50) Laymon, Stephen A. Paper: Yellow-billed Cuckoo.

(51) Lellinger, David B. 1985. A Field Manual of the Ferns and Fern-Allies of the United States and Canada, Smithsonian Institution Press, Washington, D.C.

(52) Little, Elbert L. 1980. The Audubon Society Field Guide to North American Trees – Western Region, Alfred A. Knopf, New York, New York.

(53) Little, Elbert L., Jr. December 1950. Southwestern Trees - A Guide to the Native Species of New Mexico and Arizona, Agriculture Handbook No. 9, United State Department of Agriculture, Forest Service, U.S. Government Printing Office, Washington 25 D.C.

(54) Lowe, Charles H., Cecil R. Schwalbe and Terry B. Johnson. 1986. The Venomous Reptiles of Arizona, Arizona Game and Fish Department, Phoenix, Arizona.

(55) Lowe, Charles H. 1964. The Vertebrates of Arizona with Major Section on Arizona Habitats, The University of Arizona Press, Tucson, Arizona.

(56) Maus, Kathryn. October 12, 2001. Plants of the West Branch of the Santa Cruz River, The West Branch Flora, Arid Lands Resource Sciences, University of Arizona, Tucson, Arizona.

<http://www.co.pima.az.us/cmo/sdcp/sdcp2/reports/WB/pflora.htm>

(57) Maus, Kathryn. September 9, 2002. "Checklist for the Plants of the West Branch of the Santa Cruz, Tucson, Arizona..

<http://eebweb.arizona.edu/HERB/WESTBRANCH/westbranch.html>

(58) McLaughlin, Steven P. July 18, 1990. Flora of Buenos Aires National Wildlife Refuge (including Arivaca Cienega), Office of Arid Land Studies, University of Arizona.

(59) Medina, Alvin L. 2003. Historical and Recent Flora of the Santa Rita Experimental Range, USDA Forest Service Proceedings RMRS-P-30.2003 Pages 141 thru 148.

(60) Milne, Lorus and Margery. 1980. The Audubon Society Field Guide to North American Insects and Spiders, Alfred A. Knopf, New York, New York.

(61) Minckly, W.L. 1973. Fishes of Arizona, Sims Printing Company, Inc., Phoenix, Arizona.

(62) Missouriplants.com

<http://www.missouriplants.com/index.html>

(63) National Plants Database: USDA, NRCS. 2004. The PLANTS Database, Version 3.5, National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

<http://plants.usda.gov>

with links to the following sites:

USDA, ARS, National Genetic Resources Program.

Germplasm Resources Information Network (GRIN) [Online Database].

National Germplasm Resources Laboratory, Beltsville, Maryland.

URL: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?447394> (29 November 2008)

<http://www.ars-grin.gov/npgs/aboutgrin.html>

Flora of North America

www.efloras.org

United State Department of Agriculture Forest Service, Fire Effects Information System

<http://www.fs.fed.us/database/feis/index.html>

Grass Manual on the Web

The Center for Plant Conservation

Kemper Center for Home Gardening

<http://www.mobot.org/gardeninghelp/plantinfo.shtml>

Burke Museum of Natural History and Culture

<http://www.washington.edu/burkemuseum/>

(64) Native Grasses from South Texas, Texas A&M University System, Agricultural Program.

<http://uvalde.tamu.edu/herbarium/grasses.htm>

(65) Olin, George. 1975. Mammals of the Southwest Deserts, Popular Series No. 8, Southwest Parks and Monuments Association.

(66) Owensby, Clenton. 2002. Line Drawings of Kansas Grasses
<http://spuds.agron.ksu.edu/ksgrasskey/linedraw.htm>

(67) Page, Lawrence M. and Brooks M. Burr. 1991. A Field Guide to Freshwater Fishes – North America North of Mexico, Peterson Field Guides, Houghton Mifflin Company, Boston, Massachusetts.

(68) Parker, Kittie F. 1982. An Illustrated Guide to Arizona Weeds, University of Arizona Press, Tucson, Arizona.

(69) Peterson, Roger Tory. 1961. A Field Guide to Western Birds, Houghton Mifflin Company, Boston, Massachusetts.

(70) Pima Community College – Desert Ecology of Tucson, Arizona
http://wc.pima.edu/Bfiero/tucsonecology/plants/wflow_heri.htm

(71) Pima County Parks and Recreation Department, Cienega Creek Natural Preserve Bird Checklist, Tucson, Arizona.

(72) Pima County Sonoran Desert Conservation Plan Threatened and Endangered Species
<http://www.pima.gov/cmo/sdcp/sdcp2/fsheets/facts.html>

(73) Ransom, Jay Ellis. 1981. Harper and Row's Complete Field Guide to North American Wildlife, Western Edition, Harper and Row, New York, New York.

(74) Raven, Peter H., Ray F. Evert and Helena Curtis. 1976 Biology of Plants, Second Edition, Worth Publishers, Inc.

(75) Richardson, M.L. and M.L. Miller. March 1974. United States Department of Agriculture, Soil Conservation Service in cooperation with The Pima County Natural Resource Conservation District, Reports and Interpretations for the General Soil Map of Pima County, Arizona and General Soil Map Pima County Arizona.

(76) Richmond, D.L. and M.L. Richardson. January 1974. United States Department of Agriculture, Soil Conservation Service in cooperation with the Natural Resource Conservation Districts in Mohave County, General Soil and Interpretations, Mohave County, Arizona and General Soil Map Mohave County, Arizona.

(77) Rondeau, Renee, Thomas R. Van Devender, C. David Bertelson, Philip Jenkins, Rebecca K. Wilson, Mark A. Dimmitt. December, 1996. Annotated Flora of the Tucson Mountains, Pima County, Arizona, Desert Plants, Volume 12, Number 2.
<http://eebweb.arizona.edu/herb/TUCSONS/tucsonsa-C.html>

(78) Rosen, Philip C. 15 October 2001. Biological Values of the West Branch of the Santa Cruz River, With an Outline for a Potential River Park or Reserve, Including a Preliminary Flora by Kathryn Maus (Plants of the West Branch of the Santa Cruz , The West Branch Flora has been recorded separately as Footnote 56), School of Renewable Natural Resources, University of Arizona, Tucson, Arizona.
<http://www.co.pima.az.us/cmo/sdcp/sdcp2/reports/WB/WestB.htm>

- (79) Rosenberg, Gary H. and Russel, Ruth. 1999. Checklist of North American Birds United States and Canada Including Hawaii 2000, Tucson Audubon Society.
- (80) Schmutz, Ervin M., Barry N. Freeman, Raymond E. Reed. 1968. Livestock- Poisoning Plants of Arizona, The University of Arizona Press, Tucson, Arizona.
- (81) School of Botanical Medicine - Checklist of the Vascular Plants of Arizona (excluding grasses and their allies)
<http://www.ibiblio.org/london/alternative-healthcare/Southwest-School-of-Botanical-Medicine/HOMEPAGE/Floras/AZchklist.txt>
- (82) Southeast Arizona Butterfly Association (SEABA), Plant List - SEABA's Butterfly Garden at the Tucson Audubon Society's Mason Center
<http://www.naba.org/chapters/nabasa/home.html>
- (83) Southwest Parks and Monument Association. 1991. A Checklist of Mammals, Amphibians and Reptiles of Organ Pipe Cactus National Monument, Tucson, Arizona.
- (84) Southwest Parks and Monument Association. 1999. A Checklist of the Birds of Organ Pipe Cactus National Monument, Tucson, Arizona.
- (85) Southwest Environmental Information Network (SEINet)
<http://seinet.asu.edu/collections/selection.jsp?cat=plantae>
- (86) Spellenberg, Richard. 1979. The Audubon Society Field Guide to North American Wildflowers - Western Region, Alfred A. Knopf, New York, New York.
- (87) Stebbins, Robert C. 1985. A Field Guide to Western Reptiles and Amphibians, Peterson Field Guides, Houghton Mifflin Company, Boston, Massachusetts.
- (88) Texas Native Shrubs
<http://aggie-horticulture.tamu.edu/ornamentals/nativeshrubs/indexscientific.htm>
- (89) Thornber, J.J. Vegetation Groups in the Desert Laboratory Domain *in* Spalding. 1909. The Distribution and Movements of Desert Plants, Carnegie Institution of Washington, Publication No. 113: 103 - 112.
- (90) Tohono Chul Park, Field Checklist of Birds, Tucson, Arizona.
- (91) Turner, Raymond M., Janice E. Bowers and Tony L. Burgess. 1995. Sonoran Desert Plants An Ecological Atlas, The University of Arizona.
- (92) Tuttle, Merlin D. 1988. America's Neighborhood Bats, University of Texas Press, Austin, Texas.
- (93) Udvardy, Miklos D.F. 1977. The Audubon Society Field Guide to North American Birds: Western Region, Alfred A. Knopf, Inc., New York, New York.
- (94) United States Fish and Wildlife Service, Cabeza Prieta National Wildlife Refuge: Listing of Amphibians (April 15, 2002 Update), Listing of Birds (March 2004), Listing of Mammals (April 15, 2002 Update), Listing of Plants (April 15, 2002 Update) and Listing of Reptiles (April 15, 2002 Update).
<http://www.fws.gov/southwest/refuges/arizona/cabeza.html>

(94 ES 1998) United States Department of the Interior, Endangered Species on Cabeza Prieta National Wildlife Refuge (October 1998).

(94 ETCS 1994) United States Department of the Interior, Endangered, Threatened and Candidate Species Cabeza Prieta National Wildlife Refuge (June 1994).

(95) University of Arizona

Herbarium, P.O. Box 210036 Herring Hall, 1130 East South Campus Drive, Tucson, Arizona 85721; 520-621-7243; FAX: 520-621-7186
<http://ag.arizona.edu/herbarium/>

Department of Entomology, Forbes 410, PO Box 2100: (36), Tucson, Arizona 85721-0036; 520-621-1151; FAX: 520-621-1150
<http://ag.arizona.edu/ento/insectid.htm>

(96) University of Michigan, Animal Diversity Web
<http://animaldiversity.ummz.umich.edu/>

(97) Venomous Creatures of the Southwest, Arizona-Sonora Desert Museum and the Arizona Poison Control System. University of Arizona, Poison and Drug Information Center, College of Pharmacy, Tucson 1-800-222-1222, and the Samaritan Regional Poison Center, Good Samaritan Medical Center - Phoenix and the Arizona Department of Health Services - Emergency Medical Services Division.
<http://www.pharmacy.arizona.edu/outreach/poison/>
<http://www.pharmacy.arizona.edu/outreach/poison/venom.php>
<http://www.pharmacy.arizona.edu/outreach/poison/plants.php>

(98) Walker, Henry P. and Don Bufkin. 1979. Historical Atlas of Arizona, University of Oklahoma Press, Norman, Page 4A and Map.

(99) Walters, James W. R3 78-9, A Guide to Forest Diseases of Southwestern Conifers, Forest Insect and Disease Management, State and Private Forestry, Southwestern Region, Forest Service, United States Department of Agriculture, Albuquerque, New Mexico.

(100) Whitaker, John O., Jr. 1996. National Audubon Society Field Guide to North American Mammals, Alfred A. Knopf, New York, New York.

(101) Whitson, Tom D., Larry C. Burrill, Steven A. Dewey, David W. Cudney, B.E. Nelson, Richard D. Lee, Robert Parker. 1996. Weeds of the West, Pioneer of Jackson Hole, Jackson, Wyoming.

(102) Wiens, John F. Vascular Plants of Ragged Top, compiled by John F. Wiens from 1987 - 2000, The Arizona Native Plant Society, The Plant Press, Volume 25 Number 1, Spring 2001.

(103) Wildflowers and Other Plants of Southern California, with Photographs by Michael L. Charters
<http://www.calflora.net/bloomingplants/index.html>

(104) Lehr, J. Harry. 1978. A Catalogue of the Flora of Arizona, Desert Botanical Garden, Phoenix, Arizona. Northland Press, Flagstaff, Arizona.

- (105) Humphrey, Robert H., Albert L. Brown and A.C. Everson. April 1956. Bulletin 243, Common Arizona Range Grasses, Agricultural Experiment Station, University of Arizona, Tucson, Arizona.
- (106) Wikipedia, The Free Encyclopedia
http://en.wikipedia.org/wiki/Main_Page
- (107) McGinnies, William G. 1981. Discovering the Desert, Legacy of the Carnegie Desert Botanical Laboratory, The University of Arizona Press, Tucson, Arizona.
- (108) Dodge, Natt N. 1964. Organ Pipe Cactus National Monument / Arizona, Natural History Handbook Series, No. 6, Washington, D.C.
- (109) Grow Native! Don't Plant a Pest, A Guide to Invasive Landscape Plants and Their Native Alternatives - Southeastern Arizona. Arizona Native Plant Society.
www.aznps.org
- (110) United States fish and Wildlife Service, Ecological Services Field Office, Endangered and Threatened Species of Arizona - Summer 1991.
- (111) California Register of Big Trees
<http://www.ufe.org/BigTrees/index.html>
- (112) Kitt Peak Handouts: Common Trees and Shrubs on Kitt Peak; Common Birds of Kitt Peak; Common Mammals of Kitt Peak, and Common Reptiles and Amphibians of Kitt Peak.
- (113) Halbedel, E. June 2005. The Birds of Kitt Peak, Revised 3rd Edition.
- (114) Nearctica.com, Inc. 1999, The Natural World of North America.
<http://www.nearctica.com/>
<http://www.nearctica.com/nomina/nomina.htm>
- (115) The Firefly Forest
<http://fireflyforest.net/firefly/>
and Wildflowers of Tucson, Arizona
<http://www.fireflyforest.com/flowers/index.html>
- (116) Krausman, Paul R. and Michael L. Morrison, Wildlife Ecology and Management, Santa Rita Experimental Range (1903 to 2002), USDA Forest Service Proceedings RMRS-P-30.2003: 59 - 67.
- (117) Medina, Alvin L., Historical and Recent Flora of the Santa Rita Experimental Range, USDA Forest Service Proceedings RMRS-P-30.2003: 141 - 148.
- (118) Cockrum, E. Lendell. 1960. The Recent Mammals of Arizona: Their Taxonomy and Distribution, The University of Arizona Press, Tucson, Arizona.
- (119) Stockwell, William Palmer and Lucretia Breazaele. April 1, 1933. Arizona Cacti, University of Arizona Bulletin, Vol. 4, No. 3, Biological Science Bulletin No. 1, University of Arizona, Tucson, Arizona.

(120) Duncan, Russell B. Two Rare Plants and the Warm Season Flora of a Unique Habitat in Pima County, Arizona: The Pantano Formation, Claystone Member Deposits, The Arizona Native Plant Society, The Plant Press, Autumn 2003: 7-14.

(121) Reichhardt, Karen. *Triteliopsis palmeri* - Blue Sand Lily, an Elusive Plant of the Sand Dunes, The Arizona Native Plant Society, The Plant Press, Volume 30 Number 2, October 2006: 10-11.

(122) Kaiser, Jack. Common Ferns of Southern Arizona, The Arizona Native Plant Society, The Plant Press, Volume 18 Number 2, Spring 1994: 5-12.

(123) McDonald, Christopher. Pima Pineapple Cactus, The Arizona Native Plant Society, The Plant Press, Volume 31 Number 1, April 2007: 1-4.

(124) Historical Common Names of Great Plains Plants

<http://www.unl.edu/agnicpls/gpcn/index.html>

(125) Munson, T.V. Foundations of American Grape Culture, T.V. Munson & Son, Denison, Texas, 1909.

(126) Adams, Robert P. *Juniperus* of Canada and the United States: Taxonomy, Key and Distribution, Biology Department, Baylor University, Box 727, Gruver, TX 79040 USA, December 2008.

Robert_Adams@baylor.edu

[http://www.juniperus.org/AdamsPapersPDFFiles/218-Phyto90\(3\)255-314AdamsKeytoJuniperusCanadaandUS.pdf](http://www.juniperus.org/AdamsPapersPDFFiles/218-Phyto90(3)255-314AdamsKeytoJuniperusCanadaandUS.pdf)

(127) Native American Ethnobotany, University of Michigan. A database of plants used as drugs, foods, dyes, fibers and more, by native peoples of North America.

<http://herb.umd.umich.edu/>

(128) Desert-Tropicals.com, Philippe Faucon

<http://www.desert-tropicals.com/index.html>

(129) Plants of the Southwest, Santa Fe, New Mexico 87501 U.S.A.

<http://www.plantsofthesouthwest.com/>

(130) Little, V.A. 1963. General and Applied Entomology, Harpers and Row, Publishers, Inc. New York, N.Y.

(131) The University of Arizona, Cooperative Extension, Pima county Home Horticulture.

<http://ag.arizona.edu/pima/gardening/gardening.html>

(132) The Gymnosperm Database

<http://www.conifers.org/index.html>

(132) PIER, Pacific Island Ecosystems at Risk, Plant threats to Pacific ecosystems

<http://www.hear.org/pier/index.html>

(133) USDA, ARS, National Genetic Resources Program. Germplasm Resources Information Network (GRIN) [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland.URL:

<http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?447394> (03 October 2009)

<http://www.ars-grin.gov/npgs/aboutgrin.html>

(134) Austin, Daniel F. Baboquivari Mountain Plants, The Plant Press, The Arizona Native Plant Society, Volume 33, Number 2, Fall 2009: 1-4.

(135) Encyclopedia of Life. Available from <http://www.eol.org>. Accessed 19 December 2009, 25 January 2010

(136) Flora of North America
www.efloras.org

(137) Kleinman, Dr. Russ, Associate Botanist, Dale A. Zimmerman Herbarium. Vascular Plants of the Gila Wilderness
<http://www.wnmu.edu/academic/nspages2/gilaflorea/index.html>

(138) Van Devender, T.R. and R.K., Phelps, V., Thayer, D. and ASDM Docents, Paper - 15 April, 2 Oct., 23 Dec. 1986; 11 April 1987; Waterman Mountains: limestone ridges and lower slopes; 2400-2700 ft. elev.; T12S, R8E Sec. 32+33; 32D20'30-35"N; 111D 26-27' W.

(HR) Historical Record (possibly without author and/or observation date)

(TC) Tucson Citizen (Month Day, Year Section and Page Number)

(ADS) Arizona Daily Star (Month Day, Year Section and Page Number)

(AHS) Arizona Historical Society

(ANN) Anonymous

(JFW) John F. Wiens

(MBJ) Matthew B. Johnson, Program Manager and Curator of the Desert Legume Program - Boyce Thompson Southwestern Arboretum

(PCM) Personal Communication (Date)

(PDJ) Philip D. Jenkins, Assistant Curator of the University of Arizona Herbarium

(RGM) G. Meades

(TBL) Township Bird Listing

(WTK) William T. Kendall

(KGUN) Channel 9 (ABC - Month Day, Year & Program)

(KOLD) Channel 13 (CBS - Month Day, Year & Program)

(KVOA) Channel 4 (NBC - Month Day, Year & Program)

(MIX FM) 94.9 MIX fm (Month Day, Year & Program)